

**XOLEGEL- ketoconazole gel**  
**Almirall, LLC**

-----

**HIGHLIGHTS OF PRESCRIBING INFORMATION**

These highlights do not include all the information needed to use XOLEGEL Gel safely and effectively. See full prescribing information for XOLEGEL Gel.

**XOLEGEL® (ketoconazole) gel for topical use**  
**For Topical Use Only**  
**Initial U.S. Approval: 1981**

-----  
**INDICATIONS AND USAGE**  
-----

- XOLEGEL is an azole antifungal indicated for topical treatment of seborrheic dermatitis in immunocompetent adults and children 12 years of age and older. (1, 12.1)
- Safety and efficacy of XOLEGEL for treatment of fungal infections have not been established. (1)

-----  
**DOSAGE AND ADMINISTRATION**  
-----

- XOLEGEL is for topical use only, and not for oral, ophthalmic, or intravaginal use. (2)
- XOLEGEL should be applied once daily to the affected area for 2 weeks. (2)

-----  
**DOSAGE FORMS AND STRENGTHS**  
-----

- Gel containing 2% ketoconazole. (3)

-----  
**CONTRAINDICATIONS**  
-----

- None.

-----  
**WARNINGS AND PRECAUTIONS**  
-----

XOLEGEL is flammable. Avoid using near fire, flame, or smoking during and immediately following application of XOLEGEL. (5.1)

-----  
**ADVERSE REACTIONS**  
-----

- The most common treatment-related adverse reaction was application site burning (4%). (6.1)

**To report SUSPECTED ADVERSE REACTIONS, contact Almirall at 1-866-665-2782 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

**See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.**

**Revised: 12/2019**

---

**FULL PRESCRIBING INFORMATION: CONTENTS\***

**1 INDICATIONS AND USAGE**

**2 DOSAGE AND ADMINISTRATION**

**3 DOSAGE FORMS AND STRENGTHS**

**4 CONTRAINDICATIONS**

**5 WARNINGS AND PRECAUTIONS**

5.1 Flammable Contents

5.2 Systemic Effects

5.3 Local Effects

**6 ADVERSE REACTIONS**

6.1 Clinical Trials Experience

## **7 DRUG INTERACTIONS**

## **8 USE IN SPECIFIC POPULATIONS**

8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

## **11 DESCRIPTION**

## **12 CLINICAL PHARMACOLOGY**

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

## **13 NONCLINICAL TOXICOLOGY**

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

## **14 CLINICAL STUDIES**

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

16.1 How Supplied

16.2 Storage and Handling

## **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

XOLEGEL is indicated for the topical treatment of seborrheic dermatitis in immunocompetent adults and children 12 years of age and older.

Safety and efficacy of XOLEGEL for treatment of fungal infections have not been established.

### **2 DOSAGE AND ADMINISTRATION**

XOLEGEL is for topical use only, and not for oral, ophthalmic, or intravaginal use.

XOLEGEL should be applied once daily to the affected area for 2 weeks.

### **3 DOSAGE FORMS AND STRENGTHS**

XOLEGEL is a translucent to clear amber colored gel containing 2% ketoconazole.

### **4 CONTRAINDICATIONS**

None.

### **5 WARNINGS AND PRECAUTIONS**

## 5.1 Flammable Contents

XOLEGEL is flammable. Avoid being near fire, flame, or smoking during and immediately following application of XOLEGEL.

## 5.2 Systemic Effects

Hepatitis and, at high doses, lowered testosterone and ACTH induced corticosteroid serum levels have been seen with orally administered ketoconazole; these effects have not been seen with topically administered ketoconazole.

## 5.3 Local Effects

XOLEGEL can cause local irritation at the application site. If irritation occurs or if the disease worsens, use of the medication should be discontinued and the health care provider should be contacted [*see ADVERSE REACTIONS (6.1)*].

## 6 ADVERSE REACTIONS

### 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

In the 3 safety and efficacy trials, 65 of 933 subjects (7%) experienced at least one treatment-related adverse event. The most common treatment-related adverse reaction was application site burning (4%). Treatment-related application site reactions that were reported in < 1% of subjects were: dermatitis, discharge, dryness, erythema, irritation, pain, pruritus, and pustules. Other treatment-related adverse reactions that were reported in < 1% of subjects were: eye irritation, eye swelling, keratoconjunctivitis sicca, impetigo, pyogenic granuloma, dizziness, headache, paresthesia, acne, nail discoloration, facial swelling.

## 7 DRUG INTERACTIONS

Formal drug interaction studies with XOLEGEL have not been performed. Coadministration of oral ketoconazole with CYP3A4 metabolized HMG-CoA reductase inhibitors such as simvastatin, lovastatin and atorvastatin, may increase the risk of skeletal muscle toxicity, including rhabdomyolysis. These effects have not been observed with topically administered ketoconazole.

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

#### Risk Summary

There are no available data on XOLEGEL use in pregnant women to evaluate for a drug associated risk of major birth defects, miscarriage, or adverse maternal or fetal outcomes.

In animal reproduction studies with pregnant rats, structural abnormalities (syndactylia and oligodactylia) were observed following oral doses of ketoconazole during organogenesis (see *Data*). The available data do not allow calculation of relevant comparisons between the systemic exposure of ketoconazole observed in animal studies to the systemic exposure observed in humans after topical use of XOLEGEL.

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2%-4% and 15%-20%, respectively.

## Data

### *Animal Data*

Oral administration of ketoconazole at 80 mg/kg/day to pregnant rats during organogenesis was associated with structural abnormalities (syndactylia and oligodactylia) . However, these effects may be related to maternal toxicity, which was also seen at this and higher dose levels.

In oral peri- and postnatal development studies in rats, maternal toxicity, prolonged gestation, embryoletality and fetotoxicity were observed at ketoconazole doses of 40 mg/kg/day and higher.

## **8.2 Lactation**

### Risk Summary

There are no data available on the presence of ketoconazole in human milk, its effects on the breastfed infant, or its effects on milk production. After topical application, ketoconazole concentrations in plasma are low and therefore concentrations in human breast milk are likely to be low [see *Clinical Pharmacology (12.3)*].

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for XOLEGEL and any potential adverse effects on the breastfed infant from XOLEGEL or from the underlying maternal condition.

### Clinical Considerations

Advise breastfeeding women not to apply XOLEGEL directly to the nipple and areola to avoid direct infant exposure.

## **8.4 Pediatric Use**

Safety and effectiveness in pediatric subjects below the age of 12 have not been established.

## **8.5 Geriatric Use**


Of the 933 subjects in the three safety and efficacy trials, 193 (20.7%) were 65 and older, while 61 (6.5%) were 75 and older. No overall differences in safety or effectiveness were observed between these subjects and younger subjects but greater sensitivity of some older individuals cannot be ruled out.

## **11 DESCRIPTION**

XOLEGEL contains the antifungal agent ketoconazole USP at 2% in a topical anhydrous gel vehicle for topical administration.

Chemically, ketoconazole is ( $\pm$ )-cis-1-Acetyl-4-[p-[[2-(2,4-dichlorophenyl)-2-(1H-imidazol-1-ylmethyl)-1,3-dioxolan-4-yl]methoxy]phenyl]piperazine, with the molecular formula  $C_{26}H_{28}Cl_2N_4O_4$  and a molecular weight of 531.43.

**Figure 1**



Each gram contains: 20 mg ketoconazole USP, dehydrated alcohol (34%), ascorbic acid, butylated hydroxytoluene, citric acid monohydrate, glycerin, hydroxypropyl cellulose, polyethylene glycol 400, PPG-15 stearyl ether, propylene glycol, FD&C yellow No. 6, and FD&C yellow No. 10.

XOLEGEL is a smooth, translucent to clear, amber gel.

## **12 CLINICAL PHARMACOLOGY**

### **12.1 Mechanism of Action**

The mechanism of action of ketoconazole in the treatment of seborrheic dermatitis is unknown.

### **12.2 Pharmacodynamics**

Pharmacodynamic markers for seborrheic dermatitis have not been identified.

### **12.3 Pharmacokinetics**

In a pharmacokinetic absorption trial, eighteen subjects, both males and females, with severe seborrheic dermatitis (range 1-14% of body surface area) applied XOLEGEL once daily for 2 weeks. The median total amount of gel applied was 4.6 g (range 1.65–46.3 g). Daily doses ranged from 0.05 to 3.47 g. Mean ( $\pm$  standard deviation [SD]) peak plasma levels were 1.35 ( $\pm$  3.18) ng/mL on Day 7 (range from <0.1 ng/mL, to 13.9 ng/mL), and 0.80 ( $\pm$  1.22) ng/mL on Day 14 (range from <0.1 ng/mL to 5.4 ng/mL). Median  $T_{max}$  was 8 hours on Day 7 and 7 hours on Day 14. Mean ( $\pm$  SD)  $AUC_{0-24}$  values were 20.8 ( $\pm$  44.7) ng•h/mL and 15.6 ( $\pm$  26.4) ng•h/mL on Day 7 and 14, respectively.

The plasma levels from an oral dose of 200 mg ketoconazole taken with a meal are approximately 250 times higher than the resulting plasma levels of ketoconazole

following topical application of XOLEGEL.

## 13 NONCLINICAL TOXICOLOGY

### 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

A 2-year dermal carcinogenicity study conducted in mice with topical administration of ketoconazole gel at doses up to 80 mg ketoconazole/kg/day exhibited no evidence of carcinogenic activity. A long-term feeding study in mice and in rats showed no evidence of carcinogenic activity.

Ketoconazole produced no evidence of mutagenicity in the dominant lethal mutation test in male and female mice at single oral doses up to 80 mg/kg. When tested in the Ames assay, ketoconazole was not mutagenic to *Salmonella typhimurium* in the presence or absence of metabolic activation. Ketoconazole, in combination with another drug, gave equivocal results in the mouse micronucleus test.

At oral doses of 75 mg/kg/day, ketoconazole impaired the reproductive performance in female (decreased pregnancy and implantation rates) and male (increased abnormal sperm and decreased sperm motility) rats.

## 14 CLINICAL STUDIES

Study 1 was a multicenter, double-blind, randomized, vehicle-controlled trial which enrolled 459 subjects 12 years of age and older with moderate to severe seborrheic dermatitis. A total of 229 subjects were treated with XOLEGEL, and 230 subjects were treated with vehicle. All subjects were treated once daily for 14 days, and efficacy was assessed at Day 28 (i.e., 2 weeks after end of treatment). Effective Treatment was defined as:

- an Investigator's Global Assessment score of  $\leq 1$  (completely clear or almost clear) and
- erythema and scaling scores of 0 (none) if the baseline score was 2, or 1 (mild) if the baseline score was 3.

The proportion of subjects effectively treated is shown in Table 1.

**Table 1: Trial Results**

	<i>XOLEGEL</i> N=229	<i>Vehicle</i> N=230
Number and proportion of subjects effectively treated	58 (25.3%)	32 (13.9%)

Two additional double-blind, randomized, vehicle-controlled, parallel, and multicenter trials that included a total of 316 subjects treated with XOLEGEL provided supportive evidence of the efficacy of XOLEGEL for treatment of seborrheic dermatitis. Subjects applied either XOLEGEL or vehicle study treatment to the affected area(s) once daily for 14 days and were followed through Day 28. Efficacy was assessed by the proportion of subjects who were completely clear at Day 28.

The contribution to efficacy of individual components of the vehicle has not been established.

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

### **16.1 How Supplied**

XOLEGEL<sup>®</sup> (ketoconazole) Gel, 2% is supplied in 45-gram (NDC 16110-080-45) white-coated aluminum tubes with white caps, and is dispensed with FDA-Approved Patient Labeling.

### **16.2 Storage and Handling**

Store at 25°C (77°F); excursions permitted to 15° - 30°C (59° - 86°F).

Contents are flammable.

Keep out of reach of children.

## **17 PATIENT COUNSELING INFORMATION**

*[See FDA-Approved Patient Labeling (Patient Information)]*

- This medication is to be used as directed by the health care provider. It is for external use only.
- XOLEGEL may be irritating to mucus membranes. Contact with the eyes, nostrils, and mouth should be avoided.
- As with any topical medication, patients should wash their hands after application.
- Advise breastfeeding women not to apply XOLEGEL directly to the nipple and areola to avoid direct infant exposure.
- This medication should not be used for any disorder other than that for which it has been prescribed.
- Patients should report any signs of adverse reactions to their health care provider.

## **PATIENT INFORMATION**

### **XOLEGEL<sup>®</sup> (Xol-a-gel)**

### **(ketoconazole) Gel, 2%**

Read the Patient Information that comes with XOLEGEL carefully before you start using it and each time you get a refill. There may be new information. This leaflet does not take the place of talking with your health care provider. If you have any questions about XOLEGEL, ask your health care provider.

### **What is XOLEGEL?**

XOLEGEL is a prescription medicine used on the skin to treat a skin condition called seborrheic dermatitis.

Patients with seborrheic dermatitis can have areas of dry, flaky skin on the scalp, face, ears, chest, or upper back. XOLEGEL is only to be used in adults and in children older

than 12 years of age who have a normal (healthy) immune system. XOLEGEL has not been studied in children below the age of 12.

It is not known whether XOLEGEL can be used to treat fungal infections.

XOLEGEL is a translucent to clear, amber colored gel.

### **What should I tell my health care provider before using XOLEGEL?**

- Tell your health care provider about all of your medical conditions, including if you are pregnant or planning to become pregnant, or are breastfeeding or planning to breastfeed. XOLEGEL should be used during pregnancy and breastfeeding only if needed.
- Tell your health care provider about all of the medicines you take, including prescription and non-prescription medicines, vitamins, and herbal supplements. Keep a list of your medicines and show it to your health care provider and pharmacist. Tell your health care provider and pharmacist when you get a new medicine. It is not known if XOLEGEL and other medicines can interact with each other.

### **How should I use XOLEGEL?**

- Use XOLEGEL exactly as prescribed. Talk to your health care provider if your condition gets worse or does not get better by the end of your treatment.
- Wash your hands before and after applying XOLEGEL.
- Spread a thin layer of XOLEGEL evenly on the affected skin with your fingertips. Be sure to cover all affected areas.
- Do not wash the areas where you applied XOLEGEL for at least 3 hours after you apply it.
- Wait at least 20 minutes after you spread XOLEGEL on your skin before you put makeup or sunscreens on the affected areas.
- Use XOLEGEL once daily for 2 weeks.

### **What should I avoid while using XOLEGEL?**

- XOLEGEL is only to be used on the skin. It is not for eye, mouth, or vaginal use.
- Do not touch your eyes, nose, or mouth while you are applying XOLEGEL. Wash your hands well after you apply it. Irritation may occur if you get XOLEGEL in your eyes, nose, or mouth.
- If used during breastfeeding and XOLEGEL is applied on the chest, take care to avoid accidental ingestion of XOLEGEL by the baby.
- XOLEGEL is flammable (it can catch fire). Stay away from heat, flame, or smoking while you are applying XOLEGEL and right after you apply it.
- This medication should not be used for any disorder other than that for which it has been prescribed.

### **What are the possible side effects of XOLEGEL?**

- The effects of XOLEGEL during pregnancy, including whether XOLEGEL can harm your unborn baby, are not known.
- It is not known if XOLEGEL can pass into your breastmilk or if it can harm your breastfed baby.
- Stop using XOLEGEL and talk to your health care provider if you develop itching, a rash, or any skin irritation after using XOLEGEL.
- Stop using XOLEGEL and talk to your health care provider if your skin condition (seborrheic dermatitis) gets worse.



- The most common side effect is a burning feeling where XOLEGEL is applied.
- Report any side effects to your health care provider to receive immediate medical attention. You can also report suspected side effects by calling the US Food and Drug Administration at 1-800-FDA-1088, or reporting via the internet at [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

These are not all of the side effects of XOLEGEL. For more information, ask your health care provider or pharmacist.

### **How should I store XOLEGEL?**

- Store XOLEGEL at 59°F to 86°F (15° to 30°C).
- Keep XOLEGEL and all medicines out of the reach of children.
- Contents are flammable. Avoid storing XOLEGEL near heat or flame.

### **General information about XOLEGEL**

Medicines are sometimes prescribed for conditions that are not mentioned in Patient Information leaflets. Do not use XOLEGEL for a condition for which it was not prescribed. Do not give XOLEGEL to other people, even if they have the same symptoms that you have. It may harm them.

This Patient Information leaflet summarizes the most important information about XOLEGEL. If you would like more information, talk with your health care provider. You can also ask your pharmacist or health care provider for information about XOLEGEL that is written for health professionals.

### **What are the ingredients in XOLEGEL?**

**Active ingredient:** ketoconazole, USP

**Inactive ingredients:** dehydrated alcohol, ascorbic acid, butylated hydroxytoluene, citric acid monohydrate, glycerin, hydroxypropyl cellulose, polyethylene glycol 400, PPG-15 stearyl ether, propylene glycol, FD&C yellow No. 6, and FD&C Yellow No. 10.

This Patient Information leaflet has been approved by the U.S. Food and Drug Administration.

The Patient Information leaflet was last revised: May 2012.

Manufactured for Almirall, LLC  
101 Lindenwood Drive, Suite 400, Malvern, PA 19355

Revised 12/2019  
XOLEGEL is a registered trademark of Almirall, LLC.

© 2019 Almirall, LLC

Almirall, LLC

**PRINCIPAL DISPLAY PANEL - 45 g Carton**

**NDC 16110-080-45**

**Xolegel®**

(ketoconazole) Gel, 2%

**AQUA**

PHARMACEUTICALS

**45 grams**

**Rx only. For topical use only.**



**PRINCIPAL DISPLAY PANEL - 45 g Carton**

**NDC 16110-080-45**

**Xolegel®**

(ketoconazole) Gel, 2%

**AQUA**

PHARMACEUTICALS

**45 grams**

**Rx only. For topical use only.**

**Usual Dosage:** See package insert.  
Store at 25°C (77°F); excursions permitted to 15°C – 30°C (59°F – 86°F).

**Caution:** Not for oral, ophthalmic or intravaginal use. Keep out of reach of children. If seal is damaged or punctured, do not use, and return product to place of purchase. **AVOID FIRE, FLAME, OR SMOKING DURING AND IMMEDIATELY FOLLOWING APPLICATION.**

Manufactured by DPT Laboratories, San Antonio, TX 78215  
For Aqua Pharmaceuticals, an Almirall Company, Exton, PA 19341  
XOLEGEL is a registered trademark of Aqua Pharmaceuticals.  
© 2016 Aqua Pharmaceuticals  
106677

**AQUA**  
PHARMACEUTICALS  
an Almirall company



45 grams NDC 16110-080-45

# Xolegel<sup>®</sup>

(ketoconazole) Gel, 2%

**Rx only**  
For topical use only

## XOLEGEL

ketoconazole gel

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:16110-080
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>KETOCONAZOLE</b> (UNII: R9400W927I) (KETOCONAZOLE - UNII:R9400W927I)	KETOCONAZOLE	20 mg in 1 g

### Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
ASCORBIC ACID (UNII: PQ6CK8PD0R)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
GLYCERIN (UNII: PDC6A3C0OX)	
HYDROXYPROPYL CELLULOSE (1600000 WAMW) (UNII: RFW2ET671P)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
PPG-15 STEARYL ETHER (UNII: 1II18XLS1L)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:16110-080-45	1 in 1 CARTON	07/28/2006	12/30/2022
1		45 g in 1 TUBE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA021946	07/28/2006	12/30/2022

**Labeler** - Almirall, LLC (605425912)

### Establishment

Name	Address	ID/FEI	Business Operations
DPT Laboratories, Ltd.		832224526	ANALYSIS(16110-080) , LABEL(16110-080) , MANUFACTURE(16110-080)

Revised: 1/2023

Almirall, LLC