

DRUG FACTS

Active ingredient(s)

Purpose

Use(s)

temporarily relieves minor aches and pains due to:

- headache
- a cold
- toothache
- muscular aches
- arthritis
- menstrual cramps

Warnings

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- with any other products containing acetaminophen. Taking more than directed may cause liver damage.

Ask a doctor before use if you have

- asthma
- ulcers
- bleeding problems
- stomach problems such as heartburn, upset stomach, or stomach pain that do not go away or return

Ask a doctor or pharmacist before use if you are

Ask a doctor or pharmacist before use if you are taking a prescription drug for:

- anticoagulation (thinning of the blood)
- diabetes
- gout
- arthritis

When using this product

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- new symptoms occur
- symptoms do not get better or worsen
- ringing in the ears or loss of hearing occurs
- painful area is red or swollen
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

If pregnant or breast-feeding

Keep out of reach of children

Keep out of reach of children.

Directions

- do not use more than directed (see overdose warning)
- drink a full glass of water with each dose
- adults and children 12 years of age and over: take 2 tablets every 6 hours; not more than 8 tablets in 24 hours, or as directed by a doctor
- children under 12 years: ask a doctor

Other information

- read all product information before using.
- store at 20°–25°C (68°–77°F). Protect from moisture.

Drug Facts

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen and aspirin or other pain relievers/fever reducers. Acetaminophen and aspirin may cause liver damage and stomach bleeding.

The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally, rapid heart beat.

Distributed by: **INSIGHT Pharmaceuticals Corp.**, Langhorne, PA 19047-1749

Inactive ingredients

Colloidal Silicon Dioxide, Crosppovidone, Lecithin, Microcrystalline Cellulose, Polyvinyl Alcohol, Povidone, Pregelatinized Starch, Sodium Starch Glycolate, Starch, Stearic Acid, Talc, Titanium Dioxide