

## DRUG FACTS

### Active ingredient(s)

Purpose

### Use(s)

- temporarily relieves minor aches and pains due to:
- headache
- a cold
- arthritis
- muscular aches
- sinusitis
- toothache
- premenstrual and menstrual cramps

### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 8 tablets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

### Do not use

- if you ever had an allergic reaction to aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

### Ask a doctor before use if you have

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have liver disease
- you have high blood pressure, heart disease, liver disease or kidney disease
- you have asthma
- you are taking a diuretic

### Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug or are under a doctor's care for any serious condition

### When using this product

#### Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- have bloody or black stools
- vomit blood
- have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- any new symptoms appear
- ringing in the ears or loss of hearing occurs
- pain gets worse or lasts more than 10 days
- painful area is red or swollen
- fever gets worse or lasts more than 3 days

### If pregnant or breast-feeding

#### Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

- do not use more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over
- take 2 tablets every 6 hours
- do not take more than 8 tablets in 24 hours
- children under 12 years: do not use unless directed by a doctor

### Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

### Inactive ingredients

corn starch, crospovidone, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, sodium lauryl sulfate, sodium starch glycolate, stearic acid, talc, titanium dioxide