


PENTOXIFYLLINE- pentoxifylline tablet, extended release
Preferred Pharmaceuticals, Inc.

PENTOXIFYLLINE EXTENDED-RELEASE TABLETS, USP 400 mg

DESCRIPTION

Pentoxifylline Extended-Release Tablets, USP for oral administration contain 400 mg of the active drug and the following inactive ingredients: colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, magnesium stearate, polyethylene glycol, and titanium dioxide in an extended-release formulation. Pentoxifylline is a tri-substituted xanthine derivative designated chemically as 3,7-Dihydro-3,7-dimethyl-1-(5-oxohexyl)-1H-purine-2,6-dione that, unlike theophylline, is a hemorrheologic agent, i.e., an agent that affects blood viscosity. Pentoxifylline is soluble in water and ethanol, and sparingly soluble in toluene. The molecular formula is $C_{13}H_{18}N_4O_3$ and its molecular weight is 278.31.

The structural formula is:



USP Dissolution Test 9

CLINICAL PHARMACOLOGY

Mode of Action

Pentoxifylline and its metabolites improve the flow properties of blood by decreasing its viscosity. In patients with chronic peripheral arterial disease, this increases blood flow to the affected microcirculation and enhances tissue oxygenation. The precise mode of action of pentoxifylline and the sequence of events leading to clinical improvement are still to be defined. Pentoxifylline administration has been shown to produce dose related hemorrheologic effects, lowering blood viscosity, and improving erythrocyte flexibility. Leukocyte properties of hemorrheologic importance have been modified in animal and in vitro human studies. Pentoxifylline has been shown to increase leukocyte deformability and to inhibit neutrophil adhesion and activation. Tissue oxygen levels have been shown to be significantly increased by therapeutic doses of pentoxifylline in patients with peripheral arterial disease.

Pharmacokinetics and Metabolism

After oral administration in aqueous solution pentoxifylline is almost completely absorbed. It undergoes a first-pass effect and the various metabolites appear in plasma very soon after dosing. Peak plasma levels of the parent compound and its metabolites are reached within 1 hour. The major metabolites are Metabolite I (1-[5-hydroxyhexyl]-3,7-dimethylxanthine) and Metabolite V (1-[3-carboxypropyl]-3,7-dimethylxanthine), and plasma levels of these metabolites are 5 and 8 times greater, respectively, than pentoxifylline.

Following oral administration of aqueous solutions containing 100 to 400 mg of pentoxifylline, the pharmacokinetics of the parent compound and Metabolite I are dose-related and not proportional (non-

linear), with half-life and area under the blood level time curve (AUC) increasing with dose. The elimination kinetics of Metabolite V are not dose-dependent. The apparent plasma half-life of pentoxifylline varies from 0.4 to 0.8 hours and the apparent plasma half-lives of its metabolites vary from 1 to 1.6 hours. There is no evidence of accumulation or enzyme induction (Cytochrome P450) following multiple oral doses.

Excretion is almost totally urinary; the main biotransformation product is Metabolite V. Essentially no parent drug is found in the urine. Despite large variations in plasma levels of parent compound and its metabolites, the urinary recovery of Metabolite V is consistent and shows dose proportionality. Less than 4% of the administered dose is recovered in feces. Food intake shortly before dosing delays absorption of an immediate-release dosage form but does not affect total absorption. The pharmacokinetics and metabolism of pentoxifylline have not been studied in patients with renal and/or hepatic dysfunction, but AUC was increased and elimination rate decreased in an older population (60-68 years) compared to younger individuals (22-30 years).

After administration of the 400 mg extended-release pentoxifylline tablet, plasma levels of the parent compound and its metabolites reach their maximum within 2 to 4 hours and remain constant over an extended period of time. Coadministration of pentoxifylline tablets with meals resulted in an increase in mean C_{max} and AUC by about 28% and 13% for pentoxifylline, respectively. C_{max} for metabolite M1 also increased by about 20%. The extended-release of pentoxifylline from the tablet eliminates peaks and troughs in plasma levels for improved gastrointestinal tolerance.

INDICATIONS AND USAGE

Pentoxifylline Extended-Release Tablets, USP are indicated for the treatment of patients with intermittent claudication on the basis of chronic occlusive arterial disease of the limbs. Pentoxifylline can improve function and symptoms but is not intended to replace more definitive therapy, such as surgical bypass, or removal of arterial obstructions when treating peripheral vascular disease.

CONTRAINDICATIONS

Pentoxifylline should not be used in patients with recent cerebral and/or retinal hemorrhage or in patients who have previously exhibited intolerance to this product or methylxanthines such as caffeine, theophylline, and theobromine.

PRECAUTIONS

General

Patients with chronic occlusive arterial disease of the limbs frequently show other manifestations of arteriosclerotic disease. Pentoxifylline has been used safely for treatment of peripheral arterial disease in patients with concurrent coronary artery and cerebrovascular diseases, but there have been occasional reports of angina, hypotension, and arrhythmia. Controlled trials do not show that pentoxifylline causes such adverse effects more often than placebo, but, as it is a methylxanthine derivative, it is possible some individuals will experience such responses. Patients on warfarin should have more frequent monitoring of prothrombin times, while patients with other risk factors complicated by hemorrhage (e.g., recent surgery, peptic ulceration, cerebral and/or retinal bleeding) should have periodic examinations for bleeding including, hematocrit and/or hemoglobin.

Drug Interactions

Although a causal relationship has not been established, there have been reports of bleeding and/or prolonged prothrombin time in patients treated with pentoxifylline with and without anticoagulants or platelet aggregation inhibitors. Patients on warfarin should have more frequent monitoring of prothrombin times, while patients with other risk factors complicated by hemorrhage (e.g., recent

surgery, peptic ulceration) should have periodic examinations for bleeding including hematocrit and/or hemoglobin. Concomitant administration of pentoxifylline and theophylline-containing drugs leads to increased theophylline levels and theophylline toxicity in some individuals. Such patients should be closely monitored for signs of toxicity and have their theophylline dosage adjusted as necessary. Pentoxifylline has been used concurrently with antihypertensive drugs, beta blockers, digitalis, diuretics, antidiabetic agents, and antiarrhythmics, without observed problems. Small decreases in blood pressure have been observed in some patients treated with pentoxifylline; periodic systemic blood pressure monitoring is recommended for patients receiving concomitant antihypertensive therapy. If indicated, dosage of the antihypertensive agents should be reduced.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term studies of the carcinogenic potential of pentoxifylline were conducted in mice and rats by dietary administration of the drug at doses up to 450 mg/kg (approximately 19 times the maximum recommended human daily dose (MRHD) in both species when based on body weight; 1.5 MRHD in the mouse and 3.3 times the MRHD in the rat when based on body surface area). In mice, the drug was administered for 18 months, whereas in rats, the drug was administered for 18 months followed by an additional 6 months without drug exposure. In the rat study, there was a statistically significant increase in benign mammary fibroadenomas in females of the 450 mg/kg group. The relevance of this finding to human use is uncertain. Pentoxifylline was devoid of mutagenic activity in various strains of *Salmonella* (Ames test) and in cultured mammalian cells (unscheduled DNA synthesis test) when tested in the presence and absence of metabolic activation. It was also negative in the *in vivo* mouse micronucleus test.

Pregnancy

Category C.

Teratogenicity studies have been performed in rats and rabbits using oral doses up to 576 and 264 mg/kg, respectively. On a weight basis, these doses are 24 and 11 times the maximum recommended human daily dose (MRHD); on a body-surface-area basis, they are 4.2 and 3.5 times the MRHD. No evidence of fetal malformation was observed. Increased resorption was seen in rats of the 576 mg/kg group. There are no adequate and well controlled studies in pregnant women. Pentoxifylline should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Nursing Mothers

Pentoxifylline and its metabolites are excreted in human milk. Because of the potential for tumorigenicity shown for pentoxifylline in rats, a decision should be made whether to discontinue nursing or discontinue the drug, taking into account the importance of the drug to the mother.

Pediatric Use

Safety and effectiveness in pediatric patients have not been established.

ADVERSE REACTIONS

Clinical trials were conducted using either extended-release pentoxifylline tablets for up to 60 weeks or immediate-release pentoxifylline capsules for up to 24 weeks. Dosage ranges in the tablet studies were 400 mg bid to tid and in the capsule studies, 200 to 400 mg tid. The table summarizes the incidence (in percent) of adverse reactions considered drug related, as well as the numbers of patients who received extended-release pentoxifylline tablets, immediate-release pentoxifylline capsules, or the corresponding placebos. The incidence of adverse reactions was higher in the capsule studies (where dose related increases were seen in digestive and nervous system side effects) than in the tablet studies. Studies with the capsule include domestic experience, whereas studies with the extended-release tablets were conducted outside the U.S.

The table indicates that in the tablet studies few patients discontinued because of adverse effects.

INCIDENCE (%) OF SIDE EFFECTS

	Extended-Release		Immediate	
	Tablets USP		Release Capsules	
	Commercially Available		Used Only For Controlled Clinical Trials	
	Pentoxifylline	Placebo	Pentoxifylline	Placebo
(Number of Patients at Risk)	(321)	(128)	(177)	(138)
Discontinued for Side Effect	3.21	0	9.6	7.2
CARDIOVASCULAR SYSTEM				
Angina/Chest Pain	0.3	-	1.1	2.2
Arrhythmia/Palpitation	-	-	1.7	0.7
Flushing	-	-	2.3	0.7
DIGESTIVE SYSTEM				
Abdominal Discomfort	-	-	4.0	1.4
Belching/Flatus/Bloating	0.6	-	9.0	3.6
Diarrhea	-	-	3.4	2.9
Dyspepsia	2.8	4.7	9.6	2.9
Nausea	2.2	0.8	28.8	8.7
Vomiting	1.2	-	4.5	0.7
NERVOUS SYSTEM				
Agitation/Nervousness	-	-	1.7	0.7
Dizziness	1.9	3.1	11.9	4.3
Drowsiness	-	-	1.1	5.8
Headache	1.2	1.6	6.2	5.8
Insomnia	-	-	2.3	2.2
Tremor	0.3	0.8	-	-
Blurred Vision	-	-	2.3	1.4

Pentoxifylline has been marketed in Europe and elsewhere since 1972. In addition to the above symptoms, the following have been reported spontaneously since marketing or occurred in other clinical trials with an incidence of less than 1%; the causal relationship was uncertain:

Cardiovascular - dyspnea, edema, hypotension.

Digestive - anorexia, cholecystitis, constipation, dry mouth/thirst.

Nervous - anxiety, confusion, depression, seizures, aseptic meningitis.

Respiratory - epistaxis, flu-like symptoms, laryngitis, nasal congestion.

Skin and Appendages - brittle fingernails, pruritus, rash, urticaria, angioedema.

Special Senses - blurred vision, conjunctivitis, earache, scotoma.

Miscellaneous - bad taste, excessive salivation, leukopenia, malaise, sore throat/swollen neck glands, weight change.

A few rare events have been reported spontaneously worldwide since marketing in 1972. Although they occurred under circumstances in which a causal relationship with pentoxifylline could not be established, they are listed to serve as information for physicians. Cardiovascular - angina, arrhythmia,

tachycardia, anaphylactoid reactions. Digestive - hepatitis, jaundice, increased liver enzymes; and Hemic and Lymphatic - decreased serum fibrinogen, pancytopenia, aplastic anemia, leukemia, purpura, thrombocytopenia.

OVERDOSAGE

Overdosage with pentoxifylline has been reported in pediatric patients and adults. Symptoms appear to be dose related. A report from a poison control center on 44 patients taking overdoses of enteric-coated pentoxifylline tablets noted that symptoms usually occurred 4-5 hours after ingestion and lasted about 12 hours. The highest amount ingested was 80 mg/kg; flushing, hypotension, convulsions, somnolence, loss of consciousness, fever, and agitation occurred. All patients recovered. In addition to symptomatic treatment and gastric lavage, special attention must be given to supporting respiration, maintaining systemic blood pressure, and controlling convulsions. Activated charcoal has been used to absorb pentoxifylline in patients who have overdosed.

DOSAGE AND ADMINISTRATION

The usual dosage of pentoxifylline in extended-release tablet form is one tablet (400 mg) three times a day with meals.

While the effect of pentoxifylline may be seen within 2 to 4 weeks, it is recommended that treatment be continued for at least 8 weeks. Efficacy has been demonstrated in double-blind clinical studies of 6 months duration.

Digestive and central nervous system side effects are dose related. If patients develop these effects it is recommended that the dosage be lowered to one tablet twice a day (800 mg/day). If side effects persist at this lower dosage, the administration of pentoxifylline should be discontinued.

HOW SUPPLIED

Pentoxifylline Extended-Release Tablets, USP are available for oral administration as 400 mg white, oval, unscored, film coated tablets, imprinted "APO 033" on one side and plain on the other side.

Bottle of 30 - 68788-9742-3

Store at 20° to 25°C [see Controlled Room Temperature].

Dispense in a tight, light-resistant container.

APOTEX INC.

PENTOXIFYLLINE EXTENDED-RELEASE TABLETS, USP

400mg

Manufactured by: Manufactured for:
Apotex Inc. Apotex Corp.
Toronto, Ontario Weston, Florida
Canada M9L 1T9 33326

Relabeled by:

Preferred Pharmaceuticals, Inc.

9/2012

PRINCIPAL DISPLAY PANEL

Representative sample of labeling (see **HOW SUPPLIED** section for complete listing):

Pentoxifylline Extended-Release Tablets USP

400 mg

Rx

**Pentoxifylline
Extended-Release
Tablets 400mg**
Generic for Trental
Each tablet contains 400mg pentoxifylline

Pkg Size: Exp Date:
Lot#: Batch#:
Ins:
Mfg: Apotex Inc.; Toronto, Ontario Canada
Prod#:

Warning
Store at 20° to 25°C (68° to 77°F). See USP Controlled Room Temperature. Take with meals. Keep this and all medication out of the reach of children. Rx Only. Tablet is capsule-shaped, white, imprinted with APO 033



CAUTION: Federal law PROHIBITS transfer of this drug to any person other than the patient for whom it was prescribed.

Pentoxifylline Extended-Release Tablets 400mg
Qty: Ins:
Lot#: Bat#:
Prod# (NDC):

Pentoxifylline Extended-Release Tablets 400mg
Qty: Ins:
Lot#: Bat#:
Prod# (NDC):

Pentoxifylline Extended-Release Tablets 400mg
Qty: Ins:
Insurance NDC:
Lot#: Bat#:

Pentoxifylline Extended-Release Tablets 400mg
Qty: Ins:
Lot#: Bat#:
Prod# (NDC):



Directions English
Use as directed by your doctor
Take ___ tablet(s) every ___ hours.

Instrucciones Espanol:
Usó según lo dirigido por su doctor
Toma cada ___ tableta(s) cada ___ horas.

Log

Chart

Billing

Patient

PENTOXIFYLLINE

pentoxifylline tablet, extended release

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:68788-9742(NDC:60505-0033)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PENTOXIFYLLINE (UNII: SD6QCT3TSU) (PENTOXIFYLLINE - UNII:SD6QCT3TSU)	PENTOXIFYLLINE	400 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYDROXYPROPYL CELLULOSE (UNII: RFW2ET671P)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYETHYLENE GLYCOL 8000 (UNII: Q662QK8M3B)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	WHITE	Score	no score
Shape	OVAL	Size	16mm

Flavor		Imprint Code	APO;033	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68788-9742-3	30 in 1 BOTTLE		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA075191	09/28/2012		

Labeler - Preferred Pharmaceuticals, Inc. (791119022)

Establishment

Name	Address	ID/FEI	Business Operations
Preferred Pharmaceuticals, Inc.		791119022	REPACK(68788-9742)

Revised: 9/2012

Preferred Pharmaceuticals, Inc.