DRUG FACTS

Active ingredient(s) ***Purpose***

Use(s)

- temporarily relieves
- headache
- toothache
- pain and fever of colds
- muscle pain
- menstrual pain
- minor pain of arthritis

Warnings

Do not use

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if you have

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

When using this product

Stop use and ask a doctor if

- an allergic reaction occurs (seek medical help right away)
- pain gets worse or lasts more than 10 days
- redness or swelling is present
- fever lasts more than 3 days
- new symptoms occur
- ringing in the ears or a loss of hearing occurs
- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better

If pregnant or breast-feeding

Keep out of reach of children

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

adults and children 12 years and over

- see instructions above for opening packet
- place 1-2 powders on the tongue and swallow with or without water
- repeat every 4-6 hours, while symptoms persist
- do not take more than 8 powders in 24 hours
- children under 12 years: Ask a doctor

Other information

Drug Facts

Children and teenagers who have or are recovering from chickenpox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for longer than directed

Do not use if pouch is torn or open

Critical Care Aspirin to Go
distributed by
UrgentRx®
Breakthrough Products, Inc.
Denver, CO 80202

Inactive ingredients

acesulfame potassium, citric acid, flavor, glucose, sucralose