DRUG FACTS

Active ingredient(s)	*Purpose*

Use(s)
- temporarily relieves
- muscle pain
- minor arthritis pain
- headache
- pain and fever of colds

Warnings

Do not use
Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if you have
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are
Ask a doctor or pharmacist before use if you are taking a prescription drug for
- gout
- diabetes
- arthritis

When using this product
When using this product limit use of caffeine containing drugs, foods, or drinks, because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally rapid heartbeat

Stop use and ask a doctor if
- an allergic reaction occurs (seek medical help right away)
- pain gets worse or lasts more than 10 days
- redness or swelling is present
- fever lasts more than 3 days
- new symptoms occur
- ringing in the ears or a loss of hearing occurs
- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better

If pregnant or breast-feeding

Keep out of reach of children
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
adults and children 12 years and over
- see instructions above for opening packet
- place 1 powder on the tongue and swallow with or without water
- repeat every 4-6 hours, while symptoms persist
- do not take more than 4 powders in 24 hours
- children under 12 years: Ask a doctor

Other information
Drug Facts
Children and teenagers who have or are recovering from chickenpox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Aspirin may cause a severe allergic reaction which may include:
- hives
- facial swelling
- asthma (wheezing)
- shock

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for longer than directed

Do not use if pouch is torn or open

Ache & Pain Relief to Go
distributed by
UrgentRx®
Breakthrough Products, Inc.
Denver, CO 80202

Inactive ingredients
acesulfame potassium, citric acid, flavor, glucose, sucralose