DRUG FACTS

Active ingredient(s) ***Purpose***

Use(s)
temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold

Warnings

Do not use
you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
- a cough that occurs with too much phlegm (mucus)
- a cough that lasts or is chronic as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

When using this product

Stop use and ask a doctor if
cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or presistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding

Keep out of reach of children
In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
- do not take more than 8 liquidgels in any 24-hour period
- **adults and children 12 years and over:** take 2 liquidgels every 6 to 8 hours, as needed
- **children under 12 years**: ask a doctor

Other information
- **TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**
- store at 20º-25ºC (68º-77ºF)
- avoid excessive heat above 40ºC (104ºF)
- protect from light
- use by expiration date on package

Inactive ingredients
FD&C blue #1, FD&C red #40, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol, white edible ink