

ACID CONTROLLER- famotidine tablet
Walgreen Company

Walgreen Co. Acid Controller Drug Facts

Active ingredient (in each tablet)

Famotidine 20 mg

Purpose

Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- if you have kidney disease, except under the advice and supervision of a doctor
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating, or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 12 years and over:
- to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture and light

Inactive ingredients

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, lactose (monohydrate), magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc, and titanium dioxide

Questions or comments?

1-800-719-9260

Principal Display Panel

MAXIMUM STRENGTH

ACID CONTROLLER

Famotidine Tablets, 20 mg / Acid Reducer

Just one tablet prevents & relieves heartburn due to acid indigestion

Actual Size

Compare to Maximum Strength Pepcid® AC active ingredient



DO NOT USE IF PRINTED FOIL UNDER CAP IS BROKEN OR MISSING

Well at Walgreens
MAXIMUM STRENGTH
Acid Controller
Famotidine Tablets, 20 mg
Acid Reducer
Just one tablet prevents & relieves heartburn due to acid indigestion
85 TABLETS

Active ingredient: Famotidine 20 mg (in each tablet)
Purpose: Acid reducer



WALGREENS PHARMACIST RECOMMENDED*

MAXIMUM STRENGTH

Acid Controller

Famotidine Tablets, 20 mg / Acid Reducer

- Just one tablet prevents & relieves heartburn due to acid indigestion

85 TABLETS




Actual Size

Compare to Maximum Strength Pepcid® AC active ingredient**

ACTUAL SIZE

NDC 0353-0701-01

ORG0612



LOT NO. EXP. CODE AREA

ITEM 551881

3 191708767 2

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

DO NOT USE IF PRINTED FOIL UNDER CAP IS BROKEN OR MISSING

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Famotidine 20 mg.....Acid reducer</p> <p>Purpose</p> <p>Uses</p> <ul style="list-style-type: none"> ■ relieves heartburn associated with acid indigestion and sour stomach ■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages <p>Warnings</p> <p>Allergy alert: Do not use if you are allergic to famotidine or other acid reducers</p> <p>Do not use</p> <ul style="list-style-type: none"> ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. ■ if you have kidney disease, except under the advice and supervision of a doctor ■ with other acid reducers 	<p>Drug Facts (continued)</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ heartburn with lightheadedness, sweating, or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>
--	---

Drug Facts (continued)

Directions

- adults and children 12 years and over:
 - to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
 - to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- read the directions and warnings before use

Drug Facts (continued)

- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture and light

Inactive ingredients carnauba wax, colloidal silicon dioxide, croscarmellose sodium, lactose (monohydrate), magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc, and titanium dioxide

Questions or comments?
1-800-719-9260

Walmart 100% SATISFACTION GUARANTEED

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
walmart.com ©2012 Walgreen Co.

*Walgreens Pharmacist Survey Study, November 2010.
†This product is not manufactured or distributed by Johnson & Johnson • MERCK Consumer Pharmaceuticals Co., distributor of Maximum Strength Pepcid® AG.

ACID CONTROLLER

famotidine tablet

Product Information

Product Type	HUMAN OTC DRUG LABEL	Item Code (Source)	NDC:0363-0701
Route of Administration	ORAL	DEA Schedule	

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FAMOTIDINE (FAMOTIDINE)	FAMOTIDINE	20 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX	
SILICON DIOXIDE	
CROSCARMELOSE SODIUM	
LACTOSE MONOHYDRATE	

MAGNESIUM STEARATE	
CELLULOSE, MICROCRYSTALLINE	
POLYETHYLENE GLYCOLS	
POLYVINYL ALCOHOL	
TALC	
TITANIUM DIOXIDE	

Product Characteristics

Color	WHITE	Score	no score
Shape	CAPSULE	Size	8mm
Flavor		Imprint Code	L194
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0701-71	1 in 1 CARTON		
1		50 in 1 BOTTLE		
2	NDC:0363-0701-72	1 in 1 CARTON		
2		60 in 1 BOTTLE		
3	NDC:0363-0701-01	1 in 1 CARTON		
3		85 in 1 BOTTLE		
4	NDC:0363-0701-02	5 in 1 CARTON		
4		5 in 1 BLISTER PACK		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA077351	09/26/2006	

Labeler - Walgreen Company (008965063)

Revised: 9/2012

Walgreen Company