# DRUG FACTS

Active ingredient(s) Purpose

## Use(s)

- · temporarily relieves minor aches and pains due to:
- headache
- · muscular aches

## Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 8 caplets in 24 hours, which is the maxiumum daily amount
- · with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

#### Do not no

· with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

## Ask a doctor before use if you have

liver disease

# Ask a doctor or pharmacist before use if you are

· taking the blood thinning drug warfarin

## When using this product

## Stop use and ask a doctor if

- · new symptoms appear
- symptoms do not get better or worsen
- · painful area is red and swollen
- · pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

These could be signs of a serious condition.

# If pregnant or breast-feeding

#### Keep out of reach of children

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-2222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### Directions

- · do not use more than directed
- (see overdose warning)
- adults and children12 years and over: take 2 caplets every 6 hours. Do not take mroe than 8 caplets in 24 hours.
- · children under 12 years: do not use this product in children under 12 years of age; this will provide more than the recommended dose (overdose) and may cause liver damage

# Other information

- store at controlled room temperature 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

# Inactive ingredients

corn starch, crospovidone, D&C red #27 aluminum lake, FD&C #2 aluminum lake, FD&C #6 aluminium lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, sodium starch glycolate, stearic acid, talc, titanium dioxide