DRUG FACTS

Active ingredient(s) Purpose

Use(s)

Directions

- adults and children 6 to 12 years of age (with adult supervision) 2 or 3 sprays in each nostril not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period.
- · children under 6 years of age: ask a doctor.
- To spray, hold bottle with thumb at base and nozzle between first and second fingers. Without tilting head, insert nozzle into nostril Fully depress rim with a firm, even stroke and sniff deeply. Wipe nozzle clean after use.

Warnings

Warnings

For external use only

Do not use

Ask a doctor before use if you have

Ask a doctor before use if you have

- · heart disease
- · high blood pressure
- thyroid disease
- diabetes
- · trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

When using this product

When using this product

- · do not use more than directed
- · do not use more than 3 days. Use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen
- · temporary discomfort such as burning, stinging, sneezing or an increase in nasal discharge may occur
- · use of this container by more than one person may spread infection

Stop use and ask a doctor if

Stop use and ask a doctor if symptoms persist

If pregnant or breast-feeding

Keep out of reach of children

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

DISTRIBUTED BY CARDINAL HEALTH

DUBLIN, OHIO 43017

CIN 4584645

www.myleader.com

1-800-200-6313

Made in Korea

Other information

Other information

- store between 15^O and 25^OC (59^O and 77^OF)
- · retain carton for future reference on full labeling

Inactive ingredients

Inactive ingredients: benzalkonium chloride solution, edetate disodium, polyethylene glycol, sodium phosphate dibasic, sodium phosphate monobasic, water