DRUG FACTS

Active ingredient(s) Purpose

Use(s)

- temporarily relieves minor aches and pains due to:
- headache
- menstrual pain
- minor arthritis pain
- · muscle pain
- toothache
- colds
- · or as recommended by a doctor

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- · facial swelling
- · shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- · take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- · have 3 or more alcoholic drinks every day while using this product
- · take more or for a longer time than directed

if you have ever had an allergic reaction to aspirin or any other pain reliever/ fever reducer

Ask a doctor before use if you have

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- · you have asthma

Ask a doctor or pharmacist before use if you are

taking a prescription drug for diabetes, gout, or arthritis

When using this product

Stop use and ask a doctor if

- allergic reaction occurs. Seek medical help right away. you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- · have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- · redness or swelling is present
- any new symptoms appearringing in the ears or a loss of hearing occurs

These could be signs of a serious condition

If pregnant or breast-feeding

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away: 1-800-222-1222

- adults and children 12 years of age and over: take 1 to 2 tablets every 4 hours, while symptoms persist. Drink a full glass of water with each dose.
- do not take more than 12 tablets in 24 hours unless directed by a doctor
- children under 12 years of age: ask a doctor

Other information

store at controlled room temperature 15°-30° C (59°-86° F)

Inactive ingredients

corn starch, croscarmellose sodium, D&C Yellow #10 Aluminum Lake, FD&C Yellow #6 Aluminum Lake, hypromellose, methacrylic acid copolymer, microcrystalline cellulose, mineral oil, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, talc, titanium dioxide, triethyl citrate