DRUG FACTS

Purpose Active ingredient(s)

Use(s)

for prevention and treatment of these symptoms associated with motion sickness:

- nausea
- · vomiting
- dizziness

Warnings

in children under 2 years of age, unless directed by a doctor

Do not use

Ask a doctor before use if you have

- · a breathing problem such as emphysema or chronic bronchitis
- · glaucoma
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers.

When using this product

- · marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives and tranquilizers may increase drowsiness
- · use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

If pregnant or breast-feeding

Keep out of reach of children
In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- to prevent motion sickness, the first dose should be taken ½ to 1 hour before starting activity
- do not exceed recommended dosage. Not for frequent or prolonged use except on the advice of a doctor

adults and children 12 years of age and over children 6 to under 12 years of age children 2 to under 6 years of age

take 1 to 2 tablets every 4 to 6 hours; not to exceed 8 tablets in 24 hours or as directed by a doctor take 1/2 to 1 tablet every 6 to 8 hours; not to exceed 3 tablets in 24 hours or as directed by a doctor take 1/4 to 1/2 tablet every 6 to 8 hours; not to exceed 1-1/2 tablets in 24 hours or as directed by a doctor.

Other information

Inactive ingredients

Croscarmellose Sodium, Dicalcium Phosphate, Magnesium Stearate and Microcrystalline Cellulose.