DRUG FACTS

Active ingredient(s) ***Purpose***

Use(s)

- temporarily relieves minor aches and pains due to:
- headache
- backache
- toothache
- the common cold
- minor pain of arthritis
- muscular aches
- premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Do not use
with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have
liver disease

Ask a doctor or pharmacist before use if you are
taking the blood thinning drug warfarin

When using this product

Stop use and ask a doctor if
- pain gets worse or lasts more than 10 days in adults and children
- pain gets worse or lasts more than 5 days in children under 12
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding

Keep out of reach of childrenOverdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	• take 2 tablets every 4 to 6 hours while symptoms last • do not take more than 12 tablets in 24 hours • do not take for more than 10 days unless directed by a doctor
children 6-11 years	• take 1 tablet every 4 to 6 hours while symptoms last • do not take more than 5 tablets in 24 hours • do not take for more than 5 days unless directed by a doctor
children under 6 years	• do not use this adult regular strength product in children under 6 years of age; this will provide more than the recommended dose (overdose) of acetaminophen and may cause liver damage

Other information

This product contains acetaminophen. Severe liver damage may occur if:
- adults take more than 12 tablets in 24 hours, which is the maximum daily amount
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product.
- store at controlled room temperature 25°C (77°F); excursions permitted 15º-30ºC (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, povidone, sodium starch glycolate*, stearic acid
*may contain this ingredient