

TERBINAFINE HYDROCHLORIDE - terbinafine hydrochloride tablet Bryant Ranch Prepack


Terbinafine Hydrochloride Tablets

Rx only Prescribing Information

DESCRIPTION

Terbinafine Hydrochloride Tablets contain the synthetic allylamine antifungal compound terbinafine hydrochloride. Chemically, terbinafine hydrochloride is (E)-N-(6,6dimethyl-2hepten-4-ynyl)-N-methyl-1-naphthalenemethanamine hydrochloride. The empirical formula

C₂₁H₃₁ClN with a molecular weight of 327.90, and the following structural formula: 2126



Terbinafine hydrochloride is a white to off-white fine crystalline powder. It is freely soluble in methanol and methylene chloride, soluble in ethanol, and slightly soluble in water.

Each tablet contains:

terbinafine hydrochloride (equivalent to 250 mg base) *Active Ingredients:*

colloidal silicon dioxide, NF; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF *Inactive Ingredients:*

CLINICAL PHARMACOLOGY

Pharmacokinetics

Following oral administration, terbinafine is well absorbed (>70%) and the bioavailability of Terbinafine Hydrochloride Tablets as a result of first-pass metabolism is approximately 40%.

Peak plasma concentrations of 1 appear within 2 h after a single 250 mg dose; the AUC (area under the curve) is approximately . An increase in the AUC of terbinafine of less than 20% is observed when terbinafine hydrochloride is administered with food. No clinically relevant age-dependent changes in steady-state plasma concentrations of terbinafine have been reported. In patients with renal impairment (creatinine clearance 50mL/min) or hepatic cirrhosis, the clearance of terbinafine is decreased by approximately 50% compared to normal volunteers. No effect of gender on the blood levels of terbinafine was detected in clinical trials. In plasma, terbinafine is >99% bound to plasma proteins and

there are no specific binding sites. At steady-state, in comparison to a single dose, the peak concentration of terbinafine is 25% higher and plasma AUC increases by a factor of 2.5; the increase in plasma AUC is consistent with an effective half-life of ~36 hours. Terbinafine is distributed to the sebum and skin. A terminal half-life of 200-400 h may represent the slow elimination of terbinafine from tissues such as skin and adipose. Prior to excretion, terbinafine is extensively metabolized. No metabolites have been identified that have antifungal activity similar to terbinafine. Approximately 70% of the administered dose is eliminated in the urine. $\mu\text{g/mL}$, $4.56 \mu\text{g}\cdot\text{h/mL} \leq$

Microbiology

Terbinafine hydrochloride is a synthetic allylamine derivative. Terbinafine hydrochloride is hypothesized to act by inhibiting squalene epoxidase, thus blocking the biosynthesis of ergosterol, an essential component of fungal cell membranes. , mammalian squalene epoxidase is only inhibited at higher (4000 fold) concentrations than is needed for inhibition of the dermatophyte enzyme. Depending on the concentration of the drug and the fungal species test , terbinafine hydrochloride may be fungicidal. However, the clinical significance of data is unknown. Terbinafine has been shown to be active against most strains of the following microorganisms both and in clinical infections as described in the section: *In vitro* *in vitro* *in vitro* *in vitro* INDICATIONS AND USAGE

Trichophyton mentagrophytes

Trichophyton rubrum

The following data are available, but their clinical significance is unknown. , terbinafine exhibits satisfactory MIC's against most strains of the following microorganisms; however, the safety and efficacy of terbinafine in treating clinical infections due to these microorganisms have not been established in adequate and well-controlled clinical trials: *in vitro* *In vitro*

Candida albicans

Epidermophyton floccosum

Scopulariopsis brevicaulis

CLINICAL STUDIES

The efficacy of Terbinafine Hydrochloride Tablets in the treatment of onychomycosis is illustrated by the response of patients with toenail and/or fingernail infections who participated in three US/Canadian placebo-controlled clinical trials. Results of the first toenail study, as assessed at week 48 (12 weeks of treatment with 36 weeks follow-up after completion of therapy), demonstrated mycological cure, defined as simultaneous occurrence of negative KOH plus negative culture, in 70% of patients. Fifty-nine percent (59%) of patients experienced effective treatment (mycological cure plus 0% nail involvement or >5mm of new unaffected nail growth); 38% of patients demonstrated mycological cure plus clinical cure (0% nail involvement). In a second toenail study of dermatophytic onychomycosis, in which non-dermatophytes were also cultured, similar efficacy against the dermatophytes was demonstrated. The pathogenic role of the non-dermatophytes cultured in the presence of dermatophytic onychomycosis has not been established. The clinical significance of this association is unknown. Results of the fingernail study, as assessed at week 24 (6 weeks of treatment with 18 weeks follow-up after completion of therapy), demonstrated mycological cure in 79% of patients, effective treatment in 75% of the patients, and mycological cure plus clinical cure in 59% of the patients.

The mean time to overall success was approximately 10 months for the first toenail study and 4 months for the fingernail study. In the first toenail study, for patients evaluated at least six months after achieving clinical cure and at least one year after completing terbinafine hydrochloride therapy, the clinical relapse rate was approximately 15%.

INDICATIONS AND USAGE

Terbinafine Hydrochloride Tablets are indicated for the treatment of onychomycosis of the toenail or fingernail due to dermatophytes (tinea unguium) (see and). **DOSAGE AND ADMINISTRATION**
CLINICAL STUDIES

Prior to initiating treatment, appropriate nail specimens for laboratory testing (KOH preparation, fungal culture, or nail biopsy) should be obtained to confirm the diagnosis of onychomycosis.

CONTRAINDICATIONS

Terbinafine Hydrochloride Tablets are contraindicated in individuals with hypersensitivity to terbinafine or to any other ingredients of the formulation.

WARNINGS

Rare cases of liver failure, some leading to death or liver transplant, have occurred with the use of Terbinafine Hydrochloride Tablets for the treatment of onychomycosis in individuals with and without pre-existing liver disease.

In the majority of liver cases reported in association with terbinafine hydrochloride use, the patients had serious underlying systemic conditions and an uncertain causal association with terbinafine hydrochloride. The severity of hepatic events and/or their outcome may be worse in patients with active or chronic liver disease (see). Treatment with Terbinafine Hydrochloride Tablets should be discontinued if biochemical or clinical evidence of liver injury develops (see below).

PRECAUTIONS

There have been isolated reports of serious skin reactions (e.g., Stevens-Johnson Syndrome and toxic epidermal necrolysis). If progressive skin rash occurs, treatment with terbinafine hydrochloride should be discontinued.

PRECAUTIONS

General

Terbinafine Hydrochloride Tablets are not recommended for patients with chronic or active liver disease. Before prescribing Terbinafine Hydrochloride Tablets, pre-existing liver disease should be assessed. Hepatotoxicity may occur in patients with and without pre-existing liver disease. Pretreatment serum transaminase (ALT and AST) tests are advised for all patients before taking Terbinafine Hydrochloride Tablets. Patients prescribed Terbinafine Hydrochloride Tablets should be warned to report immediately to their physician any symptoms of persistent nausea, anorexia, fatigue, vomiting, right upper abdominal pain or jaundice, dark urine or pale stools (see). Patients with these symptoms should discontinue taking oral terbinafine, and the patient's liver function should be immediately evaluated. **WARNINGS**

In patients with renal impairment (creatinine clearance <50 mL/min), the use of terbinafine hydrochloride has not been adequately studied, and therefore, is not recommended (see ,). **CLINICAL PHARMACOLOGY**
Pharmacokinetics

During postmarketing experience, precipitation and exacerbation of cutaneous and systemic lupus erythematosus have been reported infrequently in patients taking terbinafine hydrochloride. Terbinafine hydrochloride therapy should be discontinued in patients with clinical signs and symptoms suggestive of lupus erythematosus.

Changes in the ocular lens and retina have been reported following the use of Terbinafine Hydrochloride Tablets in controlled trials. The clinical significance of these changes is unknown. Transient decreases in absolute lymphocyte counts (ALC) have been observed in controlled clinical trials. In placebo-controlled trials, 8/465 terbinafine hydrochloride treated patients (1.7%) and 3/137 placebo-treated patients (2.2%) had decreases in ALC to below 1000/mm³ on two or more occasions.

The clinical significance of this observation is unknown. However, in patients with known or suspected immunodeficiency, physicians should consider monitoring complete blood counts in individuals using Terbinafine Hydrochloride therapy for greater than six weeks. Isolated cases of severe neutropenia have been reported. These were reversible upon discontinuation of terbinafine hydrochloride with or without supportive therapy. If clinical signs and symptoms suggestive of secondary infection occur, a complete blood count should be obtained. If the neutrophil count is $<1,000$ cells/mm³, terbinafine hydrochloride should be discontinued and supportive management started.

Drug Interactions

studies have shown that terbinafine is an inhibitor of the CYP450 206 isozyme. Drugs predominantly metabolized by the CYP450 206 isozyme include the following drug classes; tricyclic antidepressants, selective serotonin reuptake inhibitors, beta-blockers, antiarrhythmics class 1C (e.g. flecainide and propafenone) and monoamine oxidase inhibitors Type B. Coadministration of Terbinafine Hydrochloride Tablets should be done with careful monitoring and may require a reduction in dose of the 2D6-metabolized drug. In a study to assess the effects of terbinafine on desipramine in healthy volunteers characterized as normal metabolizers, the administration of terbinafine resulted in a 2-fold increase in C_{max} and a 5-fold increase in AUC. In this study, these effects were shown to persist at the last observation at 4 weeks after discontinuation of terbinafine hydrochloride. *In vivo*

studies with human liver microsomes showed that terbinafine does not inhibit the metabolism of tolbutamide, ethinylestradiol, ethoxycoumarin, and cyclosporine. *In vitro*

drug-drug interaction studies conducted in healthy volunteer subjects showed that terbinafine does not affect the clearance of antipyrine or digoxin. *In vivo*

Terbinafine decreases the clearance of caffeine by 19%. Terbinafine increases the clearance of cyclosporine by 15%.

There have been spontaneous reports of increase or decrease in prothrombin times in patients concomitantly taking oral terbinafine and warfarin, however, a causal relationship between Terbinafine Hydrochloride Tablets and these changes has not been established.

Terbinafine clearance is increased 100% by rifampin, a CYP450 enzyme inducer, and decreased 33% by cimetidine, a CYP450 enzyme inhibitor. Terbinafine clearance is unaffected by cyclosporine.

There is no information available from adequate drug-drug interaction studies with the following classes of drugs: oral contraceptives, hormone replacement therapies, hypoglycemics, theophyllines, phenytoins, thiazide diuretics, and calcium channel blockers.

Carcinogenesis, Mutagenesis, Impairment of Fertility

In a 28-month oral carcinogenicity study in rats, an increase in the incidence of liver tumors was observed in males at the highest dose tested, 69 mg/kg/day [2x the Maximum Recommended Human Dose (MRHD) based on AUC comparisons of the parent terbinafine]; however, even though dose limiting toxicity was not achieved at the highest tested dose, higher doses were not tested.

The results of a variety of (mutations in *E. coli* and *S. .*, DNA repair in rat hepatocytes, mutagenicity in Chinese hamster fibroblasts, chromosome aberration and sister chromatid exchanges in Chinese hamster lung cells), and (chromosome aberration in Chinese hamsters, micronucleus test in mice) genotoxicity tests gave no evidence of a mutagenic or clastogenic potential. Oral reproduction studies in rats at doses up to 300 mg/kg/day (approximately 12x the MRHD based on body surface area comparisons, BSA) did not reveal any specific effects on fertility or other reproductive parameters. Intravaginal application of terbinafine hydrochloride at 150 mg/day in pregnant rabbits did not increase the incidence of abortions or premature deliveries nor affect fetal parameters. *in vitro* typhimurium *in vivo*

Pregnancy

Oral reproduction studies have been performed in rabbits and rats at doses up to 300 mg/kg/day (12x to

23x the MRHD, in rabbits and rats, respectively, based on BSA) and have revealed no evidence of impaired fertility or harm to the fetus due to terbinafine. There are, however, no adequate and well-controlled studies in pregnant women. **Pregnancy Category B:**

Because animal reproduction studies are not always predictive of human response, and because treatment of onychomycosis can be postponed until after pregnancy is completed, it is recommended that terbinafine hydrochloride not be initiated during pregnancy.

Nursing Mothers

After oral administration, terbinafine is present in breast milk of nursing mothers. The ratio of terbinafine in milk to plasma is 7:1. Treatment with terbinafine hydrochloride is not recommended in nursing mothers.

The safety and efficacy of Terbinafine Hydrochloride Tablets have not been established in pediatric patients.

ADVERSE REACTIONS

The most frequently reported adverse events observed in the three US/Canadian placebocontrolled trials are listed in the table below. The adverse events reported encompass gastrointestinal symptoms (including diarrhea, dyspepsia, and abdominal pain), liver test abnormalities, rashes, urticaria, pruritus, and taste disturbances. In general, the adverse events were mild, transient, and did not lead to discontinuation from study participation.

	Adverse Event		Discontinuation	
	Terbinafine Hydrochloride Tablets	Placbo	Terbinafine Hydrochloride Tablets	Placbo
	(%) n=465	(%) n=137	(%) n=465	(%) n=137
Headache	12.9	9.5	0.2	0.0
Gastrointestinal Symptoms:				
Diarrhea	5.6	2.9	0.6	0.0
Dyspepsia	4.3	2.9	0.4	0.0
Abdominal Pain	2.4	1.5	0.4	0.0
Nausea	2.6	2.9	0.2	0.0
Flatulence	2.2	2.2	0.0	0.0
Dermatological Symptoms:				
Rash	5.6	2.2	0.9	0.7
Pruritus	2.8	1.5	0.2	0.0
Urticaria	1.1	0.0	0.0	0.0
Liver Enzyme Abnormalities*	3.3	1.4	0.2	0.0
Taste Disturbance	2.8	0.7	0.2	0.0
Visual Disturbance	1.1	1.5	0.9	0.0

*Liver enzyme abnormalities $\geq 2x$ the upper limit of the normal range.

Adverse events, based on worldwide experience with Terbinafine Hydrochloride Tablets use, include:

idiosyncratic and symptomatic hepatic injury and more rarely, cases of liver failure, some leading to death or liver transplant, (see and), serious skin reactions (see), severe neutropenia (see), thrombocytopenia, angioedema and allergic reactions (including anaphylaxis). Psoriasiform eruptions or exacerbation of psoriasis, acute generalized exanthematous pustulosis and precipitation and exacerbation of cutaneous and systemic lupus erythematosus have been reported in patients taking Terbinafine Hydrochloride Tablets. Terbinafine Hydrochloride Tablets may cause taste disturbance (including taste loss) which usually recovers within several weeks after discontinuation of the drug. There have been isolated reports of prolonged (greater than one year) taste disturbance. Taste disturbances associated with oral terbinafine have been reported to be severe enough to result in decreased food intake leading to significant and unwanted weight loss.

WARNINGS/PRECAUTIONS/WARNINGS/PRECAUTIONS

Other adverse reactions which have been reported include malaise, fatigue, vomiting, arthralgia, myalgia, and hair loss.

Clinical adverse effects reported spontaneously since the drug was marketed include altered prothrombin time (prolongation and reduction) in patients concomitantly treated with warfarin and Terbinafine Hydrochloride Tablets and agranulocytosis (rare).

OVERDOSAGE

Clinical experience regarding overdose with Terbinafine Hydrochloride Tablets is limited. Doses up to 5 grams (20 times the therapeutic daily dose) have been taken without inducing serious adverse reactions. The symptoms of overdose included nausea, vomiting, abdominal pain, dizziness, rash, frequent urination, and headache.

DOSAGE AND ADMINISTRATION

Terbinafine Hydrochloride Tablets, one 250 mg tablet, should be taken once daily for 6 weeks by patients with fingernail onychomycosis. Terbinafine hydrochloride, one 250 mg tablet, should be taken once daily for 12 weeks by patients with toenail onychomycosis. The optimal clinical effect is seen some months after mycological cure and cessation of treatment. This is related to the period required for outgrowth of healthy nail.

ANIMAL TOXICOLOGY

A wide range of studies in mice, rats, dogs, and monkeys, and studies using rat, monkey, and human hepatocytes suggest that peroxisome proliferation in the liver is a rat-specific finding. However, other effects, including increased liver weights and APTT, occurred in dogs and monkeys at doses giving C₅₀ trough levels of the parent terbinafine 2-3x those seen in humans at the MRHD. Higher doses were not tested. *in vivo* and *in vitro*

Manufactured by: InvaGen Pharmaceuticals, Inc., 7 Oser Avenue, Hauppauge, NY 11788
Manufactured for: Camber Pharmaceuticals, Inc. Piscataway, NJ 08854
Rev: 05/09 Barcode: 209-05-2009

Terbinafine 250mg tab.

TERBINAFINE HYDROCHLORIDE

terbinafine hydrochloride tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:63629-2807(NDC:31722-209)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TERBINAFINE HYDROCHLORIDE (UNII: 012C11ZU6G) (TERBINAFINE - UNII:G7RIW8S0XP)	TERBINAFINE	250 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	

Product Characteristics

Color	WHITE	Score	no score
Shape	ROUND	Size	11mm
Flavor		Imprint Code	209;IG
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63629-2807-1	30 in 1 BOTTLE		
2	NDC:63629-2807-2	60 in 1 BOTTLE		
3	NDC:63629-2807-3	90 in 1 BOTTLE		
4	NDC:63629-2807-4	100 in 1 BOTTLE		
5	NDC:63629-2807-5	7 in 1 BOTTLE		
6	NDC:63629-2807-6	40 in 1 BOTTLE		
7	NDC:63629-2807-7	45 in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA077533	01/01/2011	

Labeler - Bryant Ranch Prepack (171714327)

Registrant - Bryant Ranch Prepack (171714327)

Establishment

Name	Address	ID/FEI	Business Operations
Bryant Ranch Prepack		171714327	REPACK(63629-2807) , RELABEL(63629-2807)

Revised: 8/2011

Bryant Ranch Prepack