DRUG FACTS

Purpose Active ingredient(s)

Use(s)

temporarily relieves minor aches and pains due to

- · headache
- muscular aches
- minor pain of arthritis
- toothache
- backache
- the common cold
- menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- · hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chances are higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- · take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- · take more or for a longer time than directed

Do not use

if you have ever had an allergic reaction to other pain reliever/fever reduce

right before or after heart surgery

Ask a doctor before use if you have

- the stomach bleeding warning applies to you
- · you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin to prevent heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- · any new symptoms appear

If pregnant or breast-feeding

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control center right away (1-800-222-1222).

Directions

- · do not take more than directed
- the smallest effective dose should be used
- do not take longer than 10 days, unless directed by a doctor (see Warnings)
- Adults and children 12 years and older:
- take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours unless directed by a doctor
- Children under 12 years: ask a doctor

Other information

- store between 20-25 degrees Celsius (68-77 degrees Fahrenheit).
- Read all warnings and directions before use
- · Do not use if seal under bottle cap is broken or missing.

Inactive ingredients

Anhydrous Lacutose, Carnauba Wax, Colloidal Silicon Dioxide, Hypromellose, Iron Oxide Red, Magnesium Stearate, Microcrystalline Cellulose, Polydextrose, Polyethylene Glycol, Povidone, Sodium Lauryl Sulfate, Sodium Starch Glycolate, Starch (Corn), Titanium Dioxide