DRUG FACTS

Purpose Active ingredient(s)

Use(s)

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

Warnings

Do not use

if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

Ask a doctor before use if you have

liver or kidney disease. Your doctor should determine if you need a different dose.

Ask a doctor or pharmacist before use if you are

taking tranquilizers or sedatives.

When using this product

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

adults and children 6 years and over	one 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.
adults 65 years and over	ask a doctor
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information

- do not use if blister unit is broken or torn (Blister Only)
- do not use if printed foil under cap is broken or missing (Bottle Only) store between 20° to $25^\circ C$ (68° to $77^\circ F)$

Inactive ingredients

corn starch, FD&C blue no. 1 aluminum lake, hypromellose, lactose monohydrate, magnesium stearate, polydextrose, polyethylene glycol, povidone, titanium dioxide, triacetin