# DRUG FACTS

PurposeActive ingredient(s)

#### Use(s)

for the relief of

- acid indigestion
- heartburn
- sour stomach
- upset stomach associated with these symptoms pressure and bloating commonly referred to as gas

## Warnings

#### Do not use

## Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

### Ask a doctor or pharmacist before use if you are

presently taking a prescription drug. Antacids may interact with certain prescription drugs.

### When using this product

do not exceed 8 teaspoonsful in a 24-hour period, or use the maximum dosage for more than 2 weeks, unless directed by a doctor

## Stop use and ask a doctor if

#### If pregnant or breast-feeding

Keep out of reach of children

#### Directions

- shake well before using
- shake wen before using adults and children 12 years and older: take 2 to 4 teaspoonsful two times a day or as directed by a physician do not take more than 8 teaspoonsful in 24 hours or use the maximum dosage for more than 2 weeks
- children under 12 years: consult a physician

## $Other\ information$

- each teaspoonful contains: magnesium 175 mg and sodium 2 mg does not meet USP requirements for preservative effectiveness

## Inactive ingredients

 $butyl paraben, \tilde{f}lavor, glycerin, hydroxyethyl cellulose, propylene glycol, propylparaben, purified water, saccharin sodium, simethicone emulsion, sorbitol, sorbitol solution$