DURVET PENNCHLOR 50 - chlortetracycline calcium powder Durvet, Inc.
durvet
NDC: 30798-373-56
Pennchlor 50
CHLORTETRACYCLINE
Type A Medicated Article
FOR USE IN THE MANUFACTURE OF MEDICATED FEEDS
ACTIVE INGREDIENTS:
$Chlortetracycline \ as \ Chlortetracycline \ calcium \ complex \ equivalent \ to \ 50 \ grams \ Chlortetracycline$
Hydrochloride/lb.
INGREDIENTS:
Chlortetracycline, Calcium Carbonate, Roughage Products and Mineral Oil.
See back panel for directions and warnings.
CAUTION: For use in Dry Feeds ONLY - NOT FOR USE IN LIQUID FEED SUPPLEMENTS.
NADA 138-935 APPROVED BY FDA
Restricted Drug (California)-
Use Only as Directed
NOT FOR HUMAN USE
NET WT.
50 lb (22.7 kg)
durvet
Manufactured for
DURVET, INC.
Blue Springs, Missouri 64014
durvet
NDC: 30798-373-56
Pennchlor 50
CHLORTETRACYCLINE
DIRECTIONS FOR USE

Use Levels of
Chlortetracycline

Chlorie lbs. of
Pennchlor
50
per ton

Indications for Use

CHICKENS

For Broiler/fryer chickens: For an increased rate of weight gain and improved feed efficiency. WARNING: Do not feed to chickens producing eggs for human consumption. Zero-day withdrawal period.	10-50 g/ton	0.2-1.0
For Chickens: Control of infectious synovitis caused by Mycoplasma synoviae susceptible to chlortetracycline. (Feed continuously for 7-14	100-200 g/ton	2.0-4.0
days) WARNING: Do not feed to chickens producing eggs for human consumption. Zero-day withdrawal period.		
For Chickens: Control of chronic respiratory disease (CRD) and air sac infections caused by Mycoplasma gallisepticum and Escherichia coli susceptible to chlortetracycline. (Feed continuously for 7-14	200-400 g/ton	4.0-8.0
days) WARNING: Do not feed to chickens producing eggs for human consumption. Zero-day withdrawal period.		
For Chickens: Reduction of mortality due to Escherichia coli infections susceptible to chlortetracycline. (Feed for 5 days) WARNING: Do not feed to chickens producing eggs for human consumption. Withdraw 24 hours prior to slaughter.	500 g/ton	10.0
TURKEYS For Turkeys: Growing Turkeys: For an increased rate of weight gain and improved feed efficiency. WARNING: Do not feed to turkeys producing eggs for human consumption. Zero-day withdrawal period.	10-50 g/ton	0.2-1.0
For Turkeys: Control of infectious synovitis caused by Mycoplasma synoviae susceptible to chlortetracycline. (Feed continuously for 7-14 days) WARNING: Do not feed to turkeys producing eggs for human consumption. Zero-day withdrawal period.	200 g/ton	4.0
For Turkeys: Control of hexamitiasis caused by Hexamita meleagrides susceptible to chlortetracycline. (Feed continuously 7-14 days) WARNING : Do not feed to turkeys producing eggs for human consumption. Zero-day withdrawal period.	400 g/ton	8.0

For Turkeys: Turkey poults not over 4 weeks of age: Reduction of mortality due to paratyphoid caused by <u>Salmonella typhimurium</u> susceptible to chlortetracycline.	400 g/ton	8.0
For Turkeys: Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to chlortetracycline. (Feed continuously for 7-14 days) WARNING: Do not feed to turkeys producing eggs for human consumption. Zero-day withdrawal period.	25 mg/lb body weight/day	
SHEEP For Growing Sheep: For an increased rate of weight gain and improved feed efficiency.	20-50 g/ton	0.4-1.0
For Breeding Sheep: Reducing the incidence of (vibrionic) abortion caused by <u>Campylobacter fetus</u> infections susceptible to chlortetracycline. WARNING: Zero-day withdrawal period.	80 mg/head/day	
		lbs. of Pennchlor
Indications for Use	Use Levels of Chlortetracycline	50 per ton
Indications for Use SWINE For Growing Swine: For an increased rate of weight gain improved feed efficiency.		50
SWINE For Growing Swine: For an increased rate of weight gain	Chlortetracycline	50 per ton
SWINE For Growing Swine: For an increased rate of weight gain improved feed efficiency. For Swine: Reducing the incidence of cervical lymphadenitis (jowl abscesses) caused by Group E Streptococci susceptible to	Chlortetracycline 10-50 g/ton	50 per ton 0.2-1.0

WARNING: Zero-day withdrawal period.

CALVES, BEEF CATTLE, AND NONLACTATING DAIRY CATTLE

WARNING: A WITHDRAWAL PERIOD HAS NOT BEEN

ESTABLISHED

FOR THIS PRODUCT IN PRE-RUMINATING CALVES. DO

NOT USE

IN CALVES TO BE PROCESSED FOR VEAL.

For Calves (up to 250 lbs.): For an increased rate of weight gain 0.1 mg/lb body and weight/day

improved feed efficiency.

For Calves (250-400 lbs.): For an increased rate of weight gain and 25-70 mg/head/day

improved feed efficiency.

For Growing Cattle (over 400 lbs.): For an increased rate of weight 70 mg/head/day

gain, improved feed efficiency and reduction of liver condemnation

due to liver abscesses.

For Beef Cattle: For the control of bacterial pneumonia associated with 350 mg/head/day

shipping fever complex caused by <u>Pasteurella</u> spp. susceptible to chlortetracycline.

WARNING: Zero-day withdrawal period.

For Beef Cattle (under 700 lbs.): Control of active infection of anaplasmosis 350 mg/head/day

caused by Anaplasma marginale susceptible to chlortetracycline.

WARNING: Zero-day withdrawal period.

For Beef Cattle (over 700 lbs.): Control of active infection of weight/day

anaplasmosis caused by <u>Anaplasma marginale</u> susceptible to chlortetraycline.

WARNING: Zero-day withdrawal period


For Calves, Beef, and Nonlactating Dairy Cattle: For treatment 10 mg/lb body of weight/day

bacterial enteritis caused by <u>Escherichia coli</u> and bacterial pneumonia

caused by <u>Pasteurella multocida</u> susceptible to chlortetracycline.

(Treat for not more than 5 days)

Enter section text here



CAUTION: For use in Dry Feeds ONLY.

NOT FOR USE IN LIQUID FEED SUPPLEMENTS

TAKE TIME OBSERVE LABEL DIRECTIONS

REV 12/05

BAGT

durvet

Manufactured for

DURVET, INC.

Blue Springs, Missouri 64014

DURVET PENNCHLOR 50 chlortetracycline powder Product Information

Product Type		OTC TYPE A MEDICATED ARTICLE ANIMAL DRUG LABEL Item Code (Source)		NDC:30798- 373
	Route of Administration	ORAL	DEA Schedule	

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
Chlortetracycline Calcium (Chlortetracycline)	Chlortetracycline Calcium	110 g in 1 kg

Packaging				
	# Item Code	Package Description	Marketing Start Date	Marketing End Date
	1 NDC:30798-373-56	22.7 kg in 1 BAG		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA138935	12/08/2007	

Labeler - Durvet, Inc. (056387798)

Revised: 11/2012 Durvet, Inc.