

GOOD SENSE ALL DAY PAIN RELIEF- naproxen sodium tablet
L Perrigo Company

Perrigo All Day Pain Relief Drug Facts

Active ingredient (in each tablet)

Naproxen sodium 220 mg

(naproxen 200 mg) (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - you have difficulty swallowing
 - it feels like the pill is stuck in your throat
 - redness or swelling is present in the painful area
 - any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older	<ul style="list-style-type: none">• take 1 tablet every 8 to 12 hours while symptoms last• for the first dose you may take 2 tablets within the first hour• do not exceed 2 tablets in any 8- to 12-hour period• do not exceed 3 tablets in a 24-hour period
Children under 12 years	ask a doctor

Other information

- each tablet contains: sodium 21 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients

FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Questions or comments?

1-800-719-9260

Principal Display Panel

Strength to Last 12 Hours

See New Warnings Information

All Day Pain Relief

Naproxen Sodium Tablets, 220 mg

Pain Reliever & Fever Reducer (NSAID)

Actual Size


Compare to active ingredient of Aleve® Tablets

EXP.

LOT NO.

LOT NO.

N 0113-9490-78 7



GOODSENSE.

NDC 0113-9490-78

Strength To Last 12 Hours

See New Warnings Information

All Day Pain Relief

Naproxen Sodium Tablets, 220 mg
Pain Reliever & Fever Reducer (NSAID)



Actual Size

100 Tablets

Compare to active ingredient of Aleve® Tablets



Drug Facts (continued)

Questions or comments?

1-800-719-9260


KEEP CARTON FOR REFERENCE

Gluten Free

Distributed By

Perrigo®

Allegan, MI 49010



Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

■ do not take more than directed
■ the smallest effective dose should be used
■ drink a full glass of water with each dose

Adults and children 12 years and older
■ take 1 tablet every 8 to 12 hours while symptoms last
■ for the first dose you may take 2 tablets within the first hour
■ do not exceed 2 tablets in any 8- to 12-hour period
■ do not exceed 3 tablets in a 24-hour period
■ ask a doctor

Children under 12 years

Other information

■ each tablet contains: sodium 21 mg
■ store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

DO NOT USE IF PRINTED FOIL UNDER
CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet) Purposes

Naproxen sodium 220 mg Pain reliever/
(naproxen 200 mg) (NSAID)*fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed ▶

Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug



When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear ▶





GOOD SENSE ALL DAY PAIN RELIEF

naproxen sodium tablet

Product Information

Product Type	HUMAN OTC DRUG LABEL	Item Code (Source)	NDC:0113-9490
Route of Administration	ORAL	DEA Schedule	

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (NAPROXEN)	NAPROXEN SODIUM	220 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSES	
MAGNESIUM STEARATE	
CELLULOSE, MICROCRYSTALLINE	
POLYETHYLENE GLYCOLS	
POVIDONES	
TALC	
TITANIUM DIOXIDE	

Product Characteristics

Color	BLUE (Light Blue)	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	L490
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0113-9490-62	1 in 1 CARTON		
1		24 in 1 BOTTLE		

2	NDC:0113-9490-71	1 in 1 CARTON		
2		50 in 1 BOTTLE		
3	NDC:0113-9490-78	1 in 1 CARTON		
3		100 in 1 BOTTLE		
4	NDC:0113-9490-76	1 in 1 CARTON		
4		120 in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074661	01/14/1997	

Labeler - L Perrigo Company (006013346)

Registrant - L Perrigo Company (006013346)

Revised: 11/2012

L Perrigo Company