DRUG FACTS

Active ingredient(s)	*Purpose*

Use(s)

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

Warnings

Do not use

Do not use if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.

Ask a doctor before use if you have

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.

Ask a doctor or pharmacist before use if you are

Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

When using this product

When using this product

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery.

Stop use and ask a doctor if

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding

Keep out of reach of children

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- use only with enclosed dosing cup
- adults and children 6 years and over: 1 teaspoonful (5 mL) or 2 teaspoonfuls (10 mL) once daily depending upon severity of symptoms; do not take more than 2 teaspoonfuls (10 mL) in 24 hours.
- adults 65 years and older: 1 teaspoonful (5 mL) once daily; do not take more than 1 teaspoonful (5 mL) in 24 hours.
- children 2 to under 6 years of age: 1/2 teaspoonful (2.5 mL) once daily. If needed, dose can be increased to a maximum of 1 teaspoonful (5 mL) once daily or 1/2 teaspoonful (2.5 mL) every 12 hours. Do not give more than 1 teaspoonful (5 mL) in 24 hours.
- children under 2 years of age: ask a doctor
- consumers with liver or kidney disease: ask a doctor

Other information

- store between 20° to 25°C (68° to 77°F)

Dosing cup should be washed and left to air dry after each use. Do not use if carton is opened or if imprinted safety seal is torn or missing.

Dist. by:

Taro Pharmaceuticals U.S.A., Inc.

Hawthorne, NY 10532

Inactive ingredients

bubble gum artificial flavor, glacial acetic acid, glycerin, methylparaben, propylene glycol, propylparaben, purified water, sodium acetate anhydrous, sucralose