DRUG FACTS

Active ingredient(s) ***Purpose***

Use(s)

Uses
temporarily relieves the following menstrual and premenstrual symptoms:
water-weight gain # bloating
backache # headache # minor pain of menstrual and premenstrual cramps

Warnings

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
more than 8 caplets in 24 hours, which is the maximum daily amount
with other drugs containing acetaminophen
3 or more alcoholic drinks every day while using this product
Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
Ask a doctor before use if you have liver disease
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin
Stop use and ask a doctor if
pain lasts more than 10 days or gets worse
redness or swelling is present in the painful area
any new symptoms appear

Do not use

Ask a doctor before use if you have

Ask a doctor or pharmacist before use if you are

When using this product

Stop use and ask a doctor if

If pregnant or breast-feeding

Keep out of reach of children
KEEP OUT OF REACH OF CHILDREN. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Directions # adults: take 2 caplets every 6 hours, not more than 8 caplets in 24 hours # children under 12 years: ask a doctor

Other information

Inactive ingredients

Inactive ingredients
croscarmellose sodium, crospovidone, FDC red 40 aluminum lake, hydroxypropyl methylcellulose, maltodextrin*, magnesium stearate*, microcrystalline cellulose, polyethylene glycol, polysorbate 80*, povidone, sodium starch glycolate*, starch, stearic acid, titanium dioxide
*contains one or more of these ingredients