WARNING: FETAL TOXICITY

When pregnancy is detected, discontinue benazepril hydrochloride as soon as possible. Drugs that act directly on the renin-angiotensin system can cause injury and death to the developing fetus. See WARNINGS, Fetal Toxicity

DESCRIPTION

Benazepril hydrochloride, USP is a white to off-white crystalline powder, soluble (> 100 mg/mL) in water, in ethanol, and in methanol. Its chemical name is 3-[[1-(ethoxy-carbonyl)-3-phenyl-(1S)-propyl]amino]-2,3,4,5-tetrahydro-2-oxo-1H-1-(3S)-benzazepine-1-acetic acid monohydrochloride; its structural formula is



C₂₄H₂₈N₂O₅•HCl M.W. 460.96

Benazeprilat, the active metabolite of benazepril, is a non-sulfhydryl angiotensin-converting enzyme inhibitor. Benazepril is converted to benazeprilat by hepatic cleavage of the ester group.

Benazepril hydrochloride, USP is supplied as tablets containing 5 mg, 10 mg, 20 mg, and 40 mg of benazepril hydrochloride, USP for oral administration. The inactive ingredients are: colloidal silicon dioxide, corn starch, crospovidone, hypromellose, iron oxide yellow, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, titanium dioxide, and triacetin. Additionally, the 5 mg strength contains talc, the 20 mg strength contains iron oxide black and iron oxide red, and the 40 mg strength contains iron oxide red.

CLINICAL PHARMACOLOGY

Mechanism of Action

Benazepril and benazeprilat inhibit angiotensin-converting enzyme (ACE) in human subjects and animals. ACE is a peptidyl dipeptidase that catalyzes the conversion of angiotensin I to the vasoconstrictor substance, angiotensin II. Angiotensin II also stimulates aldosterone secretion by the adrenal cortex.

Inhibition of ACE results in decreased plasma angiotensin II, which leads to decreased vasopressor activity and to decreased aldosterone secretion. The latter decrease may result in a small increase of serum potassium. Hypertensive patients treated with benazepril hydrochloride alone for up to 52 weeks had elevations of serum potassium of up to 0.2 mEq/L. Similar patients treated with benazepril hydrochloride and hydrochlorothiazide for up to 24 weeks had no consistent changes in their serum potassium (see **PRECAUTIONS**).

Removal of angiotensin II negative feedback on renin secretion leads to increased plasma renin activity. In animal studies, benazepril had no inhibitory effect on the vasopressor response to angiotensin II and did not interfere with the hemodynamic effects of the autonomic neurotransmitters acetylcholine, epinephrine, and norepinephrine.

ACE is identical to kininase, an enzyme that degrades bradykinin. Whether increased levels of bradykinin, a potent vasodepressor peptide, play a role in the therapeutic effects of benazepril hydrochloride remains to be elucidated.

While the mechanism through which benazepril lowers blood pressure is believed to be primarily suppression of the reninangiotensin-aldosterone system, benazepril has an antihypertensive effect even in patients with low-renin hypertension (see **INDICATIONS AND USAGE**).

Pharmacokinetics and Metabolism

Following oral administration of benazepril hydrochloride, peak plasma concentrations of benazepril are reached within 0.5 to 1 hours. The extent of absorption is at least 37% as determined by urinary recovery and is not significantly influenced by the presence of food in the GI tract.

Cleavage of the ester group (primarily in the liver) converts benazepril to its active metabolite, benazeprilat. Peak plasma concentrations of benazeprilat are reached 1 to 2 hours after drug intake in the fasting state and 2 to 4 hours after drug intake in the nonfasting state. The serum protein binding of benazepril is about 96.7% and that of benazeprilat about 95.3%, as measured by equilibrium dialysis; on the basis of *in vitro* studies, the degree of protein binding should be unaffected by age, hepatic dysfunction, or concentration (over the concentration range of 0.24 to 23.6 µmol/L).

Benazepril is almost completely metabolized to benazeprilat, which has much greater ACE inhibitory activity than benazepril, and to the glucuronide conjugates of benazepril and benazeprilat. Only trace amounts of an administered dose of benazepril hydrochloride can be recovered in the urine as unchanged benazepril, while about 20% of the dose is excreted as benazeprilat, 4% as benazepril glucuronide, and 8% as benazeprilat glucuronide.

The kinetics of benazepril are approximately dose-proportional within the dosage range of 10 to 80 mg.

In adults, the effective half-life of accumulation of benazeprilat following multiple dosing of benazepril hydrochloride is 10 to 11 hours. Thus, steady state concentrations of benazeprilat should be reached after 2 or 3 doses of benazepril hydrochloride given once daily.

The kinetics did not change, and there was no significant accumulation during chronic administration (28 days) of once-daily doses between 5 mg and 20 mg. Accumulation ratios based on AUC and urinary recovery of benazeprilat were 1.19 and 1.27, respectively. Benazepril and benazeprilat are cleared predominantly by renal excretion in healthy subjects with normal renal function. Nonrenal (i.e., biliary) excretion accounts for approximately 11% to 12% of benazeprilat excretion in healthy subjects. In patients with renal failure, biliary clearance may compensate to an extent for deficient renal clearance.

In patients with renal insufficiency, the disposition of benazepril and benazeprilat in patients with mild-to-moderate renal insufficiency (creatinine clearance > 30 mL/min) is similar to that in patients with normal renal function. In patients with creatinine clearance \leq 30 mL/min, peak benazeprilat levels and the initial (alpha phase) half-life increase, and time to steady state may be delayed (see **DOSAGE AND ADMINISTRATION**).

When dialysis was started 2 hours after ingestion of 10 mg of benazepril, approximately 6% of benazeprilat was removed in 4 hours of dialysis. The parent compound, benazepril, was not detected in the dialysate.

In patients with hepatic insufficiency (due to cirrhosis), the pharmacokinetics of benazeprilat are essentially unaltered. The pharmacokinetics of benazepril and benazeprilat do not appear to be influenced by age.

In pediatric patients, (N = 45) hypertensive, age 6 to 16 years, given multiple daily doses of benazepril hydrochloride (0.1 to 0.5 mg/kg), the clearance of benazeprilat for children 6 to 12 years old was 0.35 L/hr/kg, more than twice that of healthy adults receiving a single dose of 10 mg (0.13 L/hr/kg). In adolescents, it was 0.17 L/hr/kg, 27% higher than that of healthy adults. The terminal elimination half-life of benazeprilat in pediatric patients was around 5 hours, one-third that observed in adults.

Pharmacodynamics

Single and multiple doses of 10 mg or more of benazepril hydrochloride cause inhibition of plasma ACE activity by at least 80% to 90% for at least 24 hours after dosing. Pressor responses to exogenous angiotensin I were inhibited by 60% to 90% (up to 4 hours post-dose) at the 10 mg dose.

Clinical Studies

Hypertension

Adult

In single-dose studies, benazepril hydrochloride lowered blood pressure within 1 hour, with peak reductions achieved 2 to 4 hours after dosing. The antihypertensive effect of a single dose persisted for 24 hours. In multiple-dose studies, once-daily doses of 20 to 80 mg decreased seated pressure (systolic/diastolic) 24 hours after dosing by about 6 to 12/4 to 7 mmHg. The trough values represent reductions of about 50% of that seen at peak.

Four dose-response studies using once-daily dosing were conducted in 470 mild-to-moderate hypertensive patients not using diuretics. The minimal effective once-daily dose of benazepril hydrochloride was 10 mg; but further falls in blood pressure, especially at morning trough, were seen with higher doses in the studied dosing range (10 to 80 mg). In studies comparing the same daily dose of benazepril hydrochloride given as a single morning dose or as a twice-daily dose, blood pressure reductions at the time of morning trough blood levels were greater with the divided regimen.

The antihypertensive effects of benazepril hydrochloride were not appreciably different in patients receiving high- or low-sodium diets.

In normal human volunteers, single doses of benazepril caused an increase in renal blood flow but had no effect on glomerular filtration rate.

Use of benazepril hydrochloride in combination with thiazide diuretics gives a blood-pressure-lowering effect greater than that seen with either agent alone. By blocking the renin-angiotensin-aldosterone axis, administration of benazepril hydrochloride tends to reduce the potassium loss associated with the diuretic.

Pediatric

In a clinical study of 107 pediatric patients, 7 to 16 years of age, with either systolic or diastolic pressure above the 95th percentile, patients were given 0.1 or 0.2 mg/kg then titrated up to 0.3 or 0.6 mg/kg with a maximum dose of 40 mg once daily. After four weeks of treatment, the 85 patients whose blood pressure was reduced on therapy were then randomized to either placebo or benazepril

and were followed up for an additional two weeks. At the end of two weeks, blood pressure (both systolic and diastolic) in children withdrawn to placebo rose by 4 to 6 mmHg more than in children on benazepril. No dose-response was observed for the three doses.

INDICATIONS AND USAGE

Benazepril hydrochloride tablets USP are indicated for the treatment of hypertension. They may be used alone or in combination with thiazide diuretics.

CONTRAINDICATIONS

Benazepril hydrochloride tablets are contraindicated in patients who are hypersensitive to this product or to any other ACE inhibitor. Benazepril hydrochloride tablets are also contraindicated in patients with a history of angioedema with or without previous ACE inhibitor treatment.

Do not coadminister aliskiren with angiotensin receptor blockers, ACE inhibitors, including benazepril hydrochloride tablets in patients with diabetes.

WARNINGS

Anaphylactoid and Possibly Related Reactions

Presumably because angiotensin-converting enzyme inhibitors affect the metabolism of eicosanoids and polypeptides, including endogenous bradykinin, patients receiving ACE inhibitors (including benazepril hydrochloride) may be subject to a variety of adverse reactions, some of them serious.

Head and Neck Angioedema

Angioedema of the face, extremities, lips, tongue, glottis, and larynx has been reported in patients treated with angiotensin-converting enzyme inhibitors. In U.S. clinical trials, symptoms consistent with angioedema were seen in none of the subjects who received placebo and in about 0.5% of the subjects who received benazepril hydrochloride. Angioedema associated with laryngeal edema can be fatal. If laryngeal stridor or angioedema of the face, tongue, or glottis occurs, treatment with benazepril hydrochloride should be discontinued and appropriate therapy instituted immediately. Where there is involvement of the tongue, glottis, or larynx, likely to cause airway obstruction, appropriate therapy, e.g., subcutaneous epinephrine injection 1:1000 (0.3 mL to 0.5 mL) should be promptly administered (see ADVERSE REACTIONS).

Black patients receiving ACE inhibitors have been reported to have a higher incidence of angioedema compared to nonblacks.

Intestinal Angioedema

Intestinal angioedema has been reported in patients treated with ACE inhibitors. These patients presented with abdominal pain (with or without nausea or vomiting); in some cases there was no prior history of facial angioedema and C-1 esterase levels were normal. The angioedema was diagnosed by procedures including abdominal CT scan or ultrasound, or at surgery, and symptoms resolved after stopping the ACE inhibitor. Intestinal angioedema should be included in the differential diagnosis of patients on ACE inhibitors presenting with abdominal pain.

Anaphylactoid Reactions During Desensitization

Two patients undergoing desensitizing treatment with hymenoptera venom while receiving ACE inhibitors sustained life-threatening anaphylactoid reactions. In the same patients, these reactions were avoided when ACE inhibitors were temporarily withheld, but they reappeared upon inadvertent rechallenge.

Anaphylactoid Reactions During Membrane Exposure

Anaphylactoid reactions have been reported in patients dialyzed with high-flux membranes and treated concomitantly with an ACE inhibitor. Anaphylactoid reactions have also been reported in patients undergoing low-density lipoprotein apheresis with dextran sulfate absorption (a procedure dependent upon devices not approved in the United States).

Hypotension

Benazepril hydrochloride can cause symptomatic hypotension. Like other ACE inhibitors, benazepril has been only rarely associated with hypotension in uncomplicated hypertensive patients. Symptomatic hypotension is most likely to occur in patients who have been volume- and/or salt-depleted as a result of prolonged diuretic therapy, dietary salt restriction, dialysis, diarrhea, or vomiting. Volume- and/or salt-depletion should be corrected before initiating therapy with benazepril hydrochloride.

In patients with congestive heart failure, with or without associated renal insufficiency, ACE inhibitor therapy may cause excessive hypotension, which may be associated with oliguria or azotemia and, rarely, with acute renal failure and death. In such patients, benazepril hydrochloride therapy should be started under close medical supervision; they should be followed closely for the first 2 weeks of treatment and whenever the dose of benazepril or diuretic is increased.

If hypotension occurs, the patient should be placed in a supine position, and, if necessary, treated with intravenous infusion of physiological saline. Benazepril hydrochloride treatment usually can be continued following restoration of blood pressure and volume.

Fetal Toxicity

Pregnancy Category D

Use of drugs that act on the renin-angiotensin system during the second and third trimesters of pregnancy reduces fetal renal function and increases fetal and neonatal morbidity and death. Resulting oligohydramnios can be associated with fetal lung hypoplasia and skeletal deformations. Potential neonatal adverse effects include skull hypoplasia, anuria, hypotension, renal failure, and death. When pregnancy is detected, discontinue benazepril hydrochloride as soon as possible. These adverse outcomes are usually associated with use of these drugs in the second and third trimester of pregnancy. Most epidemiologic studies examining fetal abnormalities after exposure to antihypertensive use in the first trimester have not distinguished drugs affecting the renin-angiotensin system from other antihypertensive agents. Appropriate management of maternal hypertension during pregnancy is important to optimize outcomes for both mother and fetus.

In the unusual case that there is no appropriate alternative to therapy with drugs affecting the renin-angiotensin system for a particular patient, apprise the mother of the potential risk to the fetus. Perform serial ultrasound examinations to assess the intra-amniotic environment. If oligohydramnios is observed, discontinue benazepril hydrochloride, unless it is considered lifesaving for the mother. Fetal testing may be appropriate, based on the week of pregnancy. Patients and physicians should be aware, however, that oligohydramnios may not appear until after the fetus has sustained irreversible injury. Closely observe infants with histories of *in utero* exposure to benazepril hydrochloride for hypotension, oliguria, and hyperkalemia (see **PRECAUTIONS, Pediatric Use**).

No teratogenic effects of benazepril hydrochloride were seen in studies of pregnant rats, mice, and rabbits. On a mg/m^2 basis, the doses used in these studies were 60 times (in rats), 9 times (in mice), and more than 0.8 times (in rabbits) the maximum recommended human dose (assuming a 50 kg woman). On a mg/kg basis these multiples are 300 times (in rats), 90 times (in mice), and more than 3 times (in rabbits) the maximum recommended human dose.

Hepatic Failure

Rarely, ACE inhibitors have been associated with a syndrome that starts with cholestatic jaundice and progresses to fulminant hepatic necrosis and (sometimes) death. The mechanism of this syndrome is not understood. Patients receiving ACE inhibitors who develop jaundice or marked elevations of hepatic enzymes should discontinue the ACE inhibitor and receive appropriate medical follow-up.

PRECAUTIONS

General

Impaired Renal Function

As a consequence of inhibiting the renin-angiotensin-aldosterone system, changes in renal function may be anticipated in susceptible individuals. In patients with severe congestive heart failure whose renal function may depend on the activity of the renin-angiotensin-aldosterone system, treatment with angiotensin-converting enzyme inhibitors, including benazepril hydrochloride, may be associated with oliguria and/or progressive azotemia and (rarely) with acute renal failure and/or death. In a small study of hypertensive patients with renal artery stenosis in a solitary kidney or bilateral renal artery stenosis, treatment with benazepril hydrochloride was associated with increases in blood urea nitrogen and serum creatinine; these increases were reversible upon discontinuation of benazepril hydrochloride or diuretic therapy, or both. When such patients are treated with ACE inhibitors, renal function should be monitored during the first few weeks of therapy. Some hypertensive patients with no apparent preexisting renal vascular disease have developed increases in blood urea nitrogen and serum creatinine, usually minor and transient, especially when benazepril hydrochloride has been given concomitantly with a diuretic. This is more likely to occur in patients with preexisting renal impairment. Dosage reduction of benazepril hydrochloride and/or discontinuation of the diuretic may be required. **Evaluation of the hypertensive patient should always include assessment of renal function (see DOSAGE AND ADMINISTRATION).**

Hyperkalemia

In clinical trials, hyperkalemia (serum potassium at least 0.5 mEq/L greater than the upper limit of normal) occurred in approximately 1% of hypertensive patients receiving benazepril hydrochloride. In most cases, these were isolated values which resolved despite continued therapy. Risk factors for the development of hyperkalemia include renal insufficiency, diabetes mellitus, and the concomitant use of potassium-sparing diuretics, potassium supplements, and/or potassium-containing salt substitutes, which should be used cautiously, if at all, with benazepril hydrochloride (see **Drug Interactions**).

Cough

Presumably due to the inhibition of the degradation of endogenous bradykinin, persistent nonproductive cough has been reported with all ACE inhibitors, always resolving after discontinuation of therapy. ACE inhibitor-induced cough should be considered in the differential diagnosis of cough.

Surgery/Anesthesia

In patients undergoing surgery or during anesthesia with agents that produce hypotension, benazepril will block the angiotensin II formation that could otherwise occur secondary to compensatory renin release. Hypotension that occurs as a result of this mechanism can be corrected by volume expansion.

Information for Patients

Pregnancy

Female patients of childbearing age should be told about the consequences of exposure to benazepril hydrochloride during pregnancy. Discuss treatment options with women planning to become pregnant. Patients should be asked to report pregnancies to their physicians as soon as possible.

Angioedema

Angioedema, including laryngeal edema, can occur at any time with treatment with ACE inhibitors. Patients should be so advised and told to report immediately any signs or symptoms suggesting angioedema (swelling of face, eyes, lips, or tongue, or difficulty in breathing) and to take no more drug until they have consulted with the prescribing physician.

Symptomatic Hypotension

Patients should be cautioned that lightheadedness can occur, especially during the first days of therapy, and it should be reported to the prescribing physician. Patients should be told that if syncope occurs, benazepril hydrochloride should be discontinued until the prescribing physician has been consulted.

All patients should be cautioned that inadequate fluid intake or excessive perspiration, diarrhea, or vomiting can lead to an excessive fall in blood pressure, with the same consequences of lightheadedness and possible syncope.

Hyperkalemia

Patients should be told not to use potassium supplements or salt substitutes containing potassium without consulting the prescribing physician.

Neutropenia

Patients should be told to promptly report any indication of infection (e.g., sore throat, fever), which could be a sign of neutropenia.

Drug Interactions

Diuretics

Patients on diuretics, especially those in whom diuretic therapy was recently instituted, may occasionally experience an excessive reduction of blood pressure after initiation of therapy with benazepril hydrochloride. The possibility of hypotensive effects with benazepril hydrochloride can be minimized by either discontinuing the diuretic or increasing the salt intake prior to initiation of treatment with benazepril hydrochloride. If this is not possible, the starting dose should be reduced (see **DOSAGE AND ADMINISTRATION**).

Potassium Supplements and Potassium-Sparing Diuretics Concomitant use with benazepril hydrochloride may effect potassium levels. Monitor potassium periodically.

Oral Anticoagulants

Interaction studies with warfarin and acenocoumarol failed to identify any clinically important effects on the serum concentrations or clinical effects of these anticoagulants.

Lithium

Increased serum lithium levels and symptoms of lithium toxicity have been reported in patients receiving ACE inhibitors (including benazepril) during therapy with lithium. Monitor lithium levels when used concomitantly with benazepril hydrochloride.

Gold

Nitritoid reactions (symptoms include facial flushing, nausea, vomiting and hypotension) have been reported rarely in patients on therapy with injectable gold (sodium aurothiomalate) and concomitant ACE inhibitor therapy.

Anti-Diabetics

In rare cases, diabetic patients receiving an ACE inhibitor (including benazepril) concomitantly with insulin or oral anti-diabetics may develop hypoglycemia. Such patients should therefore be advised about the possibility of hypoglycemic reactions and should be monitored accordingly.

Non-Steroidal Anti-Inflammatory Drugs (NSAIDs) Including Selective Cyclooxygenase-2 Inhibitors (COX-2 Inhibitors) In patients who are elderly, volume-depleted (including those on diuretic therapy), or with compromised renal function, coadministration of NSAIDs, including selective COX-2 inhibitors, with ACE inhibitors, including benazepril, may result in deterioration of renal function, including possible acute renal failure. These effects are usually reversible. Monitor renal function periodically in patients receiving benazepril and NSAID therapy.

The antihypertensive effect of ACE inhibitors, including benazepril, may be attenuated by NSAIDs.

Dual Blockade of the Renin-Angiotensin System (RAS)

Dual blockade of the RAS with angiotensin receptor blockers, ACE inhibitors, or aliskiren is associated with increased risks of hypotension, hyperkalemia, and changes in renal function (including acute renal failure) compared to monotherapy. Closely monitor blood pressure, renal function and electrolytes in patients on benazepril hydrochloride and other agents that block the RAS.

Do not coadminister aliskiren with benazepril hydrochloride in patients with diabetes. Avoid use of aliskiren with benazepril hydrochloride in patients with renal impairment (GFR < 60 ml/min).

Other

Benazepril hydrochloride has been used concomitantly with beta-adrenergic-blocking agents, calcium-channel-blocking agents, diuretics, digoxin, and hydralazine, without evidence of clinically important adverse interactions. Benazepril, like other ACE inhibitors, has had less than additive effects with beta-adrenergic blockers, presumably because both drugs lower blood pressure by inhibiting parts of the renin-angiotensin system.

The pharmacokinetics of benazepril are not affected by the following drugs: hydrochlorothiazide, furosemide, chlorthalidone, digoxin, propranolol, atenolol, nifedipine, amlodipine, naproxen, acetylsalicylic acid, or cimetidine. Likewise the administration of benazepril does not substantially affect the pharmacokinetics of these medications (cimetidine kinetics were not studied).

Carcinogenesis, Mutagenesis, Impairment of Fertility

No evidence of carcinogenicity was found when benazepril was administered to rats and mice for up to two years at doses of up to 150 mg/kg/day. When compared on the basis of body weights, this dose is 110 times the maximum recommended human dose. When compared on the basis of body surface areas, this dose is 18 and 9 times (rats and mice, respectively) the maximum recommended human dose (calculations assume a patient weight of 60 kg). No mutagenic activity was detected in the Ames test in bacteria (with or without metabolic activation), in an *in vitro* test for forward mutations in cultured mammalian cells, or in a nucleus anomaly test.

In doses of 50 to 500 mg/kg/day (6 to 60 times the maximum recommended human dose based on mg/m² comparison and 37 to 375 times the maximum recommended human dose based on a mg/kg comparison), benazepril hydrochloride had no adverse effect on the reproductive performance of male and female rats.

Nursing Mothers

Minimal amounts of unchanged benazepril and of benazeprilat are excreted into the breast milk of lactating women treated with benazepril. A newborn child ingesting entirely breast milk would receive less than 0.1% of the mg/kg maternal dose of benazepril and benazeprilat.

Geriatric Use

Of the total number of patients who received benazepril in U.S. clinical studies of benazepril hydrochloride, 18% were 65 or older while 2% were 75 or older. No overall differences in effectiveness or safety were observed between these patients and younger patients, and other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out.

Benazepril and benazeprilat are substantially excreted by the kidney. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

Pediatric Use

Neonates With a History of In Utero Exposure to Benazepril Hydrochloride

If oliguria or hypotension occurs, direct attention toward support of blood pressure and renal perfusion. Exchange transfusions or dialysis may be required as a means of reversing hypotension and/or substituting for disordered renal function. Benazepril, which crosses the placenta, can theoretically be removed from the neonatal circulation by these means; there are occasional reports of benefit from these maneuvers with another ACE inhibitor, but experience is limited.

The antihypertensive effects of benazepril hydrochloride have been evaluated in a double-blind study in pediatric patients 7 to 16 years of age (see **CLINICAL PHARMACOLOGY**, **Pharmacodynamics**, *Clinical Studies*, <u>Hypertension</u>). The pharmacokinetics of benazepril hydrochloride have been evaluated in pediatric patients 6 to 16 years of age (see **CLINICALPHARMACOLOGY**, **Pharmacokinetics and Metabolism**). Benazepril hydrochloride was generally well tolerated and adverse effects were similar to

those described in adults (see **ADVERSE REACTIONS**, **Pediatric Patients**). The long-term effects of benazepril on growth and development have not been studied. Infants below the age of 1 year should not be given benazepril hydrochloride because of the risk of effects on kidney development.

Treatment with benazepril hydrochloride is not recommended in pediatric patients less than 6 years of age (see **ADVERSE REACTIONS**), and in children with glomerular filtration rate < 30 mL/min as there are insufficient data available to support a dosing recommendation in these groups (see **CLINICAL PHARMACOLOGY**, **Pharmacokinetics and Metabolism**, *In pediatric patients* and **DOSAGE AND ADMINISTRATION**).

ADVERSE REACTIONS

Benazepril hydrochloride has been evaluated for safety in over 6000 patients with hypertension; over 700 of these patients were treated for at least one year. The overall incidence of reported adverse events was comparable in benazepril hydrochloride and placebo patients.

The reported side effects were generally mild and transient, and there was no relation between side effects and age, duration of therapy, or total dosage within the range of 2 to 80 mg. Discontinuation of therapy because of a side effect was required in approximately 5% of U.S. patients treated with benazepril hydrochloride and in 3% of patients treated with placebo.

The most common reasons for discontinuation were headache (0.6%) and cough (0.5%) (see **PRECAUTIONS**, *Cough*).

The side effects considered possibly or probably related to study drug that occurred in U.S. placebo-controlled trials in more than 1% of patients treated with benazepril hydrochloride are shown below.

PATIENTS IN U.S. PLACEBO-CONTROLLED STUDIES

| | BENAZ HYDROCI | ZEPRIL HLORIDE | PLACEBO | |
|--------------------|------------------|-------------------|-----------|-----|
| | (N = 964) | | (N = 496) | |
| | | | | |
| | N | % | N | % |
| Headache | 60 | 6.2 | 21 | 4.2 |
| Dizziness | 35 | 3.6 | 12 | 2.4 |
| Somnolence | 15 | 1.6 | 2 | 0.4 |
| Postural Dizziness | 14 | 1.5 | 1 | 0.2 |

Other adverse experiences reported in controlled clinical trials (in less than 1% of benazepril patients or with less than 1% difference in incidence between benazepril or placebo treatment), and rarer events seen in postmarketing experience, include the following (in some, a causal relationship to drug use is uncertain):

Dermatologic

Stevens-Johnson syndrome, pemphigus, apparent hypersensitivity reactions (manifested by dermatitis, pruritus, or rash), photosensitivity, and flushing.

Gastrointestinal

Nausea, pancreatitis, constipation, gastritis, vomiting, and melena.

Hematologic

Thrombocytopenia and hemolytic anemia.

Neurologic and Psychiatric

Anxiety, decreased libido, hypertonia, insomnia, nervousness, and paresthesia.

Other

Fatigue, asthma, bronchitis, dyspnea, sinusitis, urinary tract infection, frequent urination, infection, arthritis, impotence, alopecia, arthralgia, myalgia, asthenia, and sweating.

Another potentially important adverse experience, eosinophilic pneumonitis, has been attributed to other ACE inhibitors.

Pediatric Patients

The adverse experience profile for pediatric patients appears to be similar to that seen in adult patients.

Clinical Laboratory Test Findings

Hemoglobin

Decreases in hemoglobin (a low value and a decrease of 5 g/dL) were rare, occurring in only 1 of 2,014 patients receiving benazepril hydrochloride alone and in 1 of 1,357 patients receiving benazepril hydrochloride plus a diuretic. No U.S. patients discontinued treatment because of decreases in hemoglobin.

Other (Causal Relationships Unknown)

Elevations of uric acid, blood glucose, serum bilirubin, and liver enzymes (see **WARNINGS**) have been reported, as have scattered incidents of hyponatremia, electrocardiographic changes, eosinophilia, and proteinuria.

OVERDOSAGE

Single oral doses of 3 g/kg benazepril were associated with significant lethality in mice. Rats, however, tolerated single oral doses of up to 6 g/kg. Reduced activity was seen at 1 g/kg in mice and at 5 g/kg in rats. Human overdoses of benazepril have not been reported, but the most common manifestation of human benazepril overdosage is likely to be hypotension, which can be associated with electrolyte disturbances and renal failure.

Laboratory determinations of serum levels of benazepril and its metabolites are not widely available, and such determinations have, in any event, no established role in the management of benazepril overdose.

No data are available to suggest physiological maneuvers (e.g., maneuvers to change the pH of the urine) that might accelerate elimination of benazepril and its metabolites. Benazepril is only slightly dialyzable, but dialysis might be considered in overdosed patients with severely impaired renal function (see **WARNINGS**).

Angiotensin II could presumably serve as a specific antagonist-antidote in the setting of benazepril overdose, but angiotensin II is essentially unavailable outside of scattered research facilities. Because the hypotensive effect of benazepril is achieved through vasodilation and effective hypovolemia, it is reasonable to treat benazepril overdose by infusion of normal saline solution. If ingestion is recent, activated charcoal should be considered. Gastric decontamination (e.g., vomiting, gastric lavage) may be considered in individual cases, in the early period after ingestion.

Patients should be closely monitored for blood pressure and clinical symptoms. Supportive management should be employed to ensure adequate hydration and to maintain systemic blood pressure.

In the case of marked hypotension, physiological saline solution should be administered intravenously; depending on the clinical situation the use of vasopressors (e.g., catecholamines i.v.) may be considered.

DOSAGE AND ADMINISTRATION

Hypertension

Adults

The recommended initial dose for patients not receiving a diuretic is 10 mg once a day. The usual maintenance dosage range is 20 to 40 mg per day administered as a single dose or in two equally divided doses. A dose of 80 mg gives an increased response, but experience with this dose is limited. The divided regimen was more effective in controlling trough (pre-dosing) blood pressure than the same dose given as a once-daily regimen. Dosage adjustment should be based on measurement of peak (2 to 6 hours after dosing) and trough responses. If a once-daily regimen does not give adequate trough response, an increase in dosage or divided administration should be considered. If blood pressure is not controlled with benazepril hydrochloride tablets alone, a diuretic can be added.

Total daily doses above 80 mg have not been evaluated.

Concomitant administration of benazepril hydrochloride tablets with potassium supplements, potassium salt substitutes, or potassiumsparing diuretics can lead to increases of serum potassium (see **PRECAUTIONS**).

In patients who are currently being treated with a diuretic, symptomatic hypotension occasionally can occur following the initial dose of benazepril hydrochloride tablets. To reduce the likelihood of hypotension, the diuretic should, if possible, be discontinued two to three days prior to beginning therapy with benazepril hydrochloride tablets (see **WARNINGS**). Then, if blood pressure is not controlled with benazepril hydrochloride tablets alone, diuretic therapy should be resumed.

If the diuretic cannot be discontinued, an initial dose of 5 mg benazepril hydrochloride tablets should be used to avoid excessive hypotension.

Pediatrics

In children, doses of benazepril hydrochloride tablets between 0.1 and 0.6 mg/kg once daily have been studied, and doses greater than 0.1 mg/kg were shown to reduce blood pressure (see **Pharmacodynamics**). Based on this, the recommended starting dose of benazepril hydrochloride tablets is 0.2 mg/kg once per day as monotherapy. Doses above 0.6 mg/kg (or in excess of 40 mg daily) have not been studied in pediatric patients.

For pediatric patients who cannot swallow tablets, or for whom the calculated dosage (mg/kg) does not correspond to the available tablet strengths for benazepril hydrochloride tablets, follow the suspension preparation instructions below to administer benazepril HCl as a suspension.

Treatment with benazepril hydrochloride tablets is not advised for children below the age of 6 years (see **PRECAUTIONS**, **Pediatric Use**) and in pediatric patients with glomerular filtration rate < 30 mL, as there are insufficient data available to support a dosing recommendation in these groups.

For Hypertensive Patients With Renal Impairment

For patients with a creatinine clearance $< 30 \text{ mL/min/1.73 m}^2$ (serum creatinine > 3 mg/dL), the recommended initial dose is 5 mg benazepril hydrochloride tablets once daily. Dosage may be titrated upward until blood pressure is controlled or to a maximum total daily dose of 40 mg (see **WARNINGS**).

Preparation of Suspension (for 150 mL of a 2 mg/mL Suspension)

Add 75 mL of Ora-Plus^{®*} oral suspending vehicle to an amber polyethylene terephthalate (PET) bottle containing fifteen benazepril hydrochloride tablets, 20 mg, and shake for at least 2 minutes. Allow the suspension to stand for a minimum of 1 hour. After the standing time, shake the suspension for a minimum of 1 additional minute. Add 75 mL of Ora-Sweet^{®*} oral syrup vehicle to the bottle and shake the suspension to disperse the ingredients. The suspension should be refrigerated at 2 to 8°C (36 to 46°F) and can be stored for up to 30 days in the PET bottle with a child-resistant screw-cap closure. Shake the suspension before each use.

*Ora-Plus[®] and Ora-Sweet[®] are registered trademarks of Paddock Laboratories, Inc. Ora-Plus[®] contains carrageenan, citric acid, methylparaben, microcrystalline cellulose, carboxymethylcellulose sodium, potassium sorbate, simethicone, sodium phosphate monobasic, xanthan gum, and water. Ora-Sweet[®] contains citric acid, berry citrus flavorant, glycerin, methylparaben, potassium sorbate, sodium phosphate monobasic, sorbitol, sucrose, and water.

HOW SUPPLIED

Benazepril hydrochloride tablets USP, 5 mg are light yellow, arc triangle coated tablets, debossed with the number "93" on one side and "5124" on the other. Tablets are packaged in bottles of 100.

Benazepril hydrochloride tablets USP, 10 mg are mustard yellow, arc triangle coated tablets, debossed with the number "93" on one side and "5125" on the other. Tablets are packaged in bottles of 100 and 500.

Benazepril hydrochloride tablets USP, 20 mg are pink, arc triangle coated tablets, debossed with the number "93" on one side and "5126" on the other. Tablets are packaged in bottles of 100 and 500.

Benazepril hydrochloride tablets USP, 40 mg are pink to light red, arc triangle coated tablets, debossed with the number "93" on one side and "5127" on the other. Tablets are packaged in bottles of 100.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Dispense in a tight, light-resistant container as defined in the USP, with a child-resistant closure (as required).

KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

Manufactured In India By:

EMCURE PHARMACEUTICALS LTD.

Hinjwadi, Pune, India Manufactured For: **TEVA PHARMACEUTICALS USA** Sellersville, PA 18960 Rev. P 9/2012

PRINCIPAL DISPLAY PANEL

| NDC 0093-5124-01 BENAZEPRIL HYDROCHLORIDE Tablets USP 5 mg Pc only 100 TABLETS | Each tablet contains 5 mg benazepril Nydrochloride, USP. | Jsual Dosage: See package insert for 'ull prescribing information. Store at 20° to 25°C (68° to 77°F) [See JSP Controlled Room Temperature]. | Dispense in a tight, light-resistant container as defined in the USP, with a child-resistant closure (as required). KEEP THIS AND ALL MEDICATIONS DUT OF THE REACH OF CHILDREN. | Rev. B 7/2008 Manufactured In India By: EMCURE PHARMACEUTICALS LTD. Hinjwadi, Pune, India Manufactured For: Manufactured For: Seller sville, PA 18960 | Code No.: MH/DRUGS/PD/149 ⊶ ≂ | LOT: EXP: | × 5102700501N02 |
|--|--|---|---|--|---|--|---|
| | Eac | Usi Sto USI | S C S S S S S S S S S S S S S S S S S S | Hin Ma Sel Ain Sel Ain | 3 | 0093-5124-01 | 210 |
| | NDC 0093-5124-01 BENAZEPRIL HYDROCHLORIDE Tablets USP 5 mg Be only 100 TABLETS | NDC 0093-5124-01 BENAZEPRIL HYDROCHLORIDE Tablets USP 5 mg Pc only 100 TABLETS | NDC 0093-5124-01 Each tablet contains 5 mg benazepril hydrochloride, USP. Store at 20° to 25°C (68° to 77°F) [See USP Controlled Boom Temperature]. | NDC 0003-2154-01 Each tablet contains 5 mg benazepril hydrochloride, USP. Daval Dosage: See package insert for full prescribing information. Store at 20° to 25°C (8°° to 77°F) [See USP controlled Room Temperature]. Dispense in a tight, light-resistant container as defined in the USP, with a child-resistant closure (as required). KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN. | NDC 0003-2154-01 Each tablet contains 5 mg benazepril hydrochloride, USP. Usual Dosage: See package insert for full prescribing information. Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature]. Dispense in a tight, light-resistant container as defined in the USP, with a container as defined i | NDC 0003-2154-01 Each tablet contains 5 mg benazepril hydrochloride, USP. Each tablet contains 5 mg benazepril bydrochloride, USP. BENVAZEBLUR Usual Dosage: See package insert for full prescribing information. Benazepril Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature]. Dispense in a tight, light-resistant container as defined in the USP, with a child-resistant closure (as required). REEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN. Rev. B 7/2008 Manufactured In India By: EMCURE PHARMACEUTICALS LTD. Rev. B 7/2008 Code No.: MH/DRUGS/PD/149 Code No.: MH/DRUGS/PD/149 | MDC 0003-5124-01 Each tablet contains 5 mg benazepril for 0.0000 for the factor tablet contains 5 mg benazepril for 0.0000 for the factor of the factor tablet contains 5 mg benazepril for 0.0000 for the factor of the fact |

Benazepril Hydrochloride Tablets USP 5 mg 100s Label Text NDC 0093-5124-01 BENAZEPRIL HYDROCHLORIDE Tablets USP 5 mg Rx only 100 TABLETS TEVA

PRINCIPAL DISPLAY PANEL

| NDC 0093-5125-01 BENAZEPRIL HYDROCHLORIDE Tablets USP 10 mg By only 100 TABLETS | ablet contains 10 mg benazepril thloride, USP. Dosage : See package insert for scribing information. At 20° to 25°C (68° to 77°F) [See ontrolled Room Temperature]. It 20° to 25°C (68° to 77°F) [See ontrolled Room Temperature]. It a tight, light-resistant er as defined in the USP, with a seistant closure (as required). THIS AND ALL MEDICATIONS FTHE REACH OF CHILDREN. Rev. B 772008 sistant closure (as required). THIS AND ALL MEDICATIONS FTHE REACH OF CHILDREN. Rev. B 772008 actured for: PHARMACEUTICALS USA sville, PA 18960 Code No.: MH/DRUGS/PD/149 | LOT: EXP: |
|---|--|------------------|
| TEVA | Each t hydroc Usual Usual Usper contair child- | N 0093-5125-01 5 |

Benazepril Hydrochloride Tablets USP 10 mg 100s Label Text NDC 0093-5125-01 BENAZEPRIL HYDROCHLORIDE Tablets USP 10 mg Rx only 100 TABLETS TEVA

PRINCIPAL DISPLAY PANEL



Benazepril Hydrochloride Tablets USP 20 mg 100s Label Text NDC 0093-5126-01 BENAZEPRIL HYDROCHLORIDE Tablets USP 20 mg Rx only 100 TABLETS TEVA



Benazepril Hydrochloride Tablets USP 40 mg 100s Label Text NDC 0093-5127-01 BENAZEPRIL HYDROCHLORIDE Tablets USP 40 mg Rx only 100 TABLETS TEVA