DRUG FACTS

Active ingredient(s) ***Purpose***

Use(s) temporarily relieves:
- runny nose
- sneezing
- itching of the nose or throat and itchy, watery eyes due to hay fever
- headache
- minor aches and pains

Warnings

Do not use
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on skin

Ask a doctor before use if you have
- liver disease
- glaucoma
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are
- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if
- new symptoms occur
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling present

These could be signs of a serious condition.

If pregnant or breast-feeding

Keep out of reach of childrenOverdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
- **do not take more than directed (see overdose warning)**
- adults and children 12 years and over
- take 2 caplets every 4 to 6 hours
- do not take more than 8 caplets in 24 hours
- children under 12 years: do not use this adult product in children under 12 years of age; this will provide more than the recommended dose (overdose) and may cause liver damage

Other information This product contains acetaminophen. Severe liver damage may occur if you take:
- more than 8 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product
- store at controlled room temperature 15º-30ºC (59º-86ºF)
- see end flap for expiration date and lot number

Inactive ingredients carnauba wax, corn starch, croscarmellose sodium, crospovidone, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, povidone, silicon dioxide, stearic acid, titanium dioxide, triacetin