DRUG FACTS

Purpose Active ingredient(s)

Use(s)

- for prevention and treatment of these symptoms associated with motion sickness:
- nausea
- · vomiting
- dizziness

Warnings

for children under 12 years of age unless directed by a doctor

Ask a doctor before use if you have

- glaucoma
 difficulty in urination due to an enlargement of the prostate gland
 a breathing problems such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are

taking sedatives or tranqulizers

When using this product

- drowsiness may occur
 avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- · be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

If pregnant or breast-feeding

Keep out of reach of children
In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- · to prevent motion sickness, take the fitst dose one hour before starting activity
- to prevent or treat motion sickness: 1 to 2 tablets once daily for adults and children 12 years and over, or as directed by a doctor

Other information

- store at controlled room temperature 15°-30°C (59°-86°F)
- protect from heat and humidity
- used by expiration date on package

Inactive ingredients

corn starch, D&C yellow #10 aluminum lake, lactose, magnesium stearate, silica gel