

LEADER STOMACH RELIEF- bismuth subsalicylate liquid

Cardinal Health

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Cardinal Health Stomach Relief Drug Facts

Active ingredient (in each 15 mL Tablespoon)

Bismuth subsalicylate 262 mg

Purposes

Upset stomach reliever/antidiarrheal

Uses

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
- heartburn
- indigestion
- nausea
- gas
- belching
- fullness

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use

if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are

taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if

- symptoms get worse
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- shake well before use
- for accurate dosing, use dose cup
- adults and children 12 years and over: 1 dose (2 Tbsp or 30 mL) every 1/2 to 1 hour as needed
- do not exceed 8 doses (16 Tbsp or 240 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- each Tbsp contains: magnesium 5 mg, potassium 5 mg and sodium 5 mg
- salicylate 123 mg
- protect from freezing
- sugar free
- low sodium
- store at 20°-25°C (68°-77°F)
- does not meet USP requirements for pH
- for health information visit <http://www.more-info.info/>

Inactive ingredients

D&C red #22, D&C red #28, flavor, magnesium aluminum silicate, methylcellulose, purified water, saccharin sodium, salicylic acid, sodium salicylate, sorbic acid, xanthan gum

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to Original Pepto-Bismol® active ingredient

ORIGINAL STRENGTH

Stomach Relief

Upset Stomach Reliever/Antidiarrheal

Bismuth Subsalicylate 262 mg

Protective Coating Action

Soothing Relief for 5 Symptoms:

Heartburn

Indigestion

Upset Stomach

Nausea

Diarrhea

<p>Drug Facts (continued)</p> <p>Directions ■ shake well before use ■ for accurate dosing, use dose cup ■ adults and children 12 years and over: 1 dose (2 Tbsp or 30 mL) every 1/2 to 1 hour as needed ■ do not exceed 8 doses (16 Tbsp or 240 mL) in 24 hours ■ use until diarrhea stops but not more than 2 days ■ children under 12 years: ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea</p> <p>Other information ■ each Tbsp contains: magnesium 5 mg, potassium 5 mg and sodium 5 mg ■ salicylate 123 mg ■ protect from freezing ■ sugar free ■ low sodium ■ store at 20°-25°C (68°-77°F) ■ does not meet USP requirements for pH ■ for health information visit www.more-info.info</p> <p>Inactive ingredients D&C red #22, D&C red #28, flavor, magnesium aluminum silicate, methylcellulose, purified water, saccharin sodium, salicylic acid, sodium salicylate, sorbic acid, xanthan gum</p> <p>Questions or comments? 1-800-719-9260</p>	<p>NDC 37205-951-40</p> <p>LEADER®</p> <p>Compare to Original Pepto-Bismol® active ingredient*</p> <p>ORIGINAL STRENGTH</p> <h1>Stomach Relief</h1> <p>Upset Stomach Reliever/Antidiarrheal Bismuth Subsalicylate 262 mg</p> <p>Protective Coating Action</p> <p>Soothing Relief for 5 Symptoms: • Heartburn • Indigestion • Upset Stomach • Nausea • Diarrhea</p> <p>12 FL OZ (355 ml)</p> <p>SATISFACTORY GUARANTEED</p>	<p>Drug Facts</p> <table border="1"><thead><tr><th>Active ingredient (in each 15 mL Tablespoon)</th><th>Purposes</th></tr></thead><tbody><tr><td>Bismuth subsalicylate 262 mg.....</td><td>Upset stomach reliever/antidiarrheal</td></tr></tbody></table> <p>Uses relieves ■ travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence in food and drink, including: ■ heartburn ■ indigestion ■ nausea ■ gas ■ belching ■ fullness</p> <p>Warnings Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. Allergy alert: Contains salicylate. Do not take if you are ■ allergic to salicylates (including aspirin) ■ taking other salicylate products</p> <p>Do not use if you have ■ an ulcer ■ a bleeding problem ■ bloody or black stool</p> <p>Ask a doctor before use if you have ■ fever ■ mucus in the stool</p> <p>Ask a doctor or pharmacist before use if you are taking any drug for ■ anticoagulation (thinning the blood) ■ diabetes ■ gout ■ arthritis</p> <p>When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur</p> <p>Stop use and ask a doctor if ■ symptoms get worse ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)</p>	Active ingredient (in each 15 mL Tablespoon)	Purposes	Bismuth subsalicylate 262 mg.....	Upset stomach reliever/antidiarrheal
Active ingredient (in each 15 mL Tablespoon)	Purposes					
Bismuth subsalicylate 262 mg.....	Upset stomach reliever/antidiarrheal					

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING
*This product is not manufactured or distributed by Procter & Gamble, distributor of Original Pepto-Bismol®. **Gluten Free**

DISTRIBUTED BY
CARDINAL HEALTH
DUBLIN, OHIO 43017
CIN 4581229
www.myleader.com
1-800-200-6313

All Leader® Brand products are 100% satisfaction guaranteed or return to place of purchase for a full refund.

0 96295 12292 3

30240 E9 F6

bismuth subsalicylate liquid

Product Information

Product Type	HUMAN OTC DRUG LABEL	Item Code (Source)	NDC:37205-951
Route of Administration	ORAL	DEA Schedule	

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (BISMUTH CATION and SALICYLIC ACID)	BISMUTH SUBSALICYLATE	262 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
D&C RED NO. 22	
D&C RED NO. 28	
MAGNESIUM ALUMINUM SILICATE	
WATER	
SACCHARIN SODIUM	
SALICYLIC ACID	
SODIUM SALICYLATE	
SORBIC ACID	
XANTHAN GUM	

Product Characteristics

Color	PINK (viscous)	Score	
Shape		Size	
Flavor	MINT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37205-951-40	355 mL in 1 BOTTLE		
2	NDC:37205-951-34	237 mL in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part335	04/15/1990	

Labeler - Cardinal Health (097537435)