

GRISEOFULVIN- griseofulvin tablet, film coated
Oceanside Pharmaceuticals

Griseofulvin Ultramicrosize Tablets, USP

DESCRIPTION

Griseofulvin Ultramicrosize Tablets contain ultramicrosize crystals of griseofulvin, an antibiotic derived from a species of *Penicillium*.

Each Griseofulvin Ultramicrosize tablet contains:

Active Ingredient: griseofulvin ultramicrosize 125 mg

Inactive Ingredients: colloidal silicon dioxide, lactose, magnesium stearate, methylcellulose, methylparaben, polyethylene glycol 400 and 8000, povidone, and titanium dioxide.

OR

Active Ingredient: griseofulvin ultramicrosize 250 mg

Inactive Ingredients: colloidal silicon dioxide, magnesium stearate, methylcellulose, methylparaben, polyethylene glycol 400 and 8000, povidone, sodium lauryl sulfate, and titanium dioxide.

ACTION

Microbiology

Griseofulvin is fungistatic with *in vitro* activity against various species of *Microsporum*, *Epidermophyton* and *Trichophyton*. It has no effect on bacteria or other genera of fungi.

Pharmacokinetics

Following oral administration, griseofulvin is deposited in the keratin precursor cells and has a greater affinity for diseased tissue. The drug is tightly bound to the new keratin which becomes highly resistant to fungal invasions.

The efficiency of gastrointestinal absorption of ultramicrocrystalline griseofulvin is approximately one and one-half times that of the conventional microsize griseofulvin. This factor permits the oral intake of two-thirds as much ultramicrocrystalline griseofulvin as the microsize form. However, there is currently no evidence that this lower dose confers any significant clinical differences with regard to safety and/or efficacy.

In a bioequivalence study conducted in healthy volunteers (N=24) in the fasted state, 250 mg ultramicrocrystalline griseofulvin tablets were compared with 250 mg ultramicrocrystalline griseofulvin tablets that were physically altered (crushed) and administered with applesauce. The 250 mg ultramicrocrystalline griseofulvin tablets were found to be bioequivalent to the physically altered (crushed) 250 mg ultramicrocrystalline griseofulvin tablets (See Table 1).

Table 1: Mean (\pm SD) of the Pharmacokinetic Parameters for Griseofulvin administered in applesauce as a Single Dose of Griseofulvin Ultramicrosize 250-mg Tablets Uncrushed and Crushed to fasted Healthy Volunteers (N=24)

	250 mg Ultramicrocrystalline	250 mg Ultramicrocrystalline Griseofulvin Tablets
--	---------------------------------	---

	Griseofulvin Tablets Unaltered	Griseofulvin Tablets Physically Altered (Crushed and in Applesauce)
C _{max} (ng/mL)	600.61 (±167.6)	672.61 (±146.2)
T _{max} (hr)	4.04 (±2.2)	3.08 (±1.02)
AUC (ng·hr/mL)	8618.89 (±1907.2)	9023.71 (±1911.5)

INDICATIONS

Griseofulvin Ultramicrosize is indicated for the treatment of the following ringworm infections; tinea corporis (ringworm of the body), tinea pedis (athlete's foot), tinea cruris (ringworm of the groin and thigh), tinea barbae (barber's itch), tinea capitis (ringworm of the scalp), and tinea unguium (onychomycosis, ringworm of the nails), when caused by one or more of the following genera of fungi: *Trichophyton rubrum*, *Trichophyton tonsurans*, *Trichophyton mentagrophytes*, *Trichophyton interdigitalis*, *Trichophyton verrucosum*, *Trichophyton megnini*, *Trichophyton gallinae*, *Trichophyton crateriform*, *Trichophyton sulphureum*, *Trichophyton schoenleini*, *Microsporum audouini*, *Microsporum canis*, *Microsporum gypseum* and *Epidermophyton floccosum*. NOTE: Prior to therapy, the type of fungi responsible for the infection should be identified. The use of the drug is not justified in minor or trivial infections which will respond to topical agents alone. Griseofulvin is *not* effective in the following: bacterial infections, candidiasis (moniliasis), histoplasmosis, actinomycosis, sporotrichosis, chromoblastomycosis, coccidioidomycosis, North American blastomycosis, cryptococcosis (torulosis), tinea versicolor and nocardiosis.

CONTRAINDICATIONS

Two cases of conjoined twins have been reported since 1977 in patients taking griseofulvin during the first trimester of pregnancy. Griseofulvin should not be prescribed to pregnant patients. If the patient becomes pregnant while taking this drug, the patient should be apprised of the potential hazard to the fetus.

This drug is contraindicated in patients with porphyria or hepatocellular failure and in individuals with a history of hypersensitivity to griseofulvin.

WARNINGS

Prophylactic Usage

Safety and efficacy of griseofulvin for prophylaxis of fungal infections have not been established.

Serious Skin Reactions

Severe skin reactions (e.g. Stevens-Johnson syndrome, toxic epidermal necrolysis) and erythema multiforme have been reported with griseofulvin use. These reactions may be serious and may result in hospitalization or death. If severe skin reactions occur, griseofulvin should be discontinued (see ADVERSE REACTIONS section).

Hepatoxicity

Elevations in AST, ALT, bilirubin, and jaundice have been reported with griseofulvin use. These reactions may be serious and may result in hospitalization or death. Patients should be monitored for hepatic adverse events and discontinuation of griseofulvin considered if warranted (see ADVERSE REACTIONS section).

Animal Toxicology

Chronic feeding of griseofulvin, at levels ranging from 0.5%-2.5% of the diet resulted in the development of liver tumors in several strains of mice, particularly in males. Smaller particle sizes result in an enhanced effect. Lower oral dosage levels have not been tested. Subcutaneous administration of relatively small doses of griseofulvin once a week during the first three weeks of life has also been reported to induce hepatomata in mice. Thyroid tumors, mostly adenomas but some carcinomas, have been reported in male rats receiving griseofulvin at levels of 2.0%, 1.0% and 0.2% of the diet, and in female rats receiving the two higher dose levels. Although studies in other animal species have not yielded evidence of tumorigenicity, these studies were not of adequate design to form a basis for conclusion in this regard. In subacute toxicity studies, orally administered griseofulvin produced hepatocellular necrosis in mice, but this has not been seen in other species. Disturbances in porphyrin metabolism have been reported in griseofulvin-treated laboratory animals. Griseofulvin has been reported to have a colchicine-like effect on mitosis and cocarcinogenicity with methylcholanthrene in cutaneous tumor induction in laboratory animals.

Usage in Pregnancy

see CONTRAINDICATIONS section.

Animal Reproduction Studies

It has been reported in the literature that griseofulvin was found to be embryotoxic and teratogenic on oral administration to pregnant rats. Pups with abnormalities have been reported in the litters of a few bitches treated with griseofulvin. Suppression of spermatogenesis has been reported to occur in rats, but investigation in man failed to confirm this.

PRECAUTIONS

Patients on prolonged therapy with any potent medication should be under close observation. Periodic monitoring of organ system function, including renal, hepatic and hematopoietic, should be done. Since griseofulvin is derived from species of *Penicillium*, the possibility of cross-sensitivity with penicillin exists; however, known penicillin-sensitive patients have been treated without difficulty. Since a photosensitivity reaction is occasionally associated with griseofulvin therapy, patients should be warned to avoid exposure to intense natural or artificial sunlight. Lupus erythematosus or lupus-like syndromes have been reported in patients receiving griseofulvin. Griseofulvin decreases the activity of warfarin-type anticoagulants so that patients receiving these drugs concomitantly may require dosage adjustment of the anticoagulant during and after griseofulvin therapy. Barbiturates usually depress griseofulvin activity and concomitant administration may require a dosage adjustment of the antifungal agent. There have been reports in the literature of possible interactions between griseofulvin and oral contraceptives. The effect of alcohol may be potentiated by griseofulvin, producing such effects as tachycardia and flush.

ADVERSE REACTIONS

There have been post-marketing reports of severe skin and hepatic adverse events associated with griseofulvin use (see WARNINGS section).

When adverse reactions occur, they are most commonly of the hypersensitivity type such as skin rashes, urticaria, erythema multiforme-like drug reactions, and rarely, angioneurotic edema, and may necessitate withdrawal of therapy and appropriate countermeasures. Paresthesia of the hands and feet have been reported after extended therapy. Other side effects reported occasionally are oral thrush, nausea, vomiting, epigastric distress, diarrhea, headache, fatigue, dizziness, insomnia, mental confusion, and impairment of performance of routine activities. Proteinuria and leukopenia have been reported rarely. Administration of the drug should be discontinued if granulocytopenia occurs. When rare, serious reactions occur with griseofulvin, they are usually associated with high dosages, long periods of therapy, or both.

DOSAGE AND ADMINISTRATION

Accurate diagnosis of infecting organism is essential. Identification should be made either by direct microscopic examination of a mounting of infected tissue in a solution of potassium hydroxide or by culture on an appropriate medium. Medication must be continued until the infecting organism is completely eradicated as indicated by appropriate clinical or laboratory examination. Representative treatment periods are tinea capitis, 4 to 6 weeks; tinea corporis, 2 to 4 weeks; tinea pedis, 4 to 8 weeks; tinea unguium—depending on rate of growth—fingernails, at least 4 months; toenails, at least 6 months.

General measures in regard to hygiene should be observed to control sources of infection or reinfection. Concomitant use of appropriate topical agents is usually required, particularly in treatment of tinea pedis. In some forms of athlete's foot, yeasts and bacteria may be involved as well as fungi. Griseofulvin will not eradicate the bacterial or monilial infection.

Griseofulvin Ultramicrosize tablets may be swallowed whole or crushed and sprinkled onto 1 tablespoonful of applesauce and swallowed immediately without chewing.

Adults

Daily administration of 375 mg (as a single dose or in divided doses) will give a satisfactory response in most patients with tinea corporis, tinea cruris, and tinea capitis. For those fungal infections more difficult to eradicate, such as tinea pedis and tinea unguium, a divided dose of 750 mg is recommended.

Pediatric Use

Approximately 3.3 mg per pound of body weight per day of ultramicrosize griseofulvin is an effective dose for most pediatric patients. On this basis, the following dosage schedule is suggested: Children weighing 35-60 pounds - 125 mg to 187.5 mg daily. Pediatric patients weighing over 60 pounds - 187.5 mg to 375 mg daily. Children and infants 2 years of age and younger - dosage has not been established.

Clinical experience with griseofulvin in children with tinea capitis indicates that a single daily dose is effective. Clinical relapse will occur if the medication is not continued until the infecting organism is eradicated.

HOW SUPPLIED

Griseofulvin Ultramicrosize Tablets, 125 mg, white, scored, elliptical-shaped, embossed "Gris-PEG" on one side and "125" on the other. Griseofulvin Ultramicrosize Tablets, 250 mg, white, scored, capsule-shaped, embossed "Gris-PEG" on one side and "250" on the other. The 125 mg strength is available in bottles of 100 (NDC 68682-519-01). The 250 mg strength is available in bottles of 100 (NDC 68682-520-01). Both strengths are film-coated.

Rx ONLY

STORAGE

Store Griseofulvin Ultramicrosize tablets at controlled room temperature 15° - 30°C (59° - 86°F) in tight, light-resistant containers.

Distributed by:

Oceanside Pharmaceuticals, a division of Valeant Pharmaceuticals North America LLC
Bridgewater, NJ 08807

2007075

Rev. 11/12

PRINCIPAL DISPLAY PANEL - 250 mg Bottle Label

NDC 68682-520-01

Griseofulvin Ultramicrosize

Tablets USP

250 mg

Rx ONLY

100 Tablets

OCEANSIDE PHARMACEUTICALS

NON-VARNISH

NDC 68682-520-01

**Griseofulvin
Ultramicrosize**

Tablets USP

250 mg

Rx
ONLY

100 Tablets

OCEANSIDE[®]
PHARMACEUTICALS

Store at controlled room temperature at 15°-30°C (59°-86°F) in tight, light resistant containers.

Dosage and Administration:
Read package insert for prescribing information.

Distributed by:
Oceanside Pharmaceuticals,
a division of Valeant
Pharmaceuticals North America LLC,
Bridgewater, NJ 08807

Rev. 11/12 53445

LOT EXP.

3 68682 52001 9

PRINCIPAL DISPLAY PANEL - 125 mg Bottle Label

NDC 68682-519-01

Griseofulvin Ultramicrosize

Tablets USP

125 mg

Rx ONLY

100 Tablets

OCEANSIDE PHARMACEUTICALS

NDC 68682-519-01

NON-VARNISH

Griseofulvin Ultramicrosize

Tablets USP

125 mg

R
ONLY
100 Tablets



Store at controlled room temperature at 15°-30°C (59°-86°F) in tight, light resistant containers.

Dosage and Administration:
Read package insert for prescribing information.

Distributed by:
Oceanside Pharmaceuticals,
a division of Valeant Pharmaceuticals
North America LLC,
Bridgewater, NJ 08807

Rev. 11/12 53444

LOT EXP.



3 68682 51901 3

GRISEOFULVIN

griseofulvin tablet, film coated

Product Information

Product Type	HUMAN PRESCRIPTION DRUG LABEL	Item Code (Source)	NDC:68682-520
Route of Administration	ORAL	DEA Schedule	

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Griseofulvin (Griseofulvin)	Griseofulvin	250 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE	
MAGNESIUM STEARATE	
METHYLCELLULOSE (15 CPS)	
METHYLPARABEN	
POLYETHYLENE GLYCOL 400	
POLYETHYLENE GLYCOL 8000	
POVIDONE	
SODIUM LAURYL SULFATE	
TITANIUM DIOXIDE	

Product Characteristics

Color	WHITE	Score	2 pieces
-------	-------	-------	----------

Shape	OVAL (capsule-shaped)	Size	16 mm
Flavor		Imprint Code	GRIS;PEG;250
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68682-520-01	100 in 1 BOTTLE, PLASTIC		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA AUTHORIZED GENERIC	NDA050475	12/03/2012	

GRISEOFULVIN			
griseofulvin tablet, film coated			
Product Information			
Product Type	HUMAN PRESCRIPTION DRUG LABEL	Item Code (Source)	NDC:68682-519
Route of Administration	ORAL	DEA Schedule	
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
Griseofulvin (Griseofulvin)	Griseofulvin	125 mg	
Inactive Ingredients			
Ingredient Name	Strength		
SILICON DIOXIDE			
LACTOSE			
MAGNESIUM STEARATE			
METHYLCELLULOSE (15 CPS)			
METHYL PARABEN			
POLYETHYLENE GLYCOL 400			
POLYETHYLENE GLYCOL 8000			
POVIDONE			
TITANIUM DIOXIDE			
Product Characteristics			
Color	WHITE	Score	2 pieces
Shape	OVAL (elliptical-shaped)	Size	19 mm
Flavor		Imprint Code	GRIS;PEG;125
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68682-519-01	100 in 1 BOTTLE, PLASTIC		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA AUTHORIZED GENERIC	NDA050475	12/03/2012	

Labeler - Oceanside Pharmaceuticals (832011691)

Establishment

Name	Address	ID/FEI	Business Operations
Novartis Consumer Health		129836151	MANUFACTURE(68682-520, 68682-519)

Revised: 12/2012

Oceanside Pharmaceuticals