

WATER PIK ULTRA CONTROL FLUORIDE MELON - sodium fluoride aerosol, foam
WATER PIK, INC.

Indications: For the topical application of fluoride to aid in the protection against dental caries.

Contraindications: This product should not be used with individuals with known sensitivities to fluoride.

Directions for Use:

1. Remove cap from can.
2. The first time you dispense from a new can, gently lift upward on the nozzle to break the protective shipping tab (Fig. 1).
3. Shake can vigorously for at least 10 seconds before each use.
4. Hold can completely upside down to dispense (Fig. 2).
5. Point can towards applicator tray and slowly press the nozzle to fill tray. Use one press per arch, as foam will expand slightly to fill the tray.
6. For best results, dry teeth prior to application.
7. Insert the filled tray and have the patient bite down lightly, with a chewing motion to ensure interproximal coverage.
8. Leave tray(s) in contact with teeth for 1 to 4 minutes.
9. Use a saliva ejector during treatment to minimize ingestion of product.
10. Remove tray(s) and have patient expectorate.
11. Instruct patient not to eat, drink, or rinse for at least 30 minutes after application.

Contains:

1.23% Fluoride ion from sodium fluoride in a proprietary acidulated phosphate flavored foam base. Does not contain chlorofluorocarbon propellant.

Sweetened with Xylitol (sucralose). Does not contain Aspartame or Saccharin. Gluten free.

Precautions:

Use only as directed.

For dental use only. U.S. Federal Law prohibits dispensing without prescription.

Not for unsupervised home use. Safety and efficacy in patients under the age of 3 has not been established. Repeated use of APF formulations in patients with porcelain or composite restorations should be avoided due to the possibility of etching the restoration surface. Warning:

Do not swallow.

Keep out of reach of children.

Avoid spraying toward open flame.

Contents under pressure.

Do not puncture or incinerate.

Indications: For the topical application of fluoride to aid in the protection against dental caries.

Contraindications: This product should not be used with individuals with known sensitivities to fluoride.

Directions for Use:

1. Remove cap from can.
2. The first time you dispense from a new can, gently lift upward on the nozzle to break the protective shipping tab (Fig. 1).
3. Shake can vigorously for at least 10 seconds before each use.
4. Hold can completely upside down to dispense (Fig. 2).
5. Point can towards applicator tray and slowly press the nozzle to fill tray. Use one press per arch, as foam will expand slightly to fill the tray.
6. For best results, dry teeth prior to application.
7. Insert the filled tray and have the patient bite down lightly, with a chewing motion to ensure interproximal coverage.
8. Leave tray(s) in contact with teeth for 1 to 4 minutes.
9. Use a saliva ejector during treatment to minimize ingestion of product.
10. Remove tray(s) and have patient expectorate.
11. Instruct patient not to eat, drink, or rinse for at least 30 minutes after application.



Fig. 1



Fig. 2

Contains:

1.23% Fluoride ion from sodium fluoride in a proprietary acidulated phosphate flavored foam base. Does not contain chlorofluorocarbon propellant.

Sweetened with Xylitol (sucralose). Does not contain Aspartame or Saccharin. Gluten free.

Precautions:

Use only as directed.

For dental use only. U.S. Federal Law prohibits dispensing without prescription.

Not for unsupervised home use. Safety and efficacy in patients under the age of 3 has not been established. Repeated use of APF formulations in patients with porcelain or composite restorations should be avoided due to the possibility of etching the restoration surface.

Warning:

Do not swallow.
Keep out of reach of children.
Avoid spraying toward open flame.
Contents under pressure.
Do not puncture or incinerate.

waterpik Ultra Control

Fluoride Foam

60 Second APF* Topical Fluoride

- **Precise foaming action**
- **Sweetened with Xylitol**
- **Gluten free**

*Acidulated Phosphate Fluoride
1.23% Fluoride Ion (pH 3.5)



Waterpik® (stylized) is a trademark of Water Pik, Inc. registered in Canada, EU, Japan, Mexico, Switzerland, and the United States.

©2011 Water Pik, Inc.

Waterpik Item Number
20014142

FN 20014518-FAA
373482



Melon

NET WT. 4.4 oz (125g)

NET WT. 4.4 oz (125g)

waterpik
Ultra Control Fluoride
Foam
60 Second APF* Topical Fluoride



Melon

NET WT. 4.4 oz (125g)

waterpik
Ultra Control

Fluoride Foam
60 Second APF* Topical Fluoride

- **Precise foaming action**
- **Sweetened with Xylitol**
- **Gluten free**

*Acidulated Phosphate Fluoride
1.23% Fluoride Ion (pH 3.5)

For professional use only.
Store at room temperature.
Shake well before each use.



Storage: Between 55-80°F.

Keep from freezing.

To obtain a MSDS sheet, call
Water Pik, Inc. customer service
at 1-800-525-2020.

Made in the U.S.A.

Distributed by:

Water Pik, Inc.
1730 E. Prospect Road
Fort Collins, CO 80553-0001
1-800-525-2020
www.waterpik.com



Melon

NET WT. 4.4 oz (125g)

- Precise foaming action
- Sweetened with Xylitol
- Gluten free

*Acidulated Phosphate Fluoride
1.23% Fluoride Ion (pH 3.5)

For professional use only.

Store at room temperature
between 55°- 80° F.

Shake well before each use.

To obtain a MSDS sheet, please
call Water Pik, Inc. customer
service at 1-800-525-2020.

Made in the U.S.A.

Distributed by:
Water Pik, Inc.
1730 E. Prospect Road
Fort Collins, CO 80563-0001
1-800-525-2020

www.waterpik.com

Waterpik® (stylized) is a
trademark of Water Pik, Inc.,
registered Canada, EU, Japan,
Mexico, Switzerland, and the
United States.

©2011 Water Pik, Inc.
Water Pik Item Number:
20014142
FN20014517-FAA
373483



60 Second APF® Topical Fluoride

waterpik® Ultra Control Fluoride Foam



Melon

NET WT. 4.4 oz. (125g)

Indications: For the topical application of fluoride to aid in the
protection against dental caries.

Contraindications: This product should not be used with
individuals with known sensitivities to fluoride.

Directions for Use:

1. Remove cap from can.
2. The first time you dispense from a new can, gently lift upward
on the nozzle to break the protective shipping tab (Fig. 1).
3. Shake can vigorously for at least 10 seconds before each use.
4. Hold can completely upside down to dispense (Fig. 2).
5. Point can towards applicator tray and slowly press the nozzle
to fill tray. Use one press per arch, as foam will expand
slightly to fill the tray.
6. For best results, dry teeth prior to application.
7. Insert the filled tray and have the patient bite down lightly,
with a chewing motion to ensure interproximal coverage.
8. Leave tray(s) in contact with teeth for 1 to 4 minutes.
9. Use a saliva ejector during treatment to minimize ingestion
of product.
10. Remove tray(s) and have patient expectorate.
11. Instruct patient not to eat, drink, or rinse for at least 30
minutes after application.

Contains:
1.23% Fluoride ion from sodium fluoride in a proprietary
acidulated phosphate flavored foam base. Does not contain
chlorofluorocarbon propellant.
Sweetened with Xylitol (sucralose). Does not contain Aspartame
or Saccharin. Gluten free.

Precautions:

Use only as directed.
For dental use only. U.S. Federal Law prohibits dispensing
without prescription.

Not for unsupervised home use. Safety and efficacy in patients
under the age of 3 has not been established. Repeated use of
APF formulations in patients with porcelain or composite
restorations should be avoided due to the possibility of etching
the restorations surface.

Warning:

Do not swallow.
Keep out of reach of children.
Avoid spraying toward open flame.
Contents under pressure.
Do not puncture or incinerate.



Fig. 1



Fig. 2