# DRUG FACTS

PurposeActive ingredient(s)

#### Use(s)

- temporarily relieves the following symptoms due to hay fever or other upper respiratory allergies:
- sneezing
- runny nose
- itchy, watery eyes
- itching of the nose or throat

#### Warnings

#### Do not use

to make a child sleepy

# Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland

# Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

#### When using this product

- excitability may occur, especially in children
- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

## Stop use and ask a doctor if

## If pregnant or breast-feeding

#### Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## Directions adults and children 12 years and over 1 tablet every 4 to 6 hours, not to exceed 6 tablets in 24 hours ½ tablet (break tablet in half) every 4 to 6 hours, not to exceed 3 whole tablets in 24 hours children 6 to under 12 years of age children 4 to under 6 years of age do not use unless directed by a doctor children under 4 years of age do not use

# Other information

store between  $20^{\circ}$  to  $25^{\circ}C$  (68° to 77°F)

# Inactive ingredients

D&C yellow no. 10 aluminum lake, lactose monohydrate, magnesium stearate, pregelatinized starch