DRUG FACTS

Active ingredient(s) ***Purpose***

Use(s)
temporarily relieves coughs due to cold or bronchial irritation

Warnings
Do not use with any other product containing diphenhydramine, even one used on skin
Ask a doctor before use if you have
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough is accompanied by excessive phlegm (mucus)
- breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland

Do not use

Ask a doctor before use if you have

Ask a doctor or pharmacist before use if you are
if you are taking tranquilizers or sedatives

When using this product
- you may get drowsy
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if
if symptoms last for more than 1 week or recurred or accompanied by fever, rash, headache

If pregnant or breast-feeding
If pregnant or breast-feeding
- ask a health professional before use.

Keep out of reach of children.
- In case of accidental overdose, get medical help or contact a Poison Control Center right away.

Keep out of reach of children
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
Do not take more than six doses

Adults and children over 12 years	2 teaspoonfuls (TSP) every 4 hours
children under 12 years	DO NOT USE

Other information
- Store below 20°-25°C (68°-77°F)
- Protect from freezing.

Inactive ingredients
Alcohol 5%, ammonium chloride, citric acid, D&C red no. 33, FD&C red no. 40, menthol, methylparaben, propylene glycol, propylparaben, sodium citrate, strawberry flavor, sucrose, water