

## DRUG FACTS

**Active ingredient(s)**

**Purpose**

### Use(s)

temporarily:  
# reduces fever  
# relieves minor aches and pains due to: # the common cold # flu # headaches # sore throat # toothaches

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes: # more than 5 doses in 24 hours, which is the maximum daily amount # with other drugs containing acetaminophen

**Sore throat warning:** if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if your child has liver disease.**

**Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.**

**Ask a doctor before use if you have**

**Ask a doctor or pharmacist before use if you are**

### When using this product

- do not exceed recommended dosage. (see overdose warning)

### Stop use and ask a doctor if

- # new symptoms occur
- # redness or swelling is present
- # pain gets worse or lasts for more than 5 days
- # fever gets worse or lasts for more than 3 days. These could be signs of serious condition.

### If pregnant or breast-feeding

### Keep out of reach of children

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

### Directions

# this product does not contain directions or complete warnings for adult use # do not give more than directed (see overdose warning) # find right dose chart, if possible use weight to dose; otherwise, use age # use only enclosed dropper designed for use with this product, do not use any other dosing device # fill to dose level # dispense liquid slowly into child's mouth, toward inner cheek # may be given alone or mixed with formula, milk, juice etc. # if needed repeat dose every 4 hours # do not give more than 5 times in 24 hours # Replace dropper tightly to maintain child resistance

Weight (lb)	Age (yr)	Dose (mL)
Under 24	under 2 years	Ask a Doctor
24-35	2-3 years	1.6 mL (0.8mL +0.8mL)

For accurate dosing follow dosing instructions using the enclosed dropper. Fill dropper to 0.8 mL or prescribed level, and dispense with a single firm squeeze of the dropper bulb

### Other information

Store between 20°-25°C (68°-77°F)

### Other information

### Inactive ingredients

citric acid, FD&C yellow no. 6, cherry flavor, methylparaben, saccharin sodium, sodium benzoate, sodium citrate, propylene glycol and purified water.