DRUG FACTS	
Active ingredient(s)	Purpose
Use(s)Helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bother	ersome mucus and make coughs more productive.
Warnings	
Do not use	
Ask a doctor before use if you have a cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema a cough that occurs with too much phlegm (mucus)	
Ask a doctor or pharmacist before use if you are	
When using this product	
Stop use and ask a doctor ifStop use and ask doctor if cough persists for more than 1 week, tends to recur, or is accompanied by a fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.	
If pregnant or breast-feeding	
Keep out of reach of childrenKEEP OUT OF REACH OF CHILDREN. In case of overdose, get medical help or contact a Poison Control Center right away.	
Directions	
Adults and children 12 years of age and over:	1 tablet every 4 hours, not to exceed 6 tablets in 24 hours.

Children under 12 years of age: ·Take with a full glass of water

- Other information
 Tamper-evident by foil seal under cap. Do not use if foil seal is broken or missing
 Store between 20°-25°C (68°-77°F).
 Supplied in a tight, light-resistant container with a child-resistant cap.

- Bidex-400 Tablets are white to off white, biconex, capsule shaped tablets, debossed "649" bisect "649" on one side and plain on the other side.

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, hypromellose, maltodextrin, povidone, silicified microcrystalline cellulose, and stearic acid. May also contain sodium starch glycolate.

consult a doctor.