CIPROFLOXACIN - ciprofloxacin hydrochloride tablet Unit Dose Services

Ciprofloxacin Tablets 250 mg, 500 mg and 750 mg

WARNING:

Fluoroquinolones, including CIPROFLOXACIN TABLETS USP, 250 mg, 500 mg and 750 mg, are associated with an increased risk of tendinitis and tendon rupture in all ages. This risk is further increased in older patients usually over 60 years of age, in patients taking corticosteroid drugs, and in patients with kidney, heart or lung transplants (See).

WARNINGS

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Ciprofloxacin Tablets USP, 250 mg, 500 mg and 750 mg and other antibacterial drugs Ciprofloxacin Tablets USP, 250 mg, 500 mg and 750 mg should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria.

DESCRIPTION

Ciprofloxacin Hydrochloride Tablets USP, 250 mg, 500 mg and 750 mg are synthetic broad spectrum antimicrobial agents for oral administration. Ciprofloxacin hydrochloride, USP, a fluoroquinolone, is the monohydrochloride monohydrate salt of 1-cyclopropyl-6-fluoro-1, 4-dihydro-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid. It is a faintly yellowish to light yellow crystalline substance with a molecular weight of 385.8. Its empirical formula is C H FN O •HCl•H O and its chemical structure is as follows: Ciprofloxacin is 1-cyclopropyl-6-fluoro-1,4-dihydro-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid. Its empirical formula is C H FN O and its molecular weight is 331.4. It is a faintly yellowish to light yellow crystalline substance and its chemical structure is as follows: Ciprofloxacin Tablets USP are film-coated tablets and are available in 250 mg, 500 mg and 750 mg (ciprofloxacin equivalent) strengths. Ciprofloxacin Tablets are white to slightly yellowish. The inactive ingredients are pregelatinized starch, microcrystalline cellulose, colloidal silicon dioxide, crospovidone, magnesium stearate, hypromellose, titanium dioxide, polyethylene glycol and purified water.

1718332

CLINICAL PHARMACOLOGY

Absorption:

Ciprofloxacin given as an oral tablet is rapidly and well absorbed from the gastrointestinal tract after oral administration. The absolute bioavailability is approximately 70% with no substantial loss by first pass metabolism. Ciprofloxacin maximum serum concentrations and area under the curve are shown in the chart for the 250 mg to 1000 mg dose range. Maximum serum concentrations are attained 1 to 2 hours after oral dosing. Mean concentrations 12 hours after dosing with 250, 500, or 750 mg are 0.1, 0.2, and 0.4 µg/mL, respectively. The serum elimination half-life in subjects with normal renal function is approximately 4 hours. Serum concentrations increase proportionately with doses up to 1000 mg. A 500 mg oral dose given every 12 hours has been shown to produce an area under the serum concentration time curve (AUC) equivalent to that produced by an intravenous infusion of 400 mg ciprofloxacin given over 60 minutes every 12 hours. A 750 mg oral dose given every 12 hours has been shown to produce an AUC at steady-state equivalent to that produced by an intravenous infusion of 400 mg given over 60 minutes every 8 hours. A 750 mg oral dose results in a C similar to that observed with a 400 mg I.V. dose. A 250 mg oral dose given every 12 hours produces an AUC equivalent to that produced by an infusion of 200 mg ciprofloxacin given every 12 hours. AUC AUC 24h=AUC x 2 AUC 24h=AUC x 3

Dose (mg)	Maximum Serum Concentrations (μg/mL)					der Curv ug•hr/mL	
250 500 750 1000	1.2 5.4	2.4	4.3	4.8 30.8	11.6	20.2	

max

Steady-state Pharmacokinetic Parameters Following Multiple Oral and I.V. Doses					
Parameters AUC	500 mg q12h,	400 mg q12h,	750 mg q12h,	400 mg q8h,	
(μg•hr/mL) C	P.O. 13.7	I.V. 12.7	P.O. 31.6	I.V. 32.9	
(μg/mL)	2.97	4.56	3.59	4.07	
	a	a	b	С	
max					

^a0-12h ^b0-12h ^c0-8h

Distribution:

The binding of ciprofloxacin to serum proteins is 20 to 40% which is not likely to be high enough to cause significant protein binding interactions with other drugs. After oral administration, ciprofloxacin is widely distributed throughout the body. Tissue concentrations often exceed serum concentrations in both men and women, particularly in genital tissue including the prostate. Ciprofloxacin is present in active form in the saliva, nasal and bronchial secretions, mucosa of the sinuses, sputum, skin blister fluid, lymph, peritoneal fluid, bile, and prostatic secretions. Ciprofloxacin has also been detected in lung, skin, fat, muscle, cartilage, and bone. The drug diffuses into the cerebrospinal fluid (CSF); however, CSF concentrations are generally less than 10% of peak serum concentrations. Low levels of the drug have been detected in the aqueous and vitreous humors of the eye.

Metabolism:

Four metabolites have been identified in human urine which together account for approximately 15% of an oral dose. The metabolites have antimicrobial activity, but are less active than unchanged ciprofloxacin. Ciprofloxacin is an inhibitor of human cytochrome P450 1A2 (CYP1A2) mediated metabolism. Coadministration of ciprofloxacin with other drugs primarily metabolized by CYP1A2 results in increased plasma concentrations of these drugs and could lead to clinically significant adverse events of the coadministered drug (see).

CONTRAINDICATIONS; ; WARNINGSPRECAUTIONS: Drug Interactions

Excretion:

The serum elimination half-life in subjects with normal renal function is approximately 4 hours. Approximately 40 to 50% of an orally administered dose is excreted in the urine as unchanged drug. After a 250 mg oral dose, urine concentrations of ciprofloxacin usually exceed 200 $\mu g/mL$ during the first two hours and are approximately 30 $\mu g/mL$ at 8 to 12 hours after dosing. The urinary excretion of ciprofloxacin is virtually complete within 24 hours after dosing. The renal clearance of ciprofloxacin, which is approximately 300 mL/minute, exceeds the normal glomerular filtration rate of 120 mL/minute. Thus, active tubular secretion would seem to play a significant role in its elimination. Co-administration of probenecid with ciprofloxacin results in about a 50% reduction in the ciprofloxacin renal clearance and a 50% increase in its concentration in the systemic circulation. Although bile concentrations of ciprofloxacin are several fold higher than serum concentrations after oral dosing, only a small amount of the dose administered is recovered from the bile as unchanged drug. An additional 1 to 2% of the dose is recovered from the bile in the form of metabolites. Approximately 20 to 35% of an oral dose is recovered from the feces within 5 days after dosing. This may arise from either biliary clearance or transintestinal elimination.

Drug-drug Interactions:

When Ciprofloxacin Tablets USP, 250 mg, 500 mg and 750 mg is given concomitantly with food, there is a delay in the absorption of the drug, resulting in peak concentrations that occur closer to 2 hours after dosing rather than 1 hour. The overall absorption of Ciprofloxacin Tablets USP, 250 mg, 500 mg and 750 mg however, is not substantially affected. Concurrent administration of antacids containing magnesium hydroxide or aluminum hydroxide may reduce the bioavailability of ciprofloxacin by as much as 90%. (See .) The serum concentrations of ciprofloxacin and metronidazole were not altered when these two drugs were given concomitantly. Concomitant administration with tizanidine is contraindicated. (See) Concomitant administration of ciprofloxacin with theophylline decreases the

clearance of the ophylline resulting in elevated serum the ophylline levels and increased risk of a patient developing CNS or other adverse reactions. Ciprofloxacin also decreases caffeine clearance and inhibits the formation of paraxanthine after caffeine administration. (See .)

PRECAUTIONS

CONTRAINDICATIONS.

: WARNINGSPRECAUTIONS

Special Populations:

Pharmacokinetic studies of the oral (single dose) and intravenous (single and multiple dose) forms of ciprofloxacin indicate that plasma concentrations of ciprofloxacin are higher in elderly subjects (> 65 years) as compared to young adults. Although the C is increased 16- 40%, the increase in mean AUC is approximately 30%, and can be at least partially attributed to decreased renal clearance in the elderly. Elimination half-life is only slightly (~20%) prolonged in the elderly. These differences are not considered clinically significant. (See .) In patients with reduced renal function, the half-life of ciprofloxacin is slightly prolonged. Dosage adjustments may be required. (See .) In preliminary studies in patients with stable chronic liver cirrhosis, no significant changes in ciprofloxacin pharmacokinetics have been observed. The kinetics of ciprofloxacin in patients with acute hepatic insufficiency, however, have not been fully elucidated.

maxPRECAUTIONS: Geriatric Use

DOSAGE AND ADMINISTRATION

MICROBIOLOGY

Ciprofloxacin has activity against a wide range of gram-negative and gram-positive microorganisms. The bactericidal action of ciprofloxacin results from inhibition of the enzymes topoisomerase II (DNA gyrase) and topoisomerase IV, which are required for bacterial DNA replication, transcription, repair, and recombination. The mechanism of action of fluoroquinolones, including ciprofloxacin, is different from that of penicillins, cephalosporins, aminoglycosides, macrolides, and tetracyclines; therefore, microorganisms resistant to these classes of drugs may be susceptible to ciprofloxacin and other quinolones. There is no known cross-resistance between ciprofloxacin and other classes of antimicrobials. resistance to ciprofloxacin develops slowly by multiple step mutations. Ciprofloxacin is slightly less active when tested at acidic pH. The inoculum size has little effect when tested The minimal bactericidal concentration (MBC) generally does not exceed the minimal inhibitory concentration (MIC) by more than a factor of 2. Ciprofloxacin has been shown to be active against most strains of the following microorganisms, both and in clinical infections as described in the section of the package insert for Ciprofloxacin Hydrochloride Tablets USP, 250mg, 500mg and 750mg. *in vitroIn vitro*

in vitro.

in vitroINDICATIONS AND USAGE

Aerobic gram-positive microorganisms

(Many strains are only moderately susceptible.) (methicillin-susceptible strains only) (methicillin-susceptible strains only) (penicillin-susceptible strains only) *Enterococcus faecalis*

Staphylococcus aureus

Staphylococcus epidermidis Staphylococcus saprophyticus Streptococcus pneumoniae

Streptococcus pyogenes

Aerobic gram-negative microorganisms

Ciprofloxacin has been shown to be active against both and by use of serum levels as a surrogate marker (see and). The following data are available, . Ciprofloxacin exhibits minimum inhibitory concentrations (MICs) of 1 μ g/mL or less against most (\geq 90%) strains of the following microorganisms; however, the safety and effectiveness of ciprofloxacin in treating clinical infections due to these microorganisms have not been established in adequate and well-controlled clinical trials. Campylobacter jejuni Proteus mirabilis Citrobacter diversus Proteus

vulgaris Citrobacter freundii Providencia rettgeri Enterobacter

cloacae Providencia stuartii Escherichia coli Pseudomonas aeruginosa Haemophilus influenzae Salmonella typhi Haemophilus parainfluenzae

Serratia marcescens

Klebsiella pneumoniae Shigella boydii Moraxella catarrhalis Shigella dysenteriae Morganella morganii Shigella flexneri Neisseria gonorrhoeae Shigella sonnei

Bacillus anthracisin vitro INDICATIONS AND USAGEINHALATIONAL ANTHRAX – ADDITIONAL INFORMATION

in vitrobut their clinical significance is unknown

in vitro

Aerobic gram-positive microorganisms (penicillin-resistant strains only) *Staphylococcus haemolyticus Staphylococcus hominis Streptococcus pneumoniae*

Aerobic gram-negative microorganisms

Most strains of and some strains of are resistant to ciprofloxacin as are most anaerobic bacteria, including and .

Acinetobacter Iwoffii Pasteurella multocida Aeromonas hydrophila Salmonella enteritidis Edwardsiella tarda Vibrio cholerae Enterobacter aerogenes Vibrio

parahaemolyticus Klebsiella oxytoca Vibrio vulnificus Legionella pneumophila Yersinia enterocolitica

Burkholderia cepaciaStenotrophomonas maltophiliaBacteroides fragilisClostridium difficile

Susceptibility Tests

Dilution Techniques:

Quantitative methods are used to determine antimicrobial minimum inhibitory concentrations (MICs). These MICs provide estimates of the susceptibility of bacteria to antimicrobial compounds. The MICs should be determined using a standardized procedure. Standardized procedures are based on a dilution method (broth or agar) or equivalent with standardized inoculum concentrations and standardized concentrations of ciprofloxacin powder. The MIC values should be interpreted according to the following criteria: For testing aerobic microorganisms other than , , and : These interpretive standards are applicable only to broth microdilution susceptibility tests with streptococci using cation-adjusted Mueller-Hinton broth with 2-5% lysed horse blood. For testing and: This interpretive standard is applicable only to broth microdilution susceptibility tests with and using Test Medium. The current absence of data on resistant strains precludes defining any results other than "Susceptible". Strains yielding MIC results suggestive of a "nonsusceptible" category should be submitted to a reference laboratory for further testing. For testing: This interpretive standard is applicable only to agar dilution test with GC agar base and 1% defined growth supplement. A report of "Susceptible" indicates that the pathogen is likely to be inhibited if the antimicrobial compound in the blood reaches the concentrations usually achievable. A report of "Intermediate" indicates that the result should be considered equivocal, and, if the microorganism is not fully susceptible to alternative, clinically feasible drugs, the test should be repeated. This category implies possible clinical applicability in body sites where the drug is physiologically concentrated or in situations where high dosage of drug can be used. This category also provides a buffer zone, which prevents small uncontrolled technical factors from causing major discrepancies in interpretation. A report of "Resistant" indicates that the pathogen is not likely to be inhibited if the antimicrobial compound in the blood reaches the concentrations usually achievable; other therapy should be selected. Standardized susceptibility test procedures require the use of laboratory control microorganisms to control the technical aspects of the laboratory procedures. Standard ciprofloxacin powder should provide the following MIC values: This quality control range is applicable to only ATCC 49247 tested by a broth microdilution procedure using Test Medium (HTM). ATCC 33560 tested by broth microdilution procedure using cation adjusted Mueller Hinton broth with 2.5-5% lysed horse blood in a microaerophilic environment at 36-37°C for 48 hours and for 42°C at 24 hours, respectively. ATCC 49226 tested by agar dilution procedure using GC agar and 1% defined growth supplement in a 5% CO environment at 35-37°C for 20-24 hours.

Haemophilus influenzaeHaemophilus parainfluenzaeNeisseria gonorrhoeae^a

MIC (μg/mL)	Interpretation		
≤ 1	Susceptible (S)		
2	Intermediate (I)		
≥ 4	Resistant (R)		

a

 $Hae mophilus\ influenzae Hae mophilus\ para influenzae ^b$

MIC (μg/mL)	Interpretation		
≤ 1	Susceptible (S)		

Neisseria gonorrhoeae^c

MIC (μg/mL)	Interpretation
≤ 0.06	Susceptible (S)
0.12 - 0.5	Intermediate (I)
≥ 1	Resistant (R)

C

Organis m		MIC (μg/mL)
E. faecalis	ATCC 29212	0.25 – 2
E. coli	ATCC 25922	0.004 - 0.015
H. influenzae ^a	ATCC 49247	0.004 - 0.03
P. aeruginosa	ATCC 27853	0.25 - 1.0
S. aureus	ATCC 29213	0.12 - 0.5
C. jejuni ^b	ATCC 33560	0.06 - 0.25 and $0.03 - 0.12$
N. gonorrhoeae ^c	ATCC 49226	0.001-0.008

 $^{\mathrm{a}}H$. influenzaeHaemophilus $^{\mathrm{1}}$

^bC. jejuni²

^cN. gonorrhoeae₂³

Diffusion Techniques:

Quantitative methods that require measurement of zone diameters also provide reproducible estimates of the susceptibility of bacteria to antimicrobial compounds. One such standardized procedure requires the use of standardized inoculum concentrations. This procedure uses paper disks impregnated with 5μg ciprofloxacin to test the susceptibility of microorganisms to ciprofloxacin. Reports from the laboratory providing results of the standard single-disk susceptibility test with a 5-µg ciprofloxacin disk should be interpreted according to the following criteria: For testing, , methicillin-susceptible species, penicillin-susceptible,, and: These zone diameter standards are applicable only to tests performed for streptococci using Mueller-Hinton agar supplemented with 5% sheep blood incubated in 5% CO. For testing and: This zone diameter standard is applicable only to tests with and using Test Medium (HTM). The current absence of data on resistant strains precludes defining any results other than "Susceptible". Strains yielding zone diameter results suggestive of a "nonsusceptible" category should be submitted to a reference laboratory for further testing. For testing: This zone diameter standard is applicable only to disk diffusion tests with GC agar base and 1% defined growth supplement. Interpretation should be as stated above for results using dilution techniques. Interpretation involves correlation of the diameter obtained in the disk test with the MIC for ciprofloxacin. As with standardized dilution techniques, diffusion methods require the use of laboratory control microorganisms that are used to control the technical aspects of the laboratory procedures. For the diffusion technique, the 5-µg ciprofloxacin disk should provide the following zone diameters in these

laboratory test quality control strains: These quality control limits are applicable to only ATCC 49247 testing using Test Medium (HTM). These quality control limits are applicable only to tests conducted with ATCC 49226 performed by disk diffusion using GC agar base and 1% defined growth supplement.

 $Enterobacteria ceae Enterococcus\ faecalis Staphylococcus Streptococcus\ pneumoniae Streptococcus\ pyogenes Pseudomonas\ aeruginos a^a$

Zone Diameter (mm)	Interpretation		
≥ 21	Susceptible (S)		
16 – 20	Intermediate (I)		
≤ 15	Resistant (R)		

a₂

Haemophilus influenzaeHaemophilus parainfluenzae^b

Zone Diameter (mm)		Interpretation		
≥ 21		Susceptible (S)		

 $^{\mathrm{b}}$ Haemophilus influenzae Haemophilus parainfluenzae Haemophilus $^{\mathrm{3}}$

Neisseria gonorrhoeae^c

Zone Diameter (mm)	Interpretation		
≥41	Susceptible (S)		
28 - 40	Intermediate (I)		
≤ 27	Resistant (R)		

C

Organism		Zone Diameter (mm)
E. coli	ATCC 25922	30-40
H. influenzae ^a	ATCC 49247	34-42
N. gonorrhoeae ^b	ATCC 49226	48-58
P. aeruginosa	ATCC 27853	25-33
S. aureus	ATCC 25923	22-30

INDICATIONS AND USAGE

Ciprofloxacin Tablets USP, 250 mg, 500 mg and 750 mg is indicated for the treatment of infections caused by susceptible strains of the designated microorganisms in the conditions and patient populations listed below. Please see for specific recommendations.

DOSAGE AND ADMINISTRATION

Adult Patients:

caused by methicillin-susceptible or caused by or caused by or caused by or penicillin-susceptible Also, for the treatment of acute exacerbations of chronic bronchitis. NOTE: Although effective in clinical trials, ciprofloxacin is not a drug of first choice in the treatment of presumed or confirmed pneumonia secondary to caused by penicillin-susceptible or caused by methicillin-susceptible methicillin-susceptible or caused by or (used in combination with metronidazole) caused by or caused by (enterotoxigenic strains), or when antibacterial therapy is indicated. caused by NOTE: The efficacy of ciprofloxacin in the eradication of the chronic typhoid carrier state has not been demonstrated. due to **Urinary Tract Infections** Escherichia coli, Klebsiella pneumoniae, Enterobacter cloacae, Serratia marcescens, Proteus mirabilis, Providencia rettgeri, Morganella morganii, Citrobacter diversus, Citrobacter freundii, Pseudomonas aeruginosa, Staphylococcus epidermidis, Staphylococcus saprophyticus, Enterococcus faecalis. **Acute Uncomplicated Cystitis in females**

Escherichia coliStaphylococcus saprophyticus. Chronic Bacterial Prostatitis

Escherichia coliProteus mirabilis.Lower Respiratory Tract Infections

Escherichia coli, Klebsiella pneumoniae, Enterobacter cloacae, Proteus mirabilis, Pseudomonas aeruginosa, Haemophilus influenzae, Haemophilus parainfluenzae, Streptococcus pneumoniae. Moraxella catarrhalis

Streptococcus pneumoniae. Acute Sinusitis

Haemophilus influenzae, Streptococcus pneumoniae, Moraxella catarrhalis. Skin and Skin Structure Infections

Escherichia coli, Klebsiella pneumoniae, Enterobacter cloacae, Proteus mirabilis, Proteus vulgaris, Providencia stuartii, Morganella morganii, Citrobacter freundii, Pseudomonas aeruginosa, Staphylococcus aureus, Staphylococcus epidermidis, Streptococcus pyogenes. **Bone and Joint Infections**

Enterobacter cloacae, Serratia marcescens, Pseudomonas aeruginosa. **Complicated Intra-Abdominal Infections**

Escherichia coli, Pseudomonas aeruginosa, Proteus mirabilis, Klebsiella pneumoniae, Bacteroides fragilis. **Infectious Diarrhea**

Escherichia coliCampylobacter jejuni, Shigella boydii[†],Shigella dysenteriae, Shigella flexneriShigella sonnei[†]**Typhoid Fever (Enteric Fever)**

Salmonella typhi.

Uncomplicated cervical and urethral gonorrhea

Neisseria gonorrhoeae.

Pediatric patients (1 to 17 years of age):

due to NOTE: Although effective in clinical trials, ciprofloxacin is not a drug of first choice in the pediatric population due to an increased incidence of adverse events compared to controls, including events related to joints and/or surrounding tissues. (See and .) Ciprofloxacin, like other fluoroquinolones, is associated with arthropathy and histopathological changes in weight-bearing joints of juvenile animals. (See .)

Complicated Urinary Tract Infections and Pyelonephritis Escherichia coli.

, , WARNINGSPRECAUTIONS, Pediatric UseADVERSE REACTIONSCLINICAL STUDIES

ANIMAL PHARMACOLOGY

Adult and Pediatric Patients:

(post-exposure): To reduce the incidence or progression of disease following exposure to aerosolized Ciprofloxacin serum concentrations achieved in humans served as a surrogate endpoint reasonably likely to predict clinical benefit and provided the initial basis for approval of this indication. Supportive clinical information for ciprofloxacin for anthrax post-exposure prophylaxis was obtained during the anthrax bioterror attacks of October 2001. (See also,). Although treatment of infections due to this organism in this organ system demonstrated a clinically significant outcome, efficacy was studied in fewer than 10 patients. If anaerobic organisms are suspected of contributing to the infection, appropriate therapy should be administered. Appropriate culture and susceptibility tests should be performed before treatment in order to isolate and identify organisms causing infection and to determine their susceptibility to ciprofloxacin. Therapy with Ciprofloxacin Tablets USP, 250 mg, 500 mg and 750 mg may be initiated before results of these tests are known; once results become available appropriate therapy should be continued. As with other drugs, some strains of may develop resistance fairly rapidly during treatment with ciprofloxacin. Culture and susceptibility testing performed periodically during therapy will provide information not only on the therapeutic effect of the antimicrobial agent but also on the possible emergence of bacterial resistance. To reduce the development of drug-resistant bacteria and maintain the effectiveness of Ciprofloxacin Tablets USP, 250 mg, 500 mg and 750 mg and other antibacterial drugs, Ciprofloxacin Tablets USP, 250 mg, 500 mg and 750 mg should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

Inhalational anthraxBacillus anthracis.

⁵INHALATIONAL ANTHRAX – ADDITIONAL INFORMATION

†

Pseudomonas aeruginosa

CONTRAINDICATIONS

Ciprofloxacin is contraindicated in persons with a history of hypersensitivity to ciprofloxacin, any member of the quinolone class of antimicrobial agents, or any of the product components. Concomitant administration with tizanidine is contraindicated. (See)

PRECAUTIONS: Drug Interactions.

WARNINGS

Tendinopathy and Tendon Rupture:

Fluoroquinolones, including Ciprofloxacin Tablets USP, 250 mg, 500 mg and 750 mg, are associated with an increased risk of tendinitis and tendon rupture in all ages. This adverse reaction most frequently involves the Achilles tendon, and rupture of the Achilles tendon may require surgical repair. Tendinitis and tendon rupture in the rotator cuff (the shoulder), the hand, the biceps, the thumb, and other tendon sites have also been reported. The risk of developing fluoroquinolone-associated tendinitis and tendon rupture is further increased in older patients usually over 60 years of age, in patients taking corticosteroid drugs, and in patients with kidney, heart or lung transplants. Factors, in addition to age and corticosteroid use, that may independently increase the risk of tendon rupture include strenuous physical activity, renal failure, and previous tendon disorders such as rheumatoid arthritis. Tendinitis and tendon rupture have also occurred in patients taking fluoroquinolones who do not have the above risk factors. Tendon rupture can occur during or after completion of therapy; cases occurring up to several months after completion of therapy have been reported. Ciprofloxacin Tablets USP, 250 mg, 500 mg and 750 mg should be discontinued if the patient experiences pain, swelling, inflammation or rupture of a tendon. Patients should be advised to rest at the first sign of tendinitis or tendon rupture, and to contact their healthcare provider regarding changing to a non-quinolone antimicrobial drug.

Pregnant Women:

(See and subsections.)

THE SAFETY AND EFFECTIVENESS OF CIPROFLOXACIN IN PREGNANT AND LACTATING WOMEN HAVE NOT BEEN ESTABLISHED., PRECAUTIONS: PregnancyNursing Mothers

Pediatrics:

Ciprofloxacin should be used in pediatric patients (less than 18 years of age) only for infections listed in the section. An increased incidence of adverse events compared to controls, including events related to joints and/or surrounding tissues, has been observed. (See .) In pre-clinical studies, oral administration of ciprofloxacin caused lameness in immature dogs. Histopathological examination of the weight-bearing joints of these dogs revealed permanent lesions of the cartilage. Related quinolone-class drugs also produce erosions of cartilage of weight-bearing joints and other signs of arthropathy in immature animals of various species. (See .)

INDICATIONS AND USAGEADVERSE REACTIONS

ANIMAL PHARMACOLOGY

Cytochrome P450 (CYP450):

Ciprofloxacin is an inhibitor of the hepatic CYP1A2 enzyme pathway. Coadministration of ciprofloxacin and other drugs primarily metabolized by CYP1A2 (e.g. theophylline, methylxanthines, tizanidine) results in increased plasma concentrations of the coadministered drug and could lead to clinically significant pharmacodynamic side effects of the coadministered drug.

Central Nervous System Disorders:

Convulsions, increased intracranial pressure, and toxic psychosis have been reported in patients receiving quinolones, including ciprofloxacin. Ciprofloxacin may also cause central nervous system (CNS) events including: dizziness, confusion, tremors, hallucinations, depression, and, rarely, suicidal thoughts or acts. These reactions may occur following the first dose. If these reactions occur in patients receiving ciprofloxacin, the drug should be discontinued and appropriate measures instituted. As with all quinolones, ciprofloxacin should be used with caution in patients with known or suspected CNS disorders that may predispose to seizures or lower the seizure threshold (e.g. severe cerebral arteriosclerosis, epilepsy), or in the presence of other risk factors that may predispose to seizures or lower the seizure threshold (e.g. certain drug therapy, renal dysfunction). (See and .)

, , PRECAUTIONS: GeneralInformation for Patients Drug Interactions ADVERSE REACTIONS

Theophylline:

These reactions have included cardiac arrest, seizure, status epilepticus, and respiratory failure. Although similar serious adverse effects have been reported in patients receiving theophylline alone, the possibility that these reactions may be potentiated by ciprofloxacin cannot be eliminated. If concomitant use cannot be avoided, serum levels of theophylline should be monitored and dosage adjustments made as appropriate.

SERIOUS AND FATAL REACTIONS HAVE BEEN REPORTED IN PATIENTS RECEIVING CONCURRENT ADMINISTRATION OF CIPROFLOXACIN AND THEOPHYLLINE.

Hypersensitivity Reactions:

Serious and occasionally fatal hypersensitivity (anaphylactic) reactions, some following the first dose, have been reported in patients receiving quinolone therapy. Some reactions were accompanied by cardiovascular collapse, loss of consciousness, tingling, pharyngeal or facial edema, dyspnea, urticaria, and itching. Only a few patients had a history of hypersensitivity reactions. Serious anaphylactic reactions require immediate emergency treatment with epinephrine. Oxygen, intravenous steroids, and airway management, including intubation, should be administered as indicated. Other serious and sometimes fatal events, some due to hypersensitivity, and some due to uncertain etiology, have been reported rarely in patients receiving therapy with quinolones, including ciprofloxacin. These events may be severe and generally occur following the administration of multiple doses. Clinical manifestations may include one or more of the following:

- fever, rash, or severe dermatologic reactions (e.g., toxic epidermal necrolysis, Stevens-Johnson syndrome);
- vasculitis; arthralgia; myalgia; serum sickness;
- allergic pneumonitis;
- interstitial nephritis; acute renal insufficiency or failure;
- hepatitis; jaundice; acute hepatic necrosis or failure;
- anemia, including hemolytic and aplastic; thrombocytopenia, including thrombotic thrombocytopenic purpura; leukopenia; agranulocytosis; pancytopenia; and/or other hematologic abnormalities.

The drug should be discontinued immediately at the first appearance of a skin rash, jaundice, or any other sign of hypersensitivity and supportive measures instituted (See and). **PRECAUTIONS: Information for Patients ADVERSE REACTIONS**

Pseudomembranous Colitis:

associated diarrhea (CDAD) has been reported with use of nearly all antibacterial agents, including Ciprofloxacin Tablets USP, 250 mg, 500 mg and 750 mg, and may range in severity from mild diarrhea to fatal colitis. Treatment with antibacterial agents alters the normal flora of the colon leading to overgrowth of . *Clostridium difficileC. difficile*

produces toxins A and B which contribute to the development of CDAD. Hypertoxin producing strains of cause increased morbidity and mortality, as these infections can be refractory to antimicrobial therapy and may require colectomy. CDAD must be considered in all patients who present with diarrhea following antibiotic use. Careful medical history is necessary since CDAD has been reported to occur over two months after the administration of antibacterial agents. *C. difficileC. difficile*

If CDAD is suspected or confirmed, ongoing antibiotic use not directed against may need to be discontinued. Appropriate fluid and electrolyte management, protein supplementation, antibiotic

treatment of , and surgical evaluation should be instituted as clinically indicated. C. difficileC. difficile

Peripheral neuropathy:

Rare cases of sensory or sensorimotor axonal polyneuropathy affecting small and/or large axons resulting in paresthesias, hypoesthesias, dysesthesias and weakness have been reported in patients receiving quinolones, including ciprofloxacin. Ciprofloxacin should be discontinued if the patient experiences symptoms of neuropathy including pain, burning, tingling, numbness, and/or weakness, or is found to have deficits in light touch, pain, temperature, position sense, vibratory sensation, and/or motor strength in order to prevent the development of an irreversible condition.

Syphilis:

Ciprofloxacin has not been shown to be effective in the treatment of syphilis. Antimicrobial agents used in high dose for short periods of time to treat gonorrhea may mask or delay the symptoms of incubating syphilis. All patients with gonorrhea should have a serologic test for syphilis at the time of diagnosis. Patients treated with ciprofloxacin should have a follow-up serologic test for syphilis after three months.

PRECAUTIONS

General:

Crystals of ciprofloxacin have been observed rarely in the urine of human subjects but more frequently in the urine of laboratory animals, which is usually alkaline. (See .) Crystalluria related to ciprofloxacin has been reported only rarely in humans because human urine is usually acidic. Alkalinity of the urine should be avoided in patients receiving ciprofloxacin. Patients should be well hydrated to prevent the formation of highly concentrated urine. Quinolones, including ciprofloxacin, may also cause central nervous system (CNS) events, including: nervousness, agitation, insomnia, anxiety, nightmares or paranoia. (See and .) Alteration of the dosage regimen is necessary for patients with impairment of renal function. (See .) Moderate to severe photosensitivity/phototoxicity reactions, the latter of which may manifest as exaggerated sunburn reactions (e.g., burning, erythema, exudation, vesicles, blistering, edema) involving areas exposed to light (typically the face, "V" area of the neck, extensor surfaces of the forearms, dorsa of the hands), can be associated with the use of quinolones after sun or UV light exposure. Therefore, excessive exposure to these sources of light should be avoided. Drug therapy should be discontinued if phototoxicity occurs (See). As with any potent drug, periodic assessment of organ system functions, including renal, hepatic, and hematopoietic function, is advisable during prolonged therapy. Prescribing Ciprofloxacin Tablets USP, 250 mg, 500 mg and 750 mg in the absence of a proven or strongly suspected bacterial infection or a prophylactic indication is unlikely to provide benefit to the patient and increases the risk of the development of drug-resistant bacteria.

ANIMAL PHARMACOLOGYCentral Nervous System:

, , WARNINGSInformation for Patients Drug Interactions Renal Impairment:

DOSAGE AND ADMINISTRATIONPhotos ensitivity/Phototoxicity:

ADVERSE REACTIONS/Post-Marketing Adverse Events

Information for Patients:

Patients should be advised:

- to contact their healthcare provider if they experience pain, swelling, or inflammation of a tendon, or weakness or inability to use one of their joints; rest and refrain from exercise; and discontinue Ciprofloxacin Tablets USP, 250 mg, 500 mg and 750 mg treatment. The risk of severe tendon disorder with fluoroquinolones is higher in older patients usually over 60 years of age, in patients taking corticosteroid drugs, and in patients with kidney, heart or lung transplants.
- that antibacterial drugs including Ciprofloxacin Tablets USP, 250 mg, 500 mg and 750 mg should only be used to treat bacterial infections. They do not treat viral infections (e.g., the common cold). When Ciprofloxacin Tablets USP, 250 mg, 500 mg and 750 mg is prescribed to treat a bacterial infection, patients should be told that although it is common to feel better early in the course of therapy, the medication should be taken exactly as directed. Skipping doses or not completing the full course of therapy may (1) decrease the effectiveness of the immediate treatment and (2) increase the likelihood that bacteria will develop resistance and will not be treatable by Ciprofloxacin Tablets USP, 250 mg, 500 mg and 750 mg or other antibacterial drugs in the future.
- that ciprofloxacin may be taken with or without meals and to drink fluids liberally. As with other quinolones, concurrent administration of ciprofloxacin with magnesium/aluminum antacids, or sucralfate, Videx® (didanosine) chewable/buffered tablets or pediatric powder, other highly buffered drugs, or with other products containing calcium, iron or zinc should be avoided. Ciprofloxacin may be taken two hours before or six hours after taking these products. Ciprofloxacin should not be taken with dairy products (like milk or yogurt) or calcium-fortified juices alone since absorption of ciprofloxacin may be significantly reduced; however, ciprofloxacin may be taken with a meal that contains these products.
- that ciprofloxacin may be associated with hypersensitivity reactions, even following a single dose, and to discontinue the drug at the first sign of a skin rash or other allergic reaction.
- that photosensitivity/phototoxicity has been reported in patients receiving quinolones. Patients should minimize or avoid exposure to natural or artificial sunlight (tanning beds or UVA/B treatment) while taking quinolones. If patients need to be outdoors while using quinolones, they should wear loose-fitting clothes that protect skin from sun exposure and discuss other sun protection measures with their physician. If a sunburn-like reaction or skin eruption occurs, patients should contact their physician.
- that peripheral neuropathies have been associated with ciprofloxacin use. If symptoms of peripheral neuropathy including pain, burning, tingling, numbness and/or weakness develop, they should discontinue treatment and contact their physicians.
- that ciprofloxacin may cause dizziness and lightheadedness; therefore, patients should know how they react to this drug before they operate an automobile or machinery or engage in activities requiring mental alertness or coordination.
- that ciprofloxacin increases the effects of tizanidine (Zanaflex®). Patients should not use ciprofloxacin if they are already taking tizanidine.
- that ciprofloxacin may increase the effects of the ophylline and caffeine. There is a possibility of caffeine accumulation when products containing caffeine are consumed while taking quinolones.
- that convulsions have been reported in patients receiving quinolones, including ciprofloxacin, and to notify their physician before taking this drug if there is a history of this condition.
- that ciprofloxacin has been associated with an increased rate of adverse events involving joints and surrounding tissue structures (like tendons) in pediatric patients (less than 18 years of age). Parents should inform their child's physician if the child has a history of joint-related problems before taking this drug. Parents of pediatric patients should also notify their child's physician of any joint-related problems that occur during or following ciprofloxacin therapy. (See and .),

WARNINGSPRECAUTIONS, Pediatric UseADVERSE REACTIONS

• that diarrhea is a common problem caused by antibiotics which usually ends when the antibiotic is

discontinued. Sometimes after starting treatment with antibiotics, patients can develop watery and bloody stools (with or without stomach cramps and fever) even as late as two or more months after having taken the last dose of the antibiotic. If this occurs, patients should contact their physician as soon as possible.

Drug Interactions:

In a pharmacokinetic study, systemic exposure of tizanidine (4 mg single dose) was significantly increased (C 7-fold, AUC 10-fold) when the drug was given concomitantly with ciprofloxacin (500 mg bid for 3 days). The hypotensive and sedative effects of tizanidine were also potentiated. Concomitant administration of tizanidine and ciprofloxacin is contraindicated. As with some other quinolones, concurrent administration of ciprofloxacin with the ophylline may lead to elevated serum concentrations of the ophylline and prolongation of its elimination half-life. This may result in increased risk of theophylline-related adverse reactions. (See .) If concomitant use cannot be avoided, serum levels of theophylline should be monitored and dosage adjustments made as appropriate. Some quinolones, including ciprofloxacin, have also been shown to interfere with the metabolism of caffeine. This may lead to reduced clearance of caffeine and a prolongation of its serum half-life. Concurrent administration of a quinolone, including ciprofloxacin, with multivalent cation-containing products such as magnesium/aluminum antacids, sucralfate, Videx® (didanosine) chewable/buffered tablets or pediatric powder, other highly buffered drugs, or products containing calcium, iron, or zinc may substantially decrease its absorption, resulting in serum and urine levels considerably lower than desired. (See for concurrent administration of these agents with ciprofloxacin.) Histamine H -receptor antagonists appear to have no significant effect on the bioavailability of ciprofloxacin. Altered serum levels of phenytoin (increased and decreased) have been reported in patients receiving concomitant ciprofloxacin. The concomitant administration of ciprofloxacin with the sulfonylurea glyburide has, on rare occasions, resulted in severe hypoglycemia. Some quinolones, including ciprofloxacin, have been associated with transient elevations in serum creatinine in patients receiving cyclosporine concomitantly. Quinolones, including ciprofloxacin, have been reported to enhance the effects of the oral anticoagulant warfarin or its derivatives. When these products are administered concomitantly, prothrombin time or other suitable coagulation tests should be closely monitored. Probenecid interferes with renal tubular secretion of ciprofloxacin and produces an increase in the level of ciprofloxacin in the serum. This should be considered if patients are receiving both drugs concomitantly. Renal tubular transport of methotrexate may be inhibited by concomitant administration of ciprofloxacin potentially leading to increased plasma levels of methotrexate. This might increase the risk of methotrexate associated toxic reactions. Therefore, patients under methotrexate therapy should be carefully monitored when concomitant ciprofloxacin therapy is indicated. Metoclopramide significantly accelerates the absorption of oral ciprofloxacin resulting in shorter time to reach maximum plasma concentrations. No significant effect was observed on the bioavailability of ciprofloxacin. Nonsteroidal anti-inflammatory drugs (but not acetyl salicylic acid) in combination of very high doses of quinolones have been shown to provoke convulsions in pre-clinical studies.

max

WARNINGS

DOSAGE AND ADMINISTRATION

Carcinogenesis, Mutagenesis, Impairment of Fertility:

Eight mutagenicity tests have been conducted with ciprofloxacin, and the test results are listed below: Salmonella/Microsome Test (Negative) DNA Repair Assay (Negative) Mouse Lymphoma Cell Forward Mutation Assay (Positive) Chinese Hamster V Cell HGPRT Test (Negative) Syrian Hamster Embryo Cell Transformation Assay (Negative) Point Mutation Assay (Negative) Mitotic Crossover and Gene Conversion Assay (Negative) Rat Hepatocyte DNA Repair Assay (Positive) Thus, 2 of the 8 tests were positive, but results of the following 3 test systems gave negative results: Rat Hepatocyte DNA Repair Assay Micronucleus Test (Mice) Dominant Lethal Test (Mice) Long-term carcinogenicity studies in rats and mice resulted in no carcinogenic or tumorigenic effects due to ciprofloxacin at daily oral dose levels up to 250 and 750 mg/kg to rats and mice, respectively (approximately 1.7- and 2.5times the highest recommended therapeutic dose based upon mg/m). Results from photo cocarcinogenicity testing indicate that ciprofloxacin does not reduce the time to appearance of UVinduced skin tumors as compared to vehicle control. Hairless (Skh-1) mice were exposed to UVA light for 3.5 hours five times every two weeks for up to 78 weeks while concurrently being administered ciprofloxacin. The time to development of the first skin tumors was 50 weeks in mice treated concomitantly with UVA and ciprofloxacin (mouse dose approximately equal to maximum recommended human dose based upon mg/m), as opposed to 34 weeks when animals were treated with both UVA and vehicle. The times to development of skin tumors ranged from 16-32 weeks in mice treated concomitantly with UVA and other quinolones. In this model, mice treated with ciprofloxacin alone did not develop skin or systemic tumors. There are no data from similar models using pigmented mice and/or fully haired mice. The clinical significance of these findings to humans is unknown. Fertility studies performed in rats at oral doses of ciprofloxacin up to 100 mg/kg (approximately 0.7-times the highest recommended therapeutic dose based upon mg/m) revealed no evidence of impairment. in vitro

E. coli

79

Saccharomyces cerevisiae Saccharomyces cerevisiae

in vivo

2

Pregnancy

Teratogenic Effects

There are no adequate and well-controlled studies in pregnant women. An expert review of published data on experiences with ciprofloxacin use during pregnancy by TERIS - the Teratogen Information System - concluded that therapeutic doses during pregnancy are unlikely to pose a substantial teratogenic risk (quantity and quality of data=fair), but the data are insufficient to state that there is no risk. A controlled prospective observational study followed 200 women exposed to fluoroquinolones (52.5% exposed to ciprofloxacin and 68% first trimester exposures) during gestation. exposure to fluoroquinolones during embryogenesis was not associated with increased risk of major malformations. The reported rates of major congenital malformations were 2.2% for the fluoroquinolone group and 2.6% for the control group (background incidence of major malformations is 1-5%). Rates of spontaneous abortions, prematurity and low birth weight did not differ between the groups and there were no clinically significant musculoskeletal dysfunctions up to one year of age in the ciprofloxacin exposed children. Another prospective follow-up study reported on 549 pregnancies with fluoroquinolone exposure (93% first trimester exposures). There were 70 ciprofloxacin exposures, all within the first trimester. The malformation rates among live-born babies exposed to ciprofloxacin and to fluoroquinolones overall were both within background incidence ranges. No specific patterns of congenital abnormalities were found. The study did not reveal any clear adverse reactions due to exposure to ciprofloxacin. No differences in the rates of prematurity, spontaneous abortions, or birth weight were seen in women exposed to ciprofloxacin during pregnancy. However, these small postmarketing epidemiology studies, of which most experience is from short term, first trimester exposure, are insufficient to evaluate the risk for less common defects or to permit reliable and definitive conclusions regarding the safety of ciprofloxacin in pregnant women and their developing fetuses. Ciprofloxacin should not be used during pregnancy unless the potential benefit justifies the potential risk to both fetus and mother (see). Reproduction studies have been performed in rats and mice using oral doses up to 100 mg/kg (0.6 and 0.3 times the maximum daily human dose based upon body surface area, respectively) and have revealed no evidence of harm to the fetus due to ciprofloxacin. In rabbits, oral ciprofloxacin dose levels of 30 and 100 mg/kg (approximately 0.4- and 1.3-times the highest recommended therapeutic dose based upon mg/m) produced gastrointestinal toxicity resulting in maternal weight loss and an increased incidence of abortion, but no teratogenicity was observed at either dose level. After intravenous administration of doses up to 20 mg/kg (approximately 0.3-times the highest recommended therapeutic dose based upon mg/m) no maternal toxicity was produced and no embryotoxicity or teratogenicity was observed. (See .) Pregnancy Category C:

8

⁹In utero

¹⁰in utero

8,9WARNINGS

²²WARNINGS

Nursing Mothers:

Ciprofloxacin is excreted in human milk. The amount of ciprofloxacin absorbed by the nursing infant is unknown. Because of the potential for serious adverse reactions in infants nursing from mothers taking ciprofloxacin, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

Pediatric Use:

Ciprofloxacin, like other quinolones, causes arthropathy and histological changes in weight-bearing joints of juvenile animals resulting in lameness. (See .)

ANIMAL PHARMACOLOGY

Inhalational Anthrax (Post-Exposure)

Ciprofloxacin is indicated in pediatric patients for inhalational anthrax (post-exposure). The risk-benefit assessment indicates that administration of ciprofloxacin to pediatric patients is appropriate. For information regarding pediatric dosing in inhalational anthrax (post-exposure), see and .

DOSAGE AND ADMINISTRATIONINHALATIONAL ANTHRAX – ADDITIONAL INFORMATION

Complicated Urinary Tract Infection and Pyelonephritis

Ciprofloxacin is indicated for the treatment of complicated urinary tract infections and pyelonephritis due to . Although effective in clinical trials, ciprofloxacin is not a drug of first choice in the pediatric population due to an increased incidence of adverse events compared to the controls, including events related to joints and/or surrounding tissues. The rates of these events in pediatric patients with complicated urinary tract infection and pyelonephritis within six weeks of follow-up were 9.3% (31/335) versus 6% (21/349) for control agents. The rates of these events occurring at any time up to the one year follow-up were 13.7% (46/335) and 9.5% (33/349), respectively. The rate of all adverse events regardless of drug relationship at six weeks was 41% (138/335) in the ciprofloxacin arm compared to 31% (109/349) in the control arm. (See and .) *Escherichia coli*ADVERSE

REACTIONSCLINICAL STUDIES

Cystic Fibrosis

Short-term safety data from a single trial in pediatric cystic fibrosis patients are available. In a randomized, double-blind clinical trial for the treatment of acute pulmonary exacerbations in cystic fibrosis patients (ages 5-17 years), 67 patients received ciprofloxacin I.V. 10 mg/kg/dose q8h for one week followed by ciprofloxacin tablets 20 mg/kg/dose q12h to complete 10-21 days treatment and 62 patients received the combination of ceftazidime I.V. 50 mg/kg/dose q8h and tobramycin I.V. 3 mg/kg/dose q8h for a total of 10-21 days. Patients less than 5 years of age were not studied. Safety monitoring in the study included periodic range of motion examinations and gait assessments by treatment-blinded examiners. Patients were followed for an average of 23 days after completing treatment (range 0-93 days). This study was not designed to determine long term effects and the safety of repeated exposure to ciprofloxacin. Musculoskeletal adverse events in patients with cystic fibrosis were reported in 22% of the patients in the ciprofloxacin group and 21% in the comparison group. Decreased range of motion was reported in 12% of the subjects in the ciprofloxacin group and 16% in the comparison group. Arthralgia was reported in 10% of the patients in the ciprofloxacin group and 11% in the comparison group. Other adverse events were similar in nature and frequency between treatment arms. One of sixty-seven patients developed arthritis of the knee nine days after a ten day course of treatment with ciprofloxacin. Clinical symptoms resolved, but an MRI showed knee effusion without other abnormalities eight months after treatment. However, the relationship of this event to the patient's course of ciprofloxacin can not be definitively determined, particularly since patients with cystic fibrosis may develop arthralgias/arthritis as part of their underlying disease process.

Geriatric Use:

Geriatric patients are at increased risk for developing severe tendon disorders including tendon rupture

when being treated with a fluoroguinolone such as Ciprofloxacin Tablets USP, 250 mg, 500 mg and 750 mg. This risk is further increased in patients receiving concomitant corticosteroid therapy. Tendinitis or tendon rupture can involve the Achilles, hand, shoulder, or other tendon sites and can occur during or after completion of therapy; cases occurring up to several months after fluoroquinolone treatment have been reported. Caution should be used when prescribing Ciprofloxacin Tablets USP, 250 mg, 500 mg and 750 mg to elderly patients especially those on corticosteroids. Patients should be informed of this potential side effect and advised to discontinue Ciprofloxacin Tablets USP, 250 mg, 500 mg and 750 mg and contact their healthcare provider if any symptoms of tendinitis or tendon rupture occur (See and In a retrospective analysis of 23 multiple-dose controlled clinical trials of ciprofloxacin encompassing over 3500 ciprofloxacin treated patients, 25% of patients were greater than or equal to 65 years of age and 10% were greater than or equal to 75 years of age. No overall differences in safety or effectiveness were observed between these subjects and younger subjects, and other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals on any drug therapy cannot be ruled out. Ciprofloxacin is known to be substantially excreted by the kidney, and the risk of adverse reactions may be greater in patients with impaired renal function. No alteration of dosage is necessary for patients greater than 65 years of age with normal renal function. However, since some older individuals experience reduced renal function by virtue of their advanced age, care should be taken in dose selection for elderly patients, and renal function monitoring may be useful in these patients. (See and .) In general, elderly patients may be more susceptible to drug-associated effects on the QT interval. Therefore, precaution should be taken when using Ciprofloxacin Tablets USP 250 mg, 500 mg and 750 mg with concomitant drugs that can result in prolongation of the QT interval (e.g., class IA or class III antiarrhythmics) or in patients with risk factors for torsade de pointes (e.g., known QT prolongation, uncorrected

, , BOXED WARNINGWARNINGS). ADVERSE REACTIONS/Post-Marketing Adverse Event Reports

CLINICAL PHARMACOLOGYDOSAGE AND ADMINISTRATION

ADVERSE REACTIONS

Adverse Reactions in Adult Patients:

During clinical investigations with oral and parenteral ciprofloxacin, 49,038 patients received courses of the drug. Most of the adverse events reported were described as only mild or moderate in severity, abated soon after the drug was discontinued, and required no treatment. Ciprofloxacin was discontinued because of an adverse event in 1% of orally treated patients. The most frequently reported drug related events, from clinical trials of all formulations, all dosages, all drug-therapy durations, and for all indications of ciprofloxacin therapy were nausea (2.5%), diarrhea (1.6%), liver function tests abnormal (1.3%), vomiting (1%), and rash (1%). Additional medically important events that occurred in less than 1% of ciprofloxacin patients are listed below. BODY AS A WHOLE: headache, abdominal pain/discomfort, foot pain, pain, pain in extremities, injection site reaction (ciprofloxacin intravenous) CARDIOVASCULAR: palpitation, atrial flutter, ventricular ectopy, syncope, hypertension, angina pectoris, myocardial infarction, cardiopulmonary arrest, cerebral thrombosis, phlebitis, tachycardia, migraine, hypotension CENTRAL NERVOUS SYSTEM: restlessness, dizziness, lightheadedness, insomnia, nightmares, hallucinations, manic reaction, irritability, tremor, ataxia, convulsive seizures, lethargy, drowsiness, weakness, malaise, anorexia, phobia, depersonalization, depression, paresthesia, abnormal gait, grand mal convulsion GASTROINTESTINAL: painful oral mucosa, oral candidiasis, dysphagia, intestinal perforation, gastrointestinal bleeding, cholestatic jaundice, hepatitis HEMIC/LYMPHATIC: lymphadenopathy, petechia METABOLIC/NUTRITIONAL: amylase increase, lipase increase MUSCULOSKELETAL: arthralgia or back pain, joint stiffness, achiness, neck or chest pain, flare up of gout RENAL/UROGENITAL: interstitial nephritis, nephritis, renal failure, polyuria,

urinary retention, urethral bleeding, vaginitis, acidosis, breast pain RESPIRATORY: dyspnea, epistaxis, laryngeal or pulmonary edema, hiccough, hemoptysis, bronchospasm, pulmonary embolism SKIN/HYPERSENSITIVITY: allergic reaction, pruritus, urticaria, photosensitivity/phototoxicity reaction, flushing, fever, chills, angioedema, edema of the face, neck, lips, conjunctivae or hands, cutaneous candidiasis, hyperpigmentation, erythema nodosum, sweating SPECIAL SENSES: blurred vision, disturbed vision (change in color perception, overbrightness of lights), decreased visual acuity, diplopia, eye pain, tinnitus, hearing loss, bad taste, chromatopsia In several instances nausea, vomiting, tremor, irritability, or palpitation were judged by investigators to be related to elevated serum levels of theophylline possibly as a result of drug interaction with ciprofloxacin. In randomized, double-blind controlled clinical trials comparing ciprofloxacin tablets (500 mg BID) to cefuroxime axetil (250 mg-500 mg BID) and to clarithromycin (500 mg BID) in patients with respiratory tract infections, ciprofloxacin demonstrated a CNS adverse event profile comparable to the control drugs.

Adverse Reactions in Pediatric Patients:

Ciprofloxacin, administered I.V. and /or orally, was compared to a cephalosporin for treatment of complicated urinary tract infections (cUTI) or pyelonephritis in pediatric patients 1 to 17 years of age (mean age of 6 ± 4 years). The trial was conducted in the US, Canada, Argentina, Peru, Costa Rica, Mexico, South Africa, and Germany. The duration of therapy was 10 to 21 days (mean duration of treatment was 11 days with a range of 1 to 88 days). The primary objective of the study was to assess musculoskeletal and neurological safety within 6 weeks of therapy and through one year of follow-up in the 335 ciprofloxacin- and 349 comparator-treated patients enrolled. An Independent Pediatric Safety Committee (IPSC) reviewed all cases of musculoskeletal adverse events as well as all patients with an abnormal gait or abnormal joint exam (baseline or treatment-emergent). These events were

evaluated in a comprehensive fashion and included such conditions as arthralgia, abnormal gait, abnormal joint exam, joint sprains, leg pain, back pain, arthrosis, bone pain, pain, myalgia, arm pain, and decreased range of motion in a joint. The affected joints included: knee, elbow, ankle, hip, wrist, and shoulder. Within 6 weeks of treatment initiation, the rates of these events were 9.3% (31/335) in the ciprofloxacin-treated group versus 6% (21/349) in comparator-treated patients. The majority of these events were mild or moderate in intensity. All musculoskeletal events occurring by 6 weeks resolved (clinical resolution of signs and symptoms), usually within 30 days of end of treatment. Radiological evaluations were not routinely used to confirm resolution of the events. The events occurred more frequently in ciprofloxacin-treated patients than control patients, regardless of whether they received I.V. or oral therapy. Ciprofloxacin-treated patients were more likely to report more than one event and on more than one occasion compared to control patients. These events occurred in all age groups and the rates were consistently higher in the ciprofloxacin group compared to the control group. At the end of 1 year, the rate of these events reported at any time during that period was 13.7% (46/335) in the ciprofloxacin-treated group versus 9.5% (33/349) comparator-treated patients. An adolescent female discontinued ciprofloxacin for wrist pain that developed during treatment. An MRI performed 4 weeks later showed a tear in the right ulnar fibrocartilage. A diagnosis of overuse syndrome secondary to sports activity was made, but a contribution from ciprofloxacin cannot be excluded. The patient recovered by 4 months without surgical intervention. The incidence rates of neurological events within 6 weeks of treatment initiation were 3% (9/335) in the ciprofloxacin group versus 2% (7/349) in the comparator group and included dizziness, nervousness, insomnia, and somnolence. In this trial, the overall incidence rates of adverse events regardless of relationship to study drug and within 6 weeks of treatment initiation were 41% (138/335) in the ciprofloxacin group versus 31% (109/349) in the comparator group. The most frequent events were gastrointestinal: 15% (50/335) of ciprofloxacin patients compared to 9% (31/349) of comparator patients. Serious adverse events were seen in 7.5% (25/335) of ciprofloxacin-treated patients compared to 5.7% (20/349) of control patients. Discontinuation of drug due to an adverse event was observed in 3% (10/335) of ciprofloxacin-treated patients versus 1.4% (5/349) of comparator patients. Other adverse events that occurred in at least 1% of ciprofloxacin patients were diarrhea 4.8%, vomiting 4.8%, abdominal pain 3.3%, accidental injury 3%, rhinitis 3%, dyspepsia 2.7%, nausea 2.7%, fever 2.1%, asthma 1.8% and rash 1.8%. In addition to the events reported in pediatric patients in clinical trials, it should be expected that events reported in adults during clinical trials or post-marketing experience may also occur in pediatric patients.

Findings Involving Joint or Peri-articular Tissues as Assessed by the IPSC

	Ciprofloxacin	Comparator
All Patients (within 6 weeks)	31/335 (9.3%)	21/349 (6%)
95% Confidence Interval*	(-0.8%, -	+7.2%)
Age Group		
\geq 12 months < 24 months	1/36 (2.8%)	0/41
≥ 2 years < 6 years	5/124 (4%)	3/118 (2.5%)
≥ 6 years < 12 years	18/143 (12.6%)	12/153 (7.8%)
≥ 12 years to 17 years	7/32 (21.9%)	6/37 (16.2 %)
All Patients (within 1 year)	46/335 (13.7%)	33/349 (9.5%)
95% Confidence Interval*	(-0.6%, +9.1%)	

^{*}The study was designed to demonstrate that the arthropathy rate for the ciprofloxacin group did not exceed that of the control group by more than + 6%. At both the 6 week and 1 year evaluations, the 95% confidence interval indicated that it could not be concluded that ciprofloxacin group had findings

comparable to the control group.

Post-Marketing Adverse Event Reports:

The following adverse events have been reported from worldwide marketing experience with quinolones, including ciprofloxacin. Because these events are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure. Decisions to include these events in labeling are typically based on one or more of the following factors: (1) seriousness of the event, (2) frequency of the reporting, or (3) strength of causal connection to the drug. Agitation, agranulocytosis, albuminuria, anaphylactic reactions (including life-threatening anaphylactic shock), anosmia, candiduria, cholesterol elevation (serum), confusion, constipation, delirium, dyspepsia, dysphagia, erythema multiforme, exfoliative dermatitis, fixed eruption, flatulence, glucose elevation (blood), hemolytic anemia, hepatic failure (including fatal cases), hepatic necrosis, hyperesthesia, hypertonia, hypesthesia, hypotension (postural), jaundice, marrow depression (life threatening), methemoglobinemia, moniliasis (oral, gastrointestinal, vaginal), myalgia, myasthenia, myasthenia gravis (possible exacerbation), myoclonus, nystagmus, pancreatitis, pancytopenia (life threatening or fatal outcome), peripheral neuropathy, phenytoin alteration (serum), photosensitivity/phototoxicity reaction, potassium elevation (serum), prothrombin time prolongation or decrease, pseudomembranous colitis (The onset of pseudomembranous colitis symptoms may occur during or after antimicrobial treatment.), psychosis (toxic), renal calculi, serum sickness like reaction, Stevens-Johnson syndrome, taste loss, tendinitis, tendon rupture, torsade de pointes, toxic epidermal necrolysis (Lyell's Syndrome), triglyceride elevation (serum), twitching, vaginal candidiasis, and vasculitis. (See .) Adverse events were also reported by persons who received ciprofloxacin for anthrax post-exposure prophylaxis following the anthrax bioterror attacks of October 2001. (See also)

PRECAUTIONS

INHALATIONAL ANTHRAX - ADDITIONAL INFORMATION.

Adverse Laboratory Changes:

Changes in laboratory parameters listed as adverse events without regard to drug relationship are listed below: Hepatic — Elevations of ALT (SGPT) (1.9%), AST (SGOT) (1.7%), alkaline phosphatase (0.8%), LDH (0.4%), serum bilirubin (0.3%). Hematologic — Eosinophilia (0.6%), leukopenia (0.4%), decreased blood platelets (0.1%), elevated blood platelets (0.1%), pancytopenia (0.1%). Renal — Elevations of serum creatinine (1.1%), BUN (0.9%), CRYSTALLURIA, CYLINDRURIA, AND HEMATURIA HAVE BEEN REPORTED. Other changes occurring in less than 0.1% of courses were: elevation of serum gammaglutamyl transferase, elevation of serum amylase, reduction in blood glucose, elevated uric acid, decrease in hemoglobin, anemia, bleeding diathesis, increase in blood monocytes, leukocytosis.

OVERDOSAGE

In the event of acute overdosage, reversible renal toxicity has been reported in some cases. The stomach should be emptied by inducing vomiting or by gastric lavage. The patient should be carefully observed and given supportive treatment, including monitoring of renal function and administration of magnesium, aluminum, or calcium containing antacids which can reduce the absorption of ciprofloxacin. Adequate hydration must be maintained. Only a small amount of ciprofloxacin (< 10%) is removed from the body after hemodialysis or peritoneal dialysis. Single doses of ciprofloxacin were relatively nontoxic via the oral route of administration in mice, rats, and dogs. No deaths occurred within a 14-day post treatment observation period at the highest oral doses tested; up to 5000 mg/kg in either rodent species, or up to 2500 mg/kg in the dog. Clinical signs observed included hypoactivity and cyanosis in both rodent species and severe vomiting in dogs. In rabbits, significant mortality was seen at doses of ciprofloxacin > 2500 mg/kg. Mortality was delayed in these animals, occurring 10-14 days after dosing. In mice, rats, rabbits and dogs, significant toxicity including tonic/clonic convulsions was observed at intravenous doses of ciprofloxacin between 125 and 300 mg/kg.

DOSAGE AND ADMINISTRATION - ADULTS

Ciprofloxacin Tablets USP 250 mg, 500 mg and 750 mg should be administered orally to adults as described in the Dosage Guidelines table. The determination of dosage for any particular patient must take into consideration the severity and nature of the infection, the susceptibility of the causative organism, the integrity of the patient's host-defense mechanisms, and the status of renal function and hepatic function. The duration of treatment depends upon the severity of infection. The usual duration is 7 to 14 days; however, for severe and complicated infections more prolonged therapy may be required. Ciprofloxacin should be administered at least 2 hours before or 6 hours after magnesium/aluminum antacids, or sucralfate, Videx® (didanosine) chewable/buffered tablets or pediatric powder for oral solution, other highly buffered drugs, or other products containing calcium, iron or zinc. Patients whose therapy is started with CIPRO I.V. may be switched to Ciprofloxacin Tablets USP 250 mg, 500 mg and 750 mg when clinically indicated at the discretion of the physician (See and table below for the equivalent dosing regimens). Ciprofloxacin is eliminated primarily by renal excretion; however, the drug is also metabolized and partially cleared through the biliary system of the liver and through the intestine. These alternative pathways of drug elimination appear to compensate for the reduced renal excretion in patients with renal impairment. Nonetheless, some modification of dosage is recommended, particularly for patients with severe renal dysfunction. The following table provides dosage guidelines for use in patients with renal impairment: When only the serum creatinine concentration is known, the following formula may be used to estimate creatinine clearance.

Men: Creatinine clearance (mL/min) = 72 x serum creatinine (mg/dL) Women: 0.85 x the value calculated for men. The serum creatinine should represent a steady state of renal function. In patients with severe infections and severe renal impairment, a unit dose of 750 mg may be administered at the intervals noted above. Patients should be carefully monitored.

ADULT DOSAGE GUIDELINES					
Infection	Severity	Dose	Frequency	Usual Durations †	
Urinary Tract	Acute Uncomplicated	250	q 12 h	3 Days	
		mg			
	Mild/Moderate	250	q 12 h	7 to 14	
		mg		Days	
	Severe/Complicated	500	q 12 h	7 to 14	
		mg		Days	
Chronic Bacterial Prostatits	Mild/Moderate	500	q 12 h	28 Days	
		mg			
Lower Respiratory Tract	Mild/Moderate	500	q 12 h	7 to 14	
		mg		days	
	Severe/Complicated	750	q 12 h	7 to 14	
		mg		days	
Acute Sinusitis	Mild/Moderate	500	q 12 h	10 days	
		mg	_		
Skin and Skin Structure	Mild/Moderate	500	q 12 h	7 to 14	
		mg	_	Days	
	Severe/Complicated	750	q 12 h	7 to 14	
		mg		Days	
Bone and Joint	Mild/Moderate	500	q 12 h	≥4 to 6	
		mg		weeks	
	Severe/Complicated	750	q 12 h	≥4 to 6	
		mg		weeks	
Intra-Abdominal *	Complicated	500	q 12 h	7 to 14	
		mg		Days	
Infectious Diarrhea	Mild/Moderate/Severe	500	q 12 h	5 to 7 Days	
		mg	_	_	
Typhoid Fever	Mild/Moderate	500	q 12 h	10 Days	
		mg			
Urethral and Cervical Gonococcal	Uncomplicated	250	single	single dose	
Infections	_	mg	dose		
Inhalational anthrax (post-exposure) **		500	q 12 h	60 Days	
		mg			
		_			

used in conjunction with metronidazole Generally ciprofloxacin should be continued for at least 2 days after the signs and symptoms of infection have disappeared, except for inhalational anthrax (post-exposure). Drug administration should begin as soon as possible after suspected or confirmed exposure. This indication is based on a surrogate endpoint, ciprofloxacin serum concentrations achieved in humans, reasonably likely to predict clinical benefit. For a discussion of ciprofloxacin serum concentrations in various human populations, see .*

**

4

INHALATIONAL ANTHRAX - ADDITIONAL INFORMATION

Conversion of I.V. to Oral Dosing in Adults:

CLINICAL PHARMACOLOGY

Equivalent AUC Dosing Regimens

Cipro Oral Dosage	Equivalent Cipro I.V. Dosage
250 mg Tablet q 12 h	200 mg I.V. q 12 h
500 mg Tablet q 12 h	400 mg I.V. q 12 h
750 mg Tablet q 12 h	400 mg I.V. q 8 h

Adults with Impaired Renal Function:

RECOMMENDED STARTING AND MAINTENANCE DOSES FOR PATIENTS WITH IMPAIRED RENAL FUNCTION

Dose	
See Usual Dosage.	
250-500 mg q 12 h	
250-500 mg q 18 h	
250-500 mg q 24 h (after dialysis)	
2	

Weight (kg) x (140 - age)

DOSAGE AND ADMINISTRATION - PEDIATRICS

Ciprofloxacin Tablets USP 250 mg, 500 mg and 750 mg should be administered orally as described in the Dosage Guidelines table. An increased incidence of adverse events compared to controls, including events related to joints and/or surrounding tissues, has been observed. (See and .) Dosing and initial route of therapy (i.e., I.V. or oral) for complicated urinary tract infection or pyelonephritis should be determined by the severity of the infection. In the clinical trial, pediatric patients with moderate to severe infection were initiated on 6 to 10 mg/kg I.V. every 8 hours and allowed to switch to oral therapy (10 to 20 mg/kg every 12 hours), at the discretion of the physician. Pediatric patients with moderate to severe renal insufficiency were excluded from the clinical trial of complicated urinary tract infection and pyelonephritis. No information is available on dosing adjustments necessary for pediatric patients with moderate to severe renal insufficiency (i.e., creatinine clearance of < 50 mL/min/1.73m).

ADVERSE REACTIONSCLINICAL STUDIES

PEDIATRIC DOSAGE GUIDELINES

Infection	Route of Adminis tration	Dose (mg/kg)	Frequency	Total Duration
Complicated	Intravenous	6 to 10 mg/kg (maximum 400 mg per dose;	Every 8	10-21
Urinary Tract or Pyelonephritis		not to be exceeded even in patients weighing > 51 kg)	hours	days*
(patients from 1 to 17 years of age)	Oral	10 mg/kg to 20 mg/kg (maximum 750 mg per dose; not to be exceeded even in patients weighing > 51 kg)	Every 12 hours	
Inhalational Anthrax (Post-Exposure)**	Intravenous	10 mg/kg (maximum 400 mg per dose)	Every 12 hours	60 days
	Oral	15 mg/kg (maximum 500 mg per dose)	Every 12 hours	

^{*} The total duration of therapy for complicated urinary tract infection and pyelonephritis in the clinical trial was determined by the physician. The mean duration of treatment was 11 days (range 10 to 21 days). ** Drug administration should begin as soon as possible after suspected or confirmed exposure to spores. This indication is based on a surrogate endpoint, ciprofloxacin serum concentrations achieved in humans, reasonably likely to predict clinical benefit. For a discussion of ciprofloxacin serum concentrations in various human populations, see .

Bacillus anthracis

5

INHALATIONAL ANTHRAX - ADDITIONAL INFORMATION

2

ANIMAL PHARMACOLOGY

Ciprofloxacin and other quinolones have been shown to cause arthropathy in immature animals of most species tested. (See .) Damage of weight bearing joints was observed in juvenile dogs and rats. In young beagles, 100 mg/kg ciprofloxacin, given daily for 4 weeks, caused degenerative articular changes of the knee joint. At 30 mg/kg, the effect on the joint was minimal. In a subsequent study in young beagle dogs, oral ciprofloxacin doses of 30 mg/kg and 90 mg/kg ciprofloxacin (approximately 1.3- and 3.5-times the pediatric dose based upon comparative plasma AUCs) given daily for 2 weeks caused articular changes which were still observed by histopathology after a treatment-free period of 5 months. At 10 mg/kg (approximately 0.6-times the pediatric dose based upon comparative plasma AUCs), no effects on joints were observed. This dose was also not associated with arthrotoxicity after an additional treatment-free period of 5 months. In another study, removal of weight bearing from the joint reduced the lesions but did not totally prevent them. Crystalluria, sometimes associated with secondary nephropathy, occurs in laboratory animals dosed with ciprofloxacin. This is primarily related

to the reduced solubility of ciprofloxacin under alkaline conditions, which predominate in the urine of test animals; in man, crystalluria is rare since human urine is typically acidic. In rhesus monkeys, crystalluria without nephropathy was noted after single oral doses as low as 5 mg/kg. (approximately 0.07-times the highest recommended therapeutic dose based upon mg/m). After 6 months of intravenous dosing at 10 mg/kg/day, no nephropathological changes were noted; however, nephropathy was observed after dosing at 20 mg/kg/day for the same duration (approximately 0.2-times the highest recommended therapeutic dose based upon mg/m). In dogs, ciprofloxacin at 3 and 10 mg/kg by rapid I.V. injection (15 sec.) produces pronounced hypotensive effects. These effects are considered to be related to histamine release, since they are partially antagonized by pyrilamine, an antihistamine. In rhesus monkeys, rapid I.V. injection also produces hypotension but the effect in this species is inconsistent and less pronounced. In mice, concomitant administration of nonsteroidal anti-inflammatory drugs such as phenylbutazone and indomethacin with quinolones has been reported to enhance the CNS stimulatory effect of quinolones. Ocular toxicity seen with some related drugs has not been observed in ciprofloxacin-treated animals.

WARNINGS

22

CLINICAL STUDIES

Complicated Urinary Tract Infection and Pyelonephritis – Efficacy in Pediatric Patients:

NOTE: Although effective in clinical trials, ciprofloxacin is not a drug of first choice in the pediatric population due to an increased incidence of adverse events compared to controls, including events related to joints and/or surrounding tissues. Ciprofloxacin, administered I.V. and/or orally, was compared to a cephalosporin for treatment of complicated urinary tract infections (cUTI) and pyelonephritis in pediatric patients 1 to 17 years of age (mean age of 6 ± 4 years). The trial was conducted in the US, Canada, Argentina, Peru, Costa Rica, Mexico, South Africa, and Germany. The duration of therapy was 10 to 21 days (mean duration of treatment was 11 days with a range of 1 to 88 days). The primary objective of the study was to assess musculoskeletal and neurological safety. Patients were evaluated for clinical success and bacteriological eradication of the baseline organism(s) with no new infection or superinfection at 5 to 9 days post-therapy (Test of Cure or TOC). The Per Protocol population had a causative organism(s) with protocol specified colony count(s) at baseline, no protocol violation, and no premature discontinuation or loss to follow-up (among other criteria). The clinical success and bacteriologic eradication rates in the Per Protocol population were similar between ciprofloxacin and the comparator group as shown below.

Clinical Success and Bacteriologic Eradication at Test of Cure (5 to 9 Days Post-Therapy)

	Ciprofloxacin	Comparator
Randomized Patients	337	352
Per Protocol Patients	211	231

Clinical Response at 5 to 9 Days Post-Treatment	95.7%	92.6%
	(202/211)	(214/231)
	95% CI [-1.	.3%, 7.3%]
Bacteriologic Eradication by Patient at 5 to 9 Days Post-	84.4%	78.3%
Treatment*	(178/211)	(181/231)
	95% CI [-1.3	3%, 13.1%]
Bacteriologic Eradication of the Baseline Pathogen at 5 to 9 Days		
Post-Treatment		
Escherichia coli	156/178	161/179
	(88%)	(90%)

^{*} Patients with baseline pathogen(s) eradicated and no new infections or superinfections/total number of patients. There were 5.5% (6/211) ciprofloxacin and 9.5% (22/231) comparator patients with superinfections or new infections.

INHALATIONAL ANTHRAX IN ADULTS AND PEDIATRICS – ADDITIONAL INFORMATION

The mean serum concentrations of ciprofloxacin associated with a statistically significant improvement in survival in the rhesus monkey model of inhalational anthrax are reached or exceeded in adult and pediatric patients receiving oral and intravenous regimens. (See .) Ciprofloxacin pharmacokinetics have been evaluated in various human populations. The mean peak serum concentration achieved at steadystate in human adults receiving 500 mg orally every 12 hours is 2.97 µg/mL, and 4.56 µg/mL following 400 mg intravenously every 12 hours. The mean trough serum concentration at steady-state for both of these regimens is 0.2 µg/mL. In a study of 10 pediatric patients between 6 and 16 years of age, the mean peak plasma concentration achieved is 8.3 μg/mL and trough concentrations range from 0.09 to 0.26 μg/mL, following two 30-minute intravenous infusions of 10 mg/kg administered 12 hours apart. After the second intravenous infusion patients switched to 15 mg/kg orally every 12 hours achieve a mean peak concentration of 3.6 µg/mL after the initial oral dose. Long-term safety data, including effects on cartilage, following the administration of ciprofloxacin to pediatric patients are limited. (For additional information, see .) Ciprofloxacin serum concentrations achieved in humans serve as a surrogate endpoint reasonably likely to predict clinical benefit and provide the basis for this indication. A placebo-controlled animal study in rhesus monkeys exposed to an inhaled mean dose of 11 LD (~5.5 x 10 spores (range 5-30 LD) of was conducted. The minimal inhibitory concentration (MIC) of ciprofloxacin for the anthrax strain used in this study was 0.08 µg/mL. In the animals studied, mean serum concentrations of ciprofloxacin achieved at expected T (1 hour post-dose) following oral dosing to steady-state ranged from 0.98 to 1.69 µg/mL. Mean steady-state trough concentrations at 12 hours post-dose ranged from 0.12 to 0.19 µg/mL. Mortality due to anthrax for animals that received a 30-day regimen of oral ciprofloxacin beginning 24 hours post-exposure was significantly lower (1/9), compared to the placebo group (9/10) [p=0.001]. The one ciprofloxacin-treated animal that died of anthrax did so following the 30-day drug administration period. More than 9300 persons were recommended to complete a minimum of 60 days of antibiotic prophylaxis against possible inhalational exposure to during 2001. Ciprofloxacin was recommended to most of those individuals for all or part of the prophylaxis regimen. Some persons were also given anthrax vaccine or were switched to alternative antibiotics. No one who received ciprofloxacin or other therapies as prophylactic treatment subsequently developed inhalational anthrax. The number of persons who received ciprofloxacin as all or part of their post-exposure prophylaxis regimen is unknown. Among the persons surveyed by the Centers for Disease Control and Prevention, over 1000 reported receiving ciprofloxacin as sole postexposure prophylaxis for inhalational anthrax. Gastrointestinal adverse events (nausea, vomiting,

diarrhea, or stomach pain), neurological adverse events (problems sleeping, nightmares, headache, dizziness or lightheadedness) and musculoskeletal adverse events (muscle or tendon pain and joint swelling or pain) were more frequent than had been previously reported in controlled clinical trials. This higher incidence, in the absence of a control group, could be explained by a reporting bias, concurrent medical conditions, other concomitant medications, emotional stress or other confounding factors, and/or a longer treatment period with ciprofloxacin. Because of these factors and limitations in the data collection, it is difficult to evaluate whether the reported symptoms were drug-related.

DOSAGE AND ADMINISTRATIONPRECAUTIONS, Pediatric Use⁵

 50^{5} 50B. anthracis 67

B. anthracis

REFERENCES:

- 1. National Committee for Clinical Laboratory Standards, -Fifth Edition. Approved Standard NCCLS Document M7-A5, Vol. 20, No. 2, NCCLS, Wayne, PA, January, 2000. Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow Aerobically
- 2. Clinical and Laboratory Standards Institute, , CLSI Document M45-A Vol. 26, No. 19, CLSI, Wayne, PA, 2006. Methods for Antimicrobial Dilution and Disk Susceptibility Testing of Infrequently Isolated or Fastidious Bacteria; Approved Guideline.,
- 3. National Committee for Clinical Laboratory Standards, -Seventh Edition. Approved Standard NCCLS Document M2-A7, Vol. 20, No. 1, NCCLS, Wayne, PA, January, 2000. <u>Performance Standards for Antimicrobial Disk Susceptibility Tests</u>
- 4. Report presented at the FDA's Anti-Infective Drug and Dermatological Drug Product's Advisory Committee meeting, March 31, 1993, Silver Spring, MD. Report available from FDA, CDER, Advisors and Consultants Staff, HFD-21, 1901 Chapman Avenue, Room 200, Rockville, MD 20852, USA.
- 5. 21 CFR 314.510 (Subpart H Accelerated Approval of New Drugs for Life-Threatening Illnesses).
- 6. Kelly DJ, et al. Serum concentrations of penicillin, doxycycline, and ciprofloxacin during prolonged therapy in rhesus monkeys. J Infect Dis 1992; 166:1184-7.
- 7. Friedlander AM, et al. Postexposure prophylaxis against experimental inhalational anthrax. J Infect Dis 1993; 167:1239-42.
- 8. Friedman J, Polifka J. Teratogenic effects of drugs: a resource for clinicians (TERIS). Baltimore, Maryland: Johns Hopkins University Press, 2000:149-195.
- 9. Loebstein R, Addis A, Ho E, et al. Pregnancy outcome following gestational exposure to fluoroquinolones: a multicenter prospective controlled study. Antimicrob Agents Chemother. 1998;42(6):1336-1339.
- 10. Schaefer C, Amoura-Elefant E, Vial T, et al. Pregnancy outcome after prenatal quinolone exposure. Evaluation of a case registry of the European network of teratology information services (ENTIS). Eur J Obstet Gynecol Reprod Biol. 1996;69:83-89.

Ciprofloxacin Tablets USP, 250 mg, 500 mg and 750 mg

Read the Medication Guide that comes with Ciprofloxacin Tablets USP before you start taking it and each time you get a refill. There may be new information. This Medication Guide does not take the place of talking to your healthcare provider about your medical condition or your treatment.

What is the most important information I should know about Ciprofloxacin Tablets USP?

Ciprofloxacin Tablets USP belongs to a class of antibiotics called fluoroquinolones. Ciprofloxacin Tablets USP can cause side effects that may be serious or even cause death. If you get any of the following serious side effects, get medical help right away. Talk with your healthcare provider about whether you should continue to take Ciprofloxacin Tablets USP.

Tendon rupture or swelling of the tendon (tendinitis)

- Tendons are tough cords of tissue that connect muscles to bones.
- Pain, swelling, tears and inflammation of tendons including the back of the ankle (Achilles), shoulder, hand, or other tendon sites can happen in people of all ages who take fluoroquinolone antibiotics, including Ciprofloxacin Tablets USP. The risk of getting tendon problems is higher if you:
 - are over 60 years of age
 - are taking steroids (corticosteroids)
 - have had a kidney, heart or lung transplant
- Swelling of the tendon (tendinitis) and tendon rupture (breakage) have also happened in patients who take fluoroquinolones who do not have the above risk factors.
- Other reasons for tendon ruptures can include:
 - physical activity or exercise
 - kidney failure
 - tendon problems in the past, such as in people with rheumatoid arthritis (RA)
- Call your healthcare provider right away at the first sign of tendon pain, swelling or inflammation. Stop taking Ciprofloxacin Tablets USP until tendinitis or tendon rupture has been ruled out by your healthcare provider. Avoid exercise and using the affected area. The most common area of pain and swelling is the Achilles tendon at the back of your ankle. This can also happen with other tendons. Talk to your healthcare provider about the risk of tendon rupture with continued use of Ciprofloxacin Tablets USP. You may need a different antibiotic that is not a fluoroquinolone to treat your infection.
- Tendon rupture can happen while you are taking or after you have finished taking Ciprofloxacin Tablets USP. Tendon ruptures have happened up to several months after patients have finished taking their fluoroguinolone.
- Get medical help right away if you get any of the following signs or symptoms of a tendon rupture:
 - hear or feel a snap or pop in a tendon area
 - bruising right after an injury in a tendon area
 - unable to move the affected area or bear weight
- See the section for more information about side effects. "What are the possible side effects of Ciprofloxacin Tablets USP?"

What is Ciprofloxacin Tablets USP?

Ciprofloxacin Tablets USP is a fluoroquinolone antibiotic medicine used to treat certain infections caused by certain germs called bacteria.

Children less than 18 years of age have a higher chance of getting bone, joint, or tendon (musculoskeletal) problems such as pain or swelling while taking Ciprofloxacin Tablets USP. Ciprofloxacin Tablets USP should not be used as the first choice of antibiotic medicine in children under 18 years of age.

Ciprofloxacin Tablets USP should not be used in children under 18 years old, except to treat specific serious infections, such as complicated urinary tract infections and to prevent anthrax disease after breathing the anthrax bacteria germ (inhalational exposure). It is not known if Ciprofloxacin Extended Release Tablets are safe and work in children under 18 years of age.

Sometimes infections are caused by viruses rather than by bacteria. Examples include viral infections in the sinuses and lungs, such as the common cold or flu. Antibiotics, including Ciprofloxacin Tablets

USP, do not kill viruses.

Call your healthcare provider if you think your condition is not getting better while you are taking Ciprofloxacin Tablets USP.

Who should not take Ciprofloxacin Tablets USP?

Do not take Ciprofloxacin Tablets USP if you:

- have ever had a severe allergic reaction to an antibiotic known as a fluoroquinolone, or are allergic to any of the ingredients in Ciprofloxacin Tablets USP. Ask your healthcare provider if you are not sure. See the list of ingredients in Ciprofloxacin Tablets USP at the end of this Medication Guide.
- also take a medicine called tizanidine (Zanaflex®). Serious side effects from tizanidine are likely to happen.

What should I tell my healthcare provider before taking Ciprofloxacin Tablets USP?

See "What is the most important information I should know about Ciprofloxacin Tablets USP?"

Tell your healthcare provider about all your medical conditions, including if you:

- have tendon problems
- have central nervous system problems (such as epilepsy)
- have nerve problems
- have or anyone in your family has an irregular heartbeat, especially a condition called "QT prolongation"
- have a history of seizures
- have kidney problems. You may need a lower dose of Ciprofloxacin Tablets USP if your kidneys do not work well.
- have rheumatoid arthritis (RA) or other history of joint problems
- have trouble swallowing pills
- are pregnant or planning to become pregnant. It is not known if Ciprofloxacin Tablets USP will harm your unborn child.
- are breast-feeding or planning to breast-feed. Ciprofloxacin Tablets USP passes into breast milk.
 You and your healthcare provider should decide whether you will take Ciprofloxacin Tablets USP or breast-feed.

, including prescription and non-prescription medicines, vitamins and herbal and dietary supplements. Ciprofloxacin Tablets USP and other medicines can affect each other causing side effects. Especially tell your healthcare provider if you take: **Tell your healthcare provider about all the medicines you take**

- an NSAID (Non-Steroidal Anti-Inflammatory Drug). Many common medicines for pain relief are NSAIDs. Taking an NSAID while you take Ciprofloxacin Tablets USP or other fluoroquinolones may increase your risk of central nervous system effects and seizures. See "What are the possible side effects of Ciprofloxacin Tablets USP?"
- a blood thinner (warfarin, Coumadin®, Jantoven®)
- tizanidine (Zanaflex®). You should not take Ciprofloxacin Tablets USP if you are already taking tizanidine. See "Who should not take Ciprofloxacin Tablets USP?"
- theophylline (Theo-24®, Elixophyllin®, Theochron®, Uniphyl®, Theolair®)
- glyburide (Micronase®, Glynase®, Diabeta®, Glucovance®). See "What are the possible side effects of Ciprofloxacin Tablets USP?"
- phenytoin (Fosphenytoin Sodium®, Cerebyx®, Dilantin-125®, Dilantin®, Extended Phenytoin Sodium®, Prompt Penytoin Sodium®, Phenytek®)
- products that contain caffeine
- a medicine to control your heart rate or rhythm (antiarrhythmics) See "What are the possible side effects of Ciprofloxacin Tablets USP?"

- an anti-psychotic medicine
- a tricyclic antidepressant
- a water pill (diuretic)
- a steroid medicine. Corticosteroids taken by mouth or by injection may increase the chance of tendon injury. See "What is the most important information I should know about Ciprofloxacin Tablets USP?"
- methotrexate (Trexall®)
- Probenecid (Probalan®, Col-probenecid®)
- Metoclopromide (Reglan®, Reglan ODT®)
- Certain medicines may keep Ciprofloxacin Tablets USP from working correctly. Take Ciprofloxacin Tablets USP either 2 hours before or 6 hours after taking these products:
 - an antacid, multivitamin, or other product that has magnesium, calcium, aluminum, iron, or zinc
 - sucralfate (Carafate®)
 - didanosine (Videx®, Videx EC®)

Ask your healthcare provider if you are not sure if any of your medicines are listed above.

Know the medicines you take. Keep a list of your medicines and show it to your healthcare provider and pharmacist when you get a new medicine.

How should I take Ciprofloxacin Tablets USP?

- Take Ciprofloxacin Tablets USP exactly as prescribed by your healthcare provider.
- Take Ciprofloxacin Tablets USP in the morning and evening at about the same time each day.
 Swallow the tablet whole. Do not split, crush or chew the tablet. Tell your healthcare provider if you can not swallow the tablet whole.
- Ciprofloxacin Tablets USP can be taken with or without food.
- Ciprofloxacin Tablets USP should not be taken with dairy products (like milk or yogurt) or calcium-fortified juices alone, but may be taken with a meal that contains these products.
- Drink plenty of fluids while taking Ciprofloxacin Tablets USP.
- Do not skip any doses, or stop taking Ciprofloxacin Tablets USP even if you begin to feel better, until you finish your prescribed treatment, unless:
 - you have tendon effects (see), "What is the most important information I should know about Ciprofloxacin Tablets USP?"
 - you have a serious allergic reaction (see), or "What are the possible side effects
 of Ciprofloxacin Tablets USP?"
 - your healthcare provider tells you to stop.
- This will help make sure that all of the bacteria are killed and lower the chance that the bacteria will become resistant to Ciprofloxacin Tablets USP. If this happens, Ciprofloxacin Tablets USP and other antibiotic medicines may not work in the future.
- If you miss a dose of Ciprofloxacin Tablets USP, take it as soon as you remember. Do not take two doses at the same time, and do not take more than two doses in one day.
- If you take too much, call your healthcare provider or get medical help immediately.

If you have been prescribed Ciprofloxacin Tablets USP after being exposed to anthrax:

- Ciprofloxacin Tablets USP has been approved to lessen the chance of getting anthrax disease or worsening of the disease after you are exposed to the anthrax bacteria germ.
- Take Ciprofloxacin Tablets USP exactly as prescribed by your healthcare provider. Do not stop taking Ciprofloxacin Tablets USP without talking with your healthcare provider. If you stop taking Ciprofloxacin Tablets USP too soon, it may not keep you from getting the anthrax disease.
- Side effects may happen while you are taking Ciprofloxacin Tablets USP. When taking your Ciprofloxacin Tablets USP to prevent anthrax infection, you and your healthcare provider should talk about whether the risks of stopping Ciprofloxacin Tablets USP too soon are more important than

- the risks of side effects with Ciprofloxacin Tablets USP.
- If you are pregnant, or plan to become pregnant while taking Ciprofloxacin Tablets USP, you and your healthcare provider should decide whether the benefits of taking Ciprofloxacin Tablets USP for anthrax are more important than the risks.

What should I avoid while taking Ciprofloxacin Tablets USP?

- Ciprofloxacin Tablets USP can make you feel dizzy and lightheaded. Do not drive, operate
 machinery, or do other activities that require mental alertness or coordination until you know how
 Ciprofloxacin Tablets USP affects you.
- Avoid sunlamps, tanning beds, and try to limit your time in the sun. Ciprofloxacin Tablets USP can make your skin sensitive to the sun (photosensitivity) and the light from sunlamps and tanning beds. You could get severe sunburn, blisters or swelling of your skin. If you get any of these symptoms while taking Ciprofloxacin Tablets USP, call your healthcare provider right away. You should use a sunscreen and wear a hat and clothes that cover your skin if you have to be in sunlight.

What are the possible side effects of Ciprofloxacin Tablets USP?

• Ciprofloxacin Tablets USP can cause side effects that may be serious or even cause death. See "What is the most important information I should know about Ciprofloxacin Tablets USP?"

Other serious side effects of Ciprofloxacin Tablets USP include:

- Seizures have been reported in people who take fluoroquinolone antibiotics including Ciprofloxacin Tablets USP. Tell your healthcare provider if you have a history of seizures. Ask your healthcare provider whether taking Ciprofloxacin Tablets USP will change your risk of having a seizure. Central Nervous System (CNS) side effects may happen as soon as after taking the first dose of Ciprofloxacin Tablets USP. Talk to your healthcare provider right away if you get any of these side effects, or other changes in mood or behavior: Central Nervous System effects:
 - feel dizzy
 - seizures
 - hear voices, see things, or sense things that are not there (hallucinations)
 - feel restless
 - tremors
 - feel anxious or nervous
 - confusion
 - depression
 - trouble sleeping
 - nightmares
 - feel more suspicious (paranoia)
 - suicidal thoughts or acts
- Allergic reactions can happen in people taking fluoroquinolones, including Ciprofloxacin Tablets USP, even after only one dose. Stop taking Ciprofloxacin Tablets USP and get emergency medical help right away if you get any of the following symptoms of a severe allergic reaction: **Serious allergic reactions:**
 - hives
 - trouble breathing or swallowing
 - swelling of the lips, tongue, face
 - throat tightness, hoarseness
 - rapid heartbeat
 - faint
 - yellowing of the skin or eyes. Stop taking Ciprofloxacin Tablets USP and tell your healthcare provider right away if you get yellowing of your skin or white part of your eyes, or if you have dark urine. These can be signs of a serious reaction to Ciprofloxacin Tablets USP (a liver

problem).

- Skin rash may happen in people taking Ciprofloxacin Tablets USP even after only one dose. Stop taking Ciprofloxacin Tablets USP at the first sign of a skin rash and call your healthcare provider. Skin rash may be a sign of a more serious reaction to Ciprofloxacin Tablets USP. **Skin rash**
- (QT prolongation and torsade de pointes) Tell your healthcare provider right away if you have a change in your heart beat (a fast or irregular heartbeat), or if you faint. Ciprofloxacin Tablets USP may cause a rare heart problem known as prolongation of the QT interval. This condition can cause an abnormal heartbeat and can be very dangerous. The chances of this event are higher in people: Serious heart rhythm changes
 - who are elderly
 - with a family history of prolonged QT interval
 - with low blood potassium (hypokalemia)
 - who take certain medicines to control heart rhythm (antiarrhythmics)
- (Pseudomembranous colitis) Pseudomembranous colitis can happen with most antibiotics, including Ciprofloxacin Tablets USP. Call your healthcare provider right away if you get watery diarrhea, diarrhea that does not go away, or bloody stools. You may have stomach cramps and a fever. Pseudomembranous colitis can happen 2 or more months after you have finished your antibiotic. **Intestine infection**
- (Peripheral Neuropathy) Damage to the nerves in arms, hands, legs, or feet can happen in people
 who take fluoroquinolones, including Ciprofloxacin Tablets USP. Talk with your healthcare
 provider right away if you get any of the following symptoms of peripheral neuropathy in your
 arms, hands, legs, or feet: Ciprofloxacin tablets may need to be stopped to prevent permanent nerve
 damage. Changes in sensation and possible nerve damage
 - pain
 - burning
 - tingling
 - numbness
 - weakness
- (hypoglycemia) People who take Ciprofloxacin Tablets USP and other fluoroquinolone medicines
 with the oral anti-diabetes medicine glyburide (Micronase, Glynase, Diabeta, Glucovance) can get
 low blood sugar (hypoglycemia) which can sometimes be severe. Tell your healthcare provider if
 you get low blood sugar with CIPRO. Your antibiotic medicine may need to be changed. Low blood
 sugar
- (photosensitivity) See **Sensitivity to sunlight**"What should I avoid while taking Ciprofloxacin Tablets USP?"
- Increased chance of problems with joints and tissues around joints in children under 18 years old. Tell your child's healthcare provider if your child has any joint problems during or after treatment with Ciprofloxacin Tablets USP. The most common side effects of Ciprofloxacin Tablets USP include: **Joint Problems**
 - nausea
 - headache
 - diarrhea
 - vomiting
 - vaginal yeast infection
 - changes in liver function tests
 - pain or discomfort in the abdomen

These are not all the possible side effects of Ciprofloxacin Tablets USP. Tell your healthcare provider about any side effect that bothers you, or that does not go away.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store Ciprofloxacin Tablets USP?

• Store Ciprofloxacin Tablets USP below 86°F (30°C)

Keep Ciprofloxacin Tablets USP and all medicines out of the reach of children.

General Information about Ciprofloxacin Tablets USP

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use Ciprofloxacin Tablets USP for a condition for which it is not prescribed. Do not give Ciprofloxacin Tablets USP to other people, even if they have the same symptoms that you have. It may harm them.

This Medication Guide summarizes the most important information about Ciprofloxacin Tablets USP. If you would like more information about CIPRO, talk with your healthcare provider. You can ask your healthcare provider or pharmacist for information about Ciprofloxacin Tablets USP that is written for healthcare professionals.

For more information 1-(866)-562-4597

What are the ingredients in Ciprofloxacin Tablets USP?

- Active ingredient: ciprofloxacin
- Inactive ingredients: pregelatinized starch, microcrystalline cellulose, colloidal silicon dioxide, crospovidone, magnesium stearate, hypromellose, titanium dioxide, polyethylene glycol and purified water

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Manufactured by: (A Div. of J. B. Chemicals & Pharmaceuticals Ltd) Mumbai 400 030, India **Unique Pharmaceutical Laboratories**

Distributed by: Buffalo Grove, IL 60089 USA

PACK Pharmaceuticals, LLC,

CIPROFLOXACIN TABLET

NDC: 50436-3135-5 CIPROFLOXACIN HCL

250 MG 14 TAB

WARNING: KEEP OUT OF REACH OF CHILDREN LOT: XXXXXX EXP:XXXXXXX STORE AT 20-25°C (68-77°F) CONTROLLED ROOM TEMPERATURE

MFG BY: PACK PHARM XXXXXXX

MFG NDC: 16571-0411-10 MFG LOT: XXXXXXX

Pkg by: Unit Dose Services, LLC Miami, FL 33179

DRUG:CIPROFLOXACIN LOT: XXXXXXX EXP:XXXXXXX NDC:50436-314 TAB DRUG:CIPROFLOXACIN 250 MG LOT: XXXXXXX EXP: XXXXXXX

NDC:50436-3135-5 DRUG: CIPROFLOXACIN

NDC:50436-314 TAB

HCL

LOT: XXXXXXX EXP: XXXXXXX

NDC:50436-314 TAB DRUG: CIPROFLOXACIN 250 MG LOT: XXXXXXX EXP:XXXXXXX ciprofloxacin tablet

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG LABEL	Item Code (Source)	NDC:50436- 3135(NDC:16571-411)
Route of Administration	ORAL	DEA Schedule	

Active Ingredient/Active Moiety			
Ingredient Name Basis of Strength Strength			
ciprofloxacin hydrochloride (ciprofloxacin)	cipro flo xa cin	250 mg	

Inactive Ingredients			
Ingredient Name	Strength		
STARCH, CORN			
CELLULO SE, MICRO CRYSTALLINE			
SILICON DIO XIDE			
CROSPOVIDONE			
MAGNESIUM STEARATE			
HYPROMELLOSES			
TITANIUM DIO XIDE			
POLYETHYLENE GLYCOLS			
WATER			

Product Characteristics				
Color	white (white to slightly yellowish)	Score	no score	
Shape	ROUND	Size	11mm	
Flavor		Imprint Code	P;250	
Contains				

F	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:50436-3135-3	21 in 1 BOTTLE			
2	NDC:50436-3135-4	10 in 1 BOTTLE			
3	NDC:50436-3135-5	14 in 1 BOTTLE			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076639	09/10/2004	

Labeler - Unit Dose Services (831995316)

Registrant - Unit Dose Services (831995316)

Establishment			
Name	Address	ID/FEI	Business Operations
Unit Dose Services		831995316	REPACK(50436-3135)

Revised: 1/2011 Unit Dose Services