MV-502- iodine solution Michigan Veterinary Supply

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

MV-502

Michigan Veterinary Supply

MV-502

Pre and Post Iodine Teat Dip

IAn Aid in Reducing the Spread of Mastitis Causing Organisms

ACTIVE INGREDIENT: 0.5% titratable iodine

Contains 5% Glycerine

DIRECTIONS FOR PRE-DIPPING

- 1. Remove excessive organic soils from teats prior to the application of teat dip solution.
- 2. Dip or spray cow's teats thoroughly with **MV-502**.
- 3. Strip out 3-4 squirts of milk from each quarter.
- 4. To remove the Iodine Pre-Dip, wipe the teats dry with a single service towel.
- 5. Attach milking unit.

DIRECTIONS FOR POST DPPING

Immediately after milking use **MV-502** at full strength. Submerge entire teat in **MV-502** solution. Allow to air dry. Do not wipe. Always use fresh, full strength **MV-502**. If product in dip cup becomes visibly dirty, discard contents and replenish with **MV-502** undiluted product.

Do not reuse or return used product to the original container.

If frozen, thaw completely and mix well before using.

DANGER

KEEP OUT OF REACH OF CHILDREN

CAN CAUSE EYE DAMAGE.

Protect eyes when handling. DO NOT get in eyes or on clothing. Harmful is swallowed. Avoid contamination of food.

FIRST AID

Eyes: In case of contact with eyes, flush immediately with plenty of water for at least 15 minutes. Call a physician.

Internal: If swallowed, promptly rinse mouth with water then give a large quantity of milk or water. Avoid alcohol. Contact a physician immediately. Never give anything by mouth to an unconscious person.

FOR ASSISTANCE WITH MEDICAL EMERGENCY, CALL 1-800-367-5178 REFER TO MATERIAL DATA SAFETY SHEET.

PRINCIPAL DISPLAY PANEL/REPRESENTATIVE LABEL

Michigan Veterinary Supply

MV-502

Pre and Post Iodine Teat Dip

An Aid in Reducing the Spread of Mastitis Causing Organisms

ACTIVE INGREDIENT: 0.5% titratable iodine Contains 5% Glycerine

TAKE TIME OBSERVE LABEL DIRECTIONS

NET CONTENTS: 55 US GALLONS

Manufactured for: Michigan Veterinary Farm Supply Co., Inc. 7415 Lawrence Highway,

Vermontville, MI 49096



TDR-73E-VFS

DIRECTIONS FOR PRE-DIPPING

- Remove excess organic soils from udder and toats prior to foremilking.
- 2. Dip or spray cow's touts thoroughly with MV-502.
- 3. Strip out 3-4 squirts of milk from outh quarter.
 4. To remove the Iodine Pre-Dip, wipe the touts dry
- 5. Attach milking unit.

DIRECTIONS FOR POST-DIPPING

diately after milking use MV-502 at full strength. Submerge entire test in MY-502 solutio Allow to sir dry. Do not wipe. Always use fresh, fell strength MY-502. If product in dip cup becom visitly dirty, dissard contents and replenish with MY-502 undilused product.

Do not reuse or return used po to the original container.



An Aid in Reducing the Spread of Mastitis Causing Organisms

ACTIVE INGREDIENT: 0.5% titratable iodine Contains 5% Glycerine





NET CONTENTS: 55 U.S. GALLONS

DANGER

KEEP OUT OF REACH OF CHILDREN

Can Cause Eye Damage.

s when handling. DO NOT get in eyes or on swallowed. Avoid contemination of food

FIRST AID

Eves: In case of contact with eyes, flush imm

water for at least 15 minutes. Call a physician have been a common to the process of the physician have been a common to the process of the p

FOR ASSISTANCE WITH MEDICAL EMERGENCY. CALL 1-800-367-5178

REFER TO MATERIAL SAFETY DATA SHEET.



MV-502 (103)

Manufactured for: Michigan Veterinary Farm Supply Co., Inc. 7415 Lawrence Highway, Vermontville, MI 49096



MV-502

iodine solution

| Product Information | | | | | |
|-------------------------|-----------------------|--------------------|---------------|--|--|
| Product Type | OTC ANIMAL DRUG LABEL | Item Code (Source) | NDC:62183-730 | | |
| Route of Administration | TOPICAL | DEA Schedule | | | |

| Active Ingredient/Active Moiety | | | | | |
|---------------------------------|-------------------|------------|--|--|--|
| Ingredient Name | Basis of Strength | Strength | | | |
| IO DINE (IO DINE) | IODINE | 5 g in 1 L | | | |

| P | Packaging | | | | | | |
|---|------------------|---------------------|----------------------|--------------------|--|--|--|
| # | Item Code | Package Description | Marketing Start Date | Marketing End Date | | | |
| 1 | NDC:62183-730-01 | 3.8 L in 1 BOTTLE | | | | | |
| 2 | NDC:62183-730-05 | 18.9 L in 1 DRUM | | | | | |
| 3 | NDC:62183-730-15 | 56.8 L in 1 DRUM | | | | | |
| 4 | NDC:62183-730-55 | 208.2 L in 1 DRUM | | | | | |
| 5 | NDC:62183-730-75 | 1041 L in 1 TANK | | | | | |

| Marketing Information | | | | | |
|-----------------------|--|----------------------|--------------------|--|--|
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date | | |
| unapproved drug other | | 0 1/3 1/2 0 0 3 | | | |
| | | | | | |

Labeler - Michigan Veterinary Supply (020909198)

Revised: 1/2013 Michigan Veterinary Supply