

DRUG FACTS

Active ingredient(s)

Purpose

Use(s)

- temporarily relieves these symptoms of hay fever and the common cold:
- runny nose
- sneezing
- headache
- minor aches and pain
- nasal congestion
- sore throat
- temporarily relieves these additional symptoms of hay fever:
- itching of the nose or throat
- itchy, watery eyes
- temporarily reduces fever
- temporarily relieves these symptoms due to the common cold:
- nasal congestion
- headache
- minor aches and pains
- cough
- sore throat
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive
- temporarily reduces fever

Warnings

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product

- do not exceed recommended dosage**
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain or nasal congestion gets worse or lasts more than 7 days
- redness or swelling is present
- fever gets worse or lasts more than 3 days
- new symptoms occur

These could be signs of a serious condition.

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding

Keep out of reach of children

Keep out of reach of children.

Keep out of reach of children.

Directions

- do not use more than directed (see overdose warning)

adults and children 12 years and over

- take 2 caplets every 4 hours
- do not take more than 12 caplets in 24 hours

children under 12 years	<ul style="list-style-type: none"> do not use this adult product in children under 12 years of age; this will provide more than the recommended dose (overdose) and may cause liver damage
adults and children 12 years and over	<ul style="list-style-type: none"> do not use more than directed (see overdose warning) take 2 caplets every 4 hours do not take more than 12 caplets in 24 hours
children under 12 years	<ul style="list-style-type: none"> do not use this adult product in children under 12 years of age; this will provide more than the recommended dose (overdose) and may cause liver damage

Other information

- store between 20°-25°C (68°-77°F) in a dry place
- retain carton for complete product information
- store between 20°-25°C (68°-77°F) in a dry place
- retain carton for complete product information

NIGHTTIME

COLD

Drug Facts

This product contains acetaminophen. Severe liver damage may occur if you take

- with other drugs containing acetaminophen
- more than 12 caplets in 24 hours, which is the maximum daily amount
- 3 or more alcoholic drinks every day while using this product

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

DAYTIME

COLD & COUGH

Drug Facts

This product contains acetaminophen. Severe liver damage may occur if you take

- with other drugs containing acetaminophen
- more than 12 caplets in 24 hours, which is the maximum daily amount
- 3 or more alcoholic drinks every day while using this product

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Distributed by:

AAA Pharmaceutical, Inc.
681 Main Street
Lumberton, NJ 08048

Inactive ingredients

colloidal silicon dioxide, copovidone, croscarmellose sodium, FD&C blue #1, hypromellose, lactose anhydrous, magnesium stearate, methylparaben, povidone, pregelatinized starch, propylene glycol, propylparaben, sodium starch glycolate, stearic acid, titanium dioxide, triacetin
colloidal silicon dioxide, corn starch, croscarmellose sodium, crospovidone, FD&C yellow #6, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide