DRUG FACTS

Purpose Active ingredient(s)

Use(s)

temporarily relieves:

- · cough due to a cold or inhaled irritants
- · occasional minor irritation and pain due to sore throat or sore mouth

Warnings

Sore throat warnings:

- if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. These may be serious,
- do not use in children under 6 years of age unless directed by a doctor.

Do not use

Ask a doctor before use if you have

- · persistent chronic cough such as occurs with smoking, asthma, or emphysema
- cough accompanied by excessive phlegm (mucus)

Ask a doctor or pharmacist before use if you are

When using this product

Stop use and ask a doctor if

- · cough persists for more than 1 week, tends to recur, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.
- sore mouth does not improve in 7 days
- · irritation, pain, or redness persists or worsens

If pregnant or breast-feeding

Keep out of reach of children

Directions

- · adults and children 6 years and older: dissolve 2 drops (one at a time) slowly in the mouth. Repeat every 2 hours as needed or as directed by a doctor
- · children under 6 years: ask a doctor

Other information

- Phenylketonurics: contains phenylalanine 1 mg per drop
- · excessive use may have a laxative effect
- · store in a dry place
- · product may be useful for diabetics

on the advice of a doctor

Exchange information: 2 Drops = FREE Exchange 7 Drops = 1 Fruit

Dietary exchanges based on Exchange Lists for Diabetics

© 2008 American Diabetes Association, American Dietetic Association

Inactive ingredients aspartame, color (caramel), extracts of eucalyptus and a Ricola herb mixture (elder, horehound, hyssop, lemon balm, linden flowers, mallow, peppermint, sage, thyme, wild thyme), isomalt, natural flavor, sorbitol