DRUG FACTS

Active ingredient(s) ***Purpose***

Use(s)

- for temporary relief of occasional constipation and irregularity
- this product generally produces a bowel movement in 6 to 12 hours

Warnings

Do not use
if you cannot swallow without chewing.

Ask a doctor before use if you have
- stomach pain, nausea or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

Ask a doctor or pharmacist before use if you are

When using this product
- do not chew or crush tablet(s)
- it may cause stomach discomfort, faintness or cramps
- do not use within 1 hour after taking an antacid or milk

Stop use and ask a doctor if
- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding

Keep out of reach of children
In case of overdose, get medical help or contact a Poison Control Center right away.

Directions take with a glass of water

adults and children 12 yers of age and over	take 1 to 3 tablets in a single daily dose
children 6 to under 12 years of age	take 1 tablet in a single daily dose
children under 6 years of age	ask a doctor

Other information

- store at controlled room temperature 15º-30ºC (59º-86ºF)

Inactive ingredients

acacia, anhydrous calcium sulfate, anhydrous lactose, carnauba wax, colloidal silicon dioxide, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, gelatin, iron oxide, iron oxide black, iron oxide yellow (iron oxide ochre), magnesium stearate, microcrystalline cellulose, polyethylene glycol (PEG) 400, polyvinyl acetate phthalate, povidone, shellac, sodium starch glycolate, stearic acid, sugar, talc, and titanium dioxide