

DRUG FACTS

Active ingredient(s)

Purpose

Use(s)

- wash hands
- cut open tip of tube on scoremark
- use your fingertip or cotton applicator to apply a small pea-size amount of gel
- apply to affected area up to four times daily or as directed by a dentist or doctor
- for infants under 4 months of age, ask a dentist or doctor

Warnings

Allergy alert: do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other “caine” anesthetics

- Do not use more than directed for more than 7 days unless told to do so by a dentist or doctor.
- When using this product fever and nasal congestion are not symptoms of teething and may indicate the presence of infection. If these symptoms do not go away, advise your dentist or doctor.
- Stop use and ask a doctor if sore mouth symptoms do not get better in 7 days. swelling, rash or fever develops, irritation, pain or redness persists or worsens

Do not use

Ask a doctor before use if you have

Ask a doctor or pharmacist before use if you are

When using this product

Stop use and ask a doctor if

If pregnant or breast-feeding

Keep out of reach of children

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions • use up to 4 times daily, or as directed by a dentist or doctor • children under 12 years of age should be supervised in the use of this product • children under 2 years of age: consult a dentist or doctor

Other information

Inactive ingredients

FD&C Red 40, Flavor, Glycerin, Polyethylene Glycols, Purified Water, Sodium Saccharin, Sorbic Acid, Sorbitol