PEPTO-BISMOL- bismuth subsalicylate tablet, chewable
Lil' Drug Store Products, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Pepto-Bismol

Drug Facts

Active ingredient (in each tablet)

Bismuth subsalicylate 262 mg

Purposes

Upset stomach reliever and antidiarrheal

Uses

relieves
- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
  - heartburn
  - indigestion
  - nausea
  - gas
  - belching
  - fullness

Warnings

Reye's syndrome

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert

Contains salicylate. Do not take if you are
- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if you have
- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have
- fever

- mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for
- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if
- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
- chew or dissolve in mouth
- adults and children 12 years and over: 2 tablets every 1/2 to 1 hour as needed
- do not exceed 8 doses (16 tablets) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information
- **each tablet contains:** calcium 140 mg
- sodium less than 1 mg
- salicylate 102 mg
- very low sodium
- sugar free
- avoid excessive heat (over 104°F or 40°C)

Inactive ingredients

calcium carbonate, flavor, magnesium stearate, mannitol, povidone, red 27 aluminum lake, saccharin sodium, talc

Questions?

1-800-717-3786

Repackaged and distributed with permission of manufacturer by:
Lil' Drug Store Products
1201 Continental Place NE
Cedar Rapids, IA 52402

PRINCIPAL DISPLAY PANEL - 4 Tablet Pouch Carton

Pepto
BISMOL™

BISMUTH SUBSALICYLATE
UPSET STOMACH RELIEVER/ANTIDIARRHEAL

5 SYMPTOM RELIEF

Nausea • Heartburn • Indigestion
Upset Stomach • Diarrhea

4 CHEWABLE
TABLETS

Lil'
DrugStore®

Drug Facts (continued)

Other Information

- **each tablet contains:** calcium 140 mg
- sodium less than 1 mg
- salicylate 102 mg
- very low sodium
- sugar free
- avoid excessive heat (over 104°F or 40°C)

Inactive Ingredients

calcium carbonate, flavor, magnesium stearate, mannitol, povidone, red 27 aluminum lake, saccharin sodium, talc

Questions?

1-800-717-3786

Made in Mexico by Procter & Gamble
Manufactura S. de R.L. de C.V.
Dist. by Procter & Gamble, Cincinnati, OH 45202

www.pepto-bismol.com ©2011

Repackaged and distributed with permission of manufacturer by:
Lil' Drug Store Products
1201 Continental Place NE
Cedar Rapids, IA 52402
1-877-507-6516 M-F (8 AM-4:30 PM CST)
www.lildrugstore.com Rev 1-11

LOT

EXP

3 66715 97232 3

MULTI DOSE

Pepto BISMOL CHEWABLES

4 TABLETS

Pepto BISMOL™

BISMUTH SUBSALICYLATE
UPSET STOMACH RELIEVER/ANTIDIARRHEAL

5 SYMPTOM RELIEF
Nausea • Heartburn • Indigestion
Upset Stomach • Diarrhea

4 CHEWABLE TABLETS

Lil' DrugStore

Code 128
1476

TAMPER EVIDENT:
Do not use if the printed inner wrap is broken or missing.

Drug Facts

Active Ingredient (in each tablet) — Purposes

Bismuth subsalicylate 262 mg ..... Upset stomach reliever and antidiarrheal

Uses

relieves
- travelers' diarrhea • diarrhea
- upset stomach due to overindulgence in food and drink, including:
- heartburn • indigestion • nausea • gas • belching • fullness

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are
- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if you have
- an ulcer
- a bleeding problem
- bloody or black stool

LIFT PANEL FOR CONTINUED DRUG FACTS

Drug Facts (continued)

Ask a doctor before use if you have
- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for
- anticoagulation (thinning the blood)
- diabetes • gout • arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if
- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- chew or dissolve in mouth
- adults and children 12 years and over: 2 tablets every 1/2 to 1 hour as needed
- do not exceed 8 doses (16 tablets) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea



PEPTO-BISMOL

bismuth subsalicylate tablet, chewable

Product Information

Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:66715-9723(NDC:37000-477)
Route of Administration	ORAL	**DEA Schedule**	

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Bismuth subsalicylate (UNII: 62TEY51RR1) (Bismuth cation - UNII:ZS9CD1I8YE)	Bismuth subsalicylate	262 mg

Inactive Ingredients

Ingredient Name	Strength
calcium carbonate (UNII: H0G9379FGK)	
magnesium stearate (UNII: 70097M6I30)	
mannitol (UNII: 3OWL53L36A)	
povidones (UNII: FZ989GH94E)	
D&C red no. 27 (UNII: 2LRS185U6K)	
aluminum oxide (UNII: LMI26O6933)	
saccharin sodium (UNII: SB8ZUX40TY)	
talc (UNII: 7SEV7J4R1U)	

Product Characteristics

Color	PINK	**Score**	no score

Shape	ROUND	**Size**	16mm
Flavor		**Imprint Code**	Pepto;Bismol
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-9723-2	2 in 1 CARTON		
1		2 in 1 POUCH		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part335	09/16/2010	

Labeler - Lil' Drug Store Products, Inc. (093103646)

Establishment

Name	Address	ID/FEI	Business Operations
CCB Packaging Inc.		362374464	REPACK(66715-9723) , RELABEL(66715-9723)

Establishment

Name	Address	ID/FEI	Business Operations
The Proctor & Gamble Manufacturing Company		007130032	MANUFACTURE(66715-9723)

Revised: 1/2013 Lil' Drug Store Products, Inc.