DRUG FACTS

Active ingredient(s) Purpose

Use(s)

- relieves occasional constipation to help restore and maintain regularity
- this product generally produces bowel movement in 12 to 72 hours

Warnings

Choking: Taking this product without adequate fluid may cause it to swell and block your throat or esophagus and may cause choking. Do not take this product if you have difficulty in swallowing. If you experience chest pain, vomiting, or difficulty in swallowing or breathing after taking this product, seek immediate medical attention.

Do not use

Ask a doctor before use if you have

- abdominal pain, nausea, or vomiting
- a sudden change in bowel habits that persists over a period of 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug. Take this product 2 or more hours before or after other drugs. All laxatives may affect how other drugs work.

When using this product

- do not use for more than 7 days unless directed by a doctor
- do not take more than 8 caplets in a 24 hour period unless directed by a doctor

Stop use and ask a doctor if

rectal bleeding occurs or if you fail to have a bowel movement after use of this or any other laxative. These could be signs of a serious condition.

If pregnant or breast-feeding

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- take each dose of this product with at least 8 ounces (a full glass) of water or other fluid. Taking this product without enough liquid may cause choking. See choking warning.
- this product works naturally so continued use for one to three days is normally required to provide full benefit. Dosage may vary according to diet, exercise, previous laxative use or severity of constipation.

age	recommended dose	daily maximum
adults and children 12 years of age and over	2 caplets once a day	up to 4 times a day
children under 12 years	consult a physician	

$Other\ information$

- each caplet contains: calcium 140 mg and magnesium 10 mg
- store at 20-25 $^{\rm O}$ C (68-77 $^{\rm O}$ F)

Inactive ingredients

caramel, crospovidone, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium lauryl sulfate