MILK OF MAGNESIA- magnesium hydroxide liquid Meijer Distribution Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Meijer Distribution, Inc. Milk of Magnesia Drug Facts

Active ingredient (in each 15 mL TBSP)

Magnesium hydroxide 1200 mg

Purpose

Saline laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in ½ to 6 hours

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

Ask a doctor or pharmacist before use if you are

taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of
 a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

shake well before use

- do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided
- TBSP = tablespoonful

adults and children 12 years and older	2 to 4 TBSP.
children 6 to 11 years	1 to 2 TBSP.
children under 6 years	ask a doctor

Other information

- each tablespoon (15 mL) contains: calcium 20 mg, magnesium 595 mg, potassium 2 mg and sodium 2 mg
- does not meet USP requirements for preservative effectiveness
- store at 20-25°C (68-77°F)
- keep tightly closed

Inactive ingredients

anhydrous citric acid, D&C red #28, flavor, glycerin, purified water, sorbitol solution, sucrose, xanthan gum

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to Phillips'® active ingredient

Wild Cherry Flavor

Milk of Magnesia

Saline Laxative - Magnesium Hydroxide

Comfortable, Cramp Free Relief

Stimulant Free

ALCOHOL 0.4%

NDC 41250-949-40

meijer

Compare to Phillips'® active ingredient*

Wild Cherry Flavor

Milkof Magnesia

Saline Laxative • Magnesium Hydroxide

Comfortable, Cramp Free Relief

Stimulant Free

12 FL OZ (355 mL)

ALCOHOL 0.4%

: 94940 bE F7

Drug Facts

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING.

Active ingredient (in each 15 mL TBSP)

Purpose

Magnesium hydroxide 1200 mg......Saline laxative

- **Uses** relieves occasional constipation (irregularity)
- generally produces bowel movement in 1/2 to 6 hours

Warnings

Ask a doctor before use if you have

- kidney disease
 a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- shake well before use
- do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
 drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided
- TBSP = tablespoonful

adults and children 12 years and older	2 to 4 TBSP.	
children 6 to 11 years	1 to 2 TBSP.	
children under 6 years	ask a doctor	

Other information

■ each tablespoon (15 mL) contains: calcium 20 mg, magnesium 595 mg, potassium 2 mg and sodium 2 mg store at 20-25°C (68-77°F) keep tightly closed does not meet USP requirements for preservative effectiveness

Inactive ingredients anhydrous citric acid, D&C red #28, flavor, glycerin, purified water, sorbitol solution, sucrose, xanthan gum

(Questions or comments? 1-800-719-9260





MILK OF MAGNESIA

magnesium hydroxide liquid

Product Information			
Product Type	HUMAN OTC DRUG LABEL	Item Code (Source)	NDC:41250-949
Route of Administration	ORAL	DEA Schedule	

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
MAGNESIUM HYDRO XIDE (MAGNESIUM CATION and HYDRO XIDE ION)	MAGNESIUM HYDRO XIDE	1200 mg in 15 mL	

Inactive Ingredients			
Ingredient Name	Strength		
ANHYDRO US CITRIC ACID			
GLYCERIN			
WATER			
SORBITOL			
SUCROSE			
XANTHAN GUM			
D&C RED NO. 28			

Product Characteristics			
Color	PINK (Opaque Light pink)	Score	
Shape		Size	
Flavor	CHERRY (wild cherry)	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1 1	NDC:41250-949-40	355 mL in 1 BOTTLE		
2 1	NDC:41250-949-35	769 mL in 1 BOTTLE		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	10/18/2004	

Labeler - Meijer Distribution Inc (006959555)

Revised: 1/2013 Meijer Distribution Inc