TOPCARE MILK OF MAGNESIA- magnesium hydroxide liquid
Topco Associates LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Topco Associates LLC. Milk of Magnesia Drug Facts

Active ingredient (in each 15 mL TBSP)

Magnesium hydroxide 1200 mg

Purpose

Saline laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in ½ to 6 hours

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

Ask a doctor or pharmacist before use if you are

taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- shake well before use

- do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided
- TBSP = tablespoonful

adults and children 12 years and older	2 to 4 TBSP.
children 6 to 11 years	1 to 2 TBSP.
children under 6 years	ask a doctor

Other information

- each tablespoon (15 mL) contains: calcium 20 mg, magnesium 595 mg, potassium 2 mg and sodium 2 mg
- store at 20-25ºC (68-77ºF)
- keep tightly closed
- does not meet USP requirements for preservative effectiveness

Inactive ingredients

anhydrous citric acid, D&C red #28, flavor, glycerin, purified water, sorbitol solution, sucrose, xanthan gum

Questions or comments?

1-888-423-0139

Principal Display Panel

STIMULANT FREE

Milk of Magnesia

SALINE LAXATIVE

MAGNESIUM HYDROXIDE 1200 mg

Comfortable, cramp free relief

CHERRY FLAVOR

ALCOHOL 0.4%

COMPARE TO PHILLIPS'® WILD CHERRY active ingredient

TopCare®

NDC 36800-949-40

STIMULANT FREE

Milk of Magnesia

SALINE LAXATIVE

MAGNESIUM HYDROXIDE 1200 mg

- **Comfortable, cramp free relief**



CHERRY FLAVOR

ALCOHOL 0.4%

QUALITY GUARANTEED

COMPARE TO PHILLIPS'® WILD CHERRY active ingredient

12 FL OZ (355 mL)

: 94940 88 F6

Drug Facts

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING.

Active ingredient (in each 15 mL TBSP) — Purpose

Magnesium hydroxide 1200 mg.....Saline laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 1/2 to 6 hours

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- shake well before use
- do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided
- TBSP = tablespoonful

adults and children 12 years and older	2 to 4 TBSP.
children 6 to 11 years	1 to 2 TBSP.
children under 6 years	ask a doctor

Other information

- **each tablespoon (15 mL) contains:** calcium 20 mg, magnesium 595 mg, potassium 2 mg and sodium 2 mg
- store at 20-25°C (68-77°F)
- keep tightly closed
- does not meet USP requirements for preservative effectiveness

Inactive ingredients

anhydrous citric acid, D&C red #28, flavor, glycerin, purified water, sorbitol solution, sucrose, xanthan gum

Questions or comments? 1-888-423-0139

DISTRIBUTED BY TOPCO ASSOCIATES LLC

TOPCARE MILK OF MAGNESIA

magnesium hydroxide liquid

Product Information

Product Type	HUMAN OTC DRUG LABEL	**Item Code (Source)**	NDC:36800-949
Route of Administration	ORAL	**DEA Schedule**	

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MAGNESIUM HYDROXIDE (MAGNESIUM CATION and HYDROXIDE ION)	MAGNESIUM HYDROXIDE	1200 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID	
D&C RED NO. 28	
GLYCERIN	
WATER	
SORBITOL	
SUCROSE	
XANTHAN GUM	

Product Characteristics

Color	PINK (Opaque Light pink)	**Score**	
Shape		**Size**	
Flavor	CHERRY (wild cherry)	**Imprint Code**	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-949-40	355 mL in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	02/20/2006	

Labeler - Topco Associates LLC (006935977)

Revised: 1/2013 Topco Associates LLC