DRUG FACTS

Purpose Active ingredient(s)

Use(s)

- temporarily relieves minor aches and pains due to:
- · headache
- a cold
- arthritis
- muscular aches
- sinusitis
- toothache
- premenstrual & menstrual cramps

Warnings

Do not use

- · if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis. or kidney disease
- vou are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

taking

- any other drug containing an NSAID (prescription or nonprescription)
- · a blood thinning (anticoagulant) or steroid drug
- a prescription drug for diabetes, gout, or arthritis
- · any other drug, or are under a doctor's care for any serious condition

When using this product

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away
- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- · have bloody or black stools
- · have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- painful area is red or swollen
- pain gets worse or lasts for more than 10 days fever gets worse or lasts for more than 3 days
- any new symptoms appear

If pregnant or breast-feeding

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed
- · drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours
- · children under 12 years; ask a doctor

Other information

- store at controlled room temperature 20°-25°C (68°-77°F)
- excursions permitted to 15°C to 30°C (59°-86°F)

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness

Aspirin may cause a severe allergic reaction, which may include:

- · facial swelling
- asthma (wheezing)

This product contains acetaminophen. Severe liver damage may occur if you take

- · more than 8 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may $cause\ nervousness, irritability, sleeplessness, and, occasionally, rapid\ heartbeat.$

Inactive ingredients

hypromellose, maltodextrin, microcrystalline cellulose, polydextrose, polyvinylpyrrolidone, pregelatinized starch, silica, starch, stearic acid, talc, titanium dioxide, and triglycerides