DRUG FACTS

Purpose Active ingredient(s)

Use(s)

For relief of occasional constipation.

Warnings

May cause abdominal discomfort, faintness and cramps.

If pregnant or breast-feeding, ask a health professional before use.

Ask a doctor before using any laxative if you have abdominal pain, nausea or vomiting

- · a sudden change in bowel habits lasting more than 2 weeks
- · already used a laxative for more than 1 week

Swallow tablets whole. It is important that you do not chew or crush. Do not give to children under 6 years of age, or to persons who cannot swallow without chewing, unless directed by a doctor. Do not take within 1 hour after taking an antacid, milk or milk products.

Do not use

Ask a doctor before use if you have

Ask a doctor or pharmacist before use if you are

When using this product

Stop use and ask a doctor if

- have rectal bleeding
- · have no bowel movement after use

These symptoms may indicate a serious condition.

If pregnant or breast-feeding

Keep out of reach of children

 $\label{eq:case_equation} \mbox{In case of overdose, get medical help or contact a Poison Control Center right away.}$

Directions

Single Daily Dosage

adults and children 12 years and over 1 - 3 tablets

children 6 to under 12 years 1 tablet

children under 6 years ask a doctor

Other information

Inactive ingredients

acacia, anhydrous lactose, beeswax, calcium sulfate, carnauba wax, collodial silicon dioxide, corn starch, DandC yellow 10 lake, FDandC yellow 6 lake, gelatin, magnesium stearate, microcrystalline cellulose, pharmaceutical glaze, polyvinyl acetate phthalate, povidone, sodium starch glycolate, stearic acid, sucrose, talc, titanium dioxide