

HYDROCORTISONE- hydrocortisone cream
HYDROCORTISONE- hydrocortisone ointment
Perrigo New York Inc

HYDROCORTISONE CREAM USP, 2.5%
HYDROCORTISONE OINTMENT USP, 2.5%


Rx Only

DESCRIPTION

Each gram of Hydrocortisone Cream USP, 2.5% contains 25 mg of hydrocortisone in a cream base of cetyl alcohol, methylparaben, propylene glycol, propylparaben, purified water, sodium lauryl sulfate, and stearyl alcohol.

Each gram of Hydrocortisone Ointment USP, 2.5% contains 25 mg of hydrocortisone in ointment base of light mineral oil and white petrolatum.

Chemically, hydrocortisone is [Pregn-4-ene-3,20-dione,11,17,21-trihydroxy-, (11 β)-] with the molecular formula (C₂₁H₃₀O₅) and is represented by the following structural formula:



Its molecular weight is 362.47 and its CAS Registry Number is 50-23-7. The topical corticosteroids, including hydrocortisone, constitute a class of primarily synthetic steroids used as anti-inflammatory and antipruritic agents.

CLINICAL PHARMACOLOGY

Topical corticosteroids share anti-inflammatory, antipruritic, and vasoconstrictive actions. The mechanism of anti-inflammatory activity of the topical corticosteroids is unclear. Various laboratory methods, including vasoconstrictor assays, are used to compare and predict potencies and/or clinical efficacies of the topical corticosteroids. There is some evidence to suggest that a recognizable correlation exists between vasoconstrictor potency and therapeutic efficacy in man.

Pharmacokinetics -

The extent of percutaneous absorption of topical corticosteroids is determined by many factors including the vehicle, the integrity of the epidermal barrier, and the use of occlusive dressings. Topical corticosteroids can be absorbed from normal intact skin. Inflammation and/or other disease processes in the skin increase percutaneous absorption. Occlusive dressings substantially increase the percutaneous absorption of topical corticosteroids. Thus, occlusive dressings may be a valuable therapeutic adjunct for treatment of resistant dermatoses (see DOSAGE AND ADMINISTRATION).

Once absorbed through the skin, topical corticosteroids are handled through pharmacokinetic pathways

similar to systematically administered corticosteroids. Corticosteroids are bound to plasma proteins in varying degrees. Corticosteroids are metabolized primarily in the liver and are then excreted by the kidneys. Some of the topical corticosteroids and their metabolites are also excreted into the bile.

INDICATIONS AND USAGE

Topical corticosteroids are indicated for the relief of the inflammatory and pruritic manifestations of corticosteroid-responsive dermatosis.

CONTRAINDICATIONS

Topical corticosteroids are contraindicated in those patients with a history of hypersensitivity to any of the components of the preparation.

PRECAUTIONS

General -

Systemic absorption of topical corticosteroids has produced reversible hypothalamic-pituitary-adrenal (HPA) axis suppression, manifestations of Cushing's syndrome, hyperglycemia, and glucosuria in some patients. Conditions which augment systemic absorption include the application of the more potent steroids, use over large surface areas, prolonged use, and the addition of occlusive dressings.

Therefore, patients receiving a large dose of a potent topical steroid applied to a large surface area or under an occlusive dressing should be evaluated periodically for evidence of HPA axis suppression by using the urinary free cortisol and ACTH stimulation tests. If HPA axis suppression is noted, an attempt should be made to withdraw the drug, to reduce the frequency of application, or to substitute a less potent steroid. Recovery of HPA axis function is generally prompt and complete upon discontinuation of the drug. Infrequently, signs and symptoms of steroid withdrawal may occur, requiring supplemental systemic corticosteroids.

Children may absorb proportionally larger amounts of topical corticosteroids and thus be more susceptible to systemic toxicity (see PRECAUTIONS - Pediatric Use). If irritation develops, topical corticosteroids should be discontinued and appropriate therapy instituted.

In the presence of dermatological infections, the use of an appropriate antifungal or antibacterial agent should be instituted. If a favorable response does not occur promptly, the corticosteroid should be discontinued until the infection has been adequately controlled.

Information for the Patient -

Patients using topical corticosteroids should receive the following information and instructions:

1. This medication is to be used as directed by the physician. It is for external use only. Avoid contact with the eyes.
2. Patients should be advised not to use this medication for any disorder other than for which it was prescribed.
3. The treated skin area should not be bandaged or otherwise covered or wrapped as to be occlusive unless directed by the physician.
4. Patients should report any signs of local adverse reactions, especially under occlusive dressing.
5. Parents of pediatric patients should be advised not to use tight-fitting diapers or plastic pants on a child being treated in the diaper area, as these garments may constitute occlusive dressings.

Laboratory Tests

The following tests may be helpful in evaluating the HPA axis suppression:

Urinary free cortisol test

ACTH stimulation test

Carcinogenesis, Mutagenesis, Impairment of Fertility -

Long-term animal studies have not been performed to evaluate the carcinogenic potential or the effect on fertility of topical corticosteroids.

Studies to determine mutagenicity with prednisolone and hydrocortisone have revealed negative results.

Pregnancy:

Teratogenic Effects:

Pregnancy category C -

Corticosteroids are generally teratogenic in laboratory animals when administered systemically at relatively low dosage levels. The more potent corticosteroids have been shown to be teratogenic after dermal application in laboratory animals. There are no adequate and well-controlled studies in pregnant women on teratogenic effects from topically applied corticosteroids. Therefore, topical corticosteroids should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. Drugs of this class should not be used extensively on pregnant patients, in large amounts, or for prolonged periods of time.

Nursing Mothers -

It is not known whether topical administration of corticosteroids could result in sufficient systemic absorption to produce detectable quantities in breast milk. Systemically administered corticosteroids are secreted into breast milk in quantities not likely to have a deleterious effect on the infant. Nevertheless, caution should be exercised when topical corticosteroids are administered to a nursing woman.

Pediatric Use -

Pediatric patients may demonstrate greater susceptibility to topical corticosteroid-induced HPA axis suppression and Cushing's syndrome than mature patients because of a larger skin surface area to body weight ratio.

Hypothalamic-pituitary-adrenal (HPA) axis suppression, Cushing's syndrome, and intracranial hypertension have been reported in pediatric patients receiving topical corticosteroids. Manifestations of adrenal suppression in pediatric patients include linear growth retardation, delayed weight gain, low plasma cortisol levels, and absence of response to ACTH stimulation. Manifestations of intracranial hypertension include bulging fontanelles, headaches, and bilateral papilledema.

Administration of topical corticosteroids to pediatric patients should be limited to the least amount compatible with an effective therapeutic regimen. Chronic corticosteroid therapy may interfere with the growth and development of pediatric patients.

ADVERSE REACTIONS

The following local adverse reactions are reported infrequently with topical corticosteroids, but may occur more frequently with the use of occlusive dressings. These reactions are listed in an approximate decreasing order of occurrence: burning, itching, irritation, dryness, folliculitis, hypertrichosis, acneiform eruptions, hypopigmentation, perioral dermatitis, allergic contact dermatitis, maceration of the skin, secondary infection, skin atrophy, striae, and miliaria.

OVERDOSAGE

Topically applied corticosteroids can be absorbed in sufficient amounts to produce systemic effects (see PRECAUTIONS).

DOSAGE AND ADMINISTRATION

Topical corticosteroids are generally applied to the affected area as a thin film from two to four times daily depending on the severity of the condition. Occlusive dressings may be used for the management of psoriasis or recalcitrant conditions. If an infection develops, the use of occlusive dressings should be discontinued and appropriate antimicrobial therapy instituted.

HOW SUPPLIED

Hydrocortisone Cream USP, 2.5% is available as follows:

20 g tube (NDC 45802-**004**-02)

1 oz. (28 g) tube (NDC 45802-**004**-03)

Hydrocortisone Ointment USP, 2.5% is available as follows:

20 g tube (NDC 45802-**014**-02)

1 lb. jar (NDC 45802-**014**-05)

STORAGE

Store at 20-25°C (68-77°F) [see USP Controlled Room Temperature].

Keep out of the reach of children.

Manufactured By Perrigo

Bronx, NY 10457

Distributed By Perrigo

Allegan, MI 49010 · www.perrigo.com

Rev. 12/12

:1F300 RC J3

Principal Display Panel - Hydrocortisone Cream USP, 2.5% Carton

Rx Only


Hydrocortisone Cream USP, 2.5%



Principal Display Panel - Hydrocortisone Cream USP, 2.5% Tube

Rx Only


Hydrocortisone Cream USP, 2.5%



Principal Display Panel - Hydrocortisone Ointment USP, 2.5% Carton

Rx Only

Hydrocortisone Ointment USP, 2.5%



Principal Display Panel - Hydrocortisone Ointment USP, 2.5% Tube

Rx Only

Hydrocortisone Ointment USP, 2.5%

Rx Only

NDC 45802-014-02

Hydrocortisone Ointment USP, 2.5%

NET WT 20 g

Perrigo®

Each gram contains: 25 mg of hydrocortisone in an ointment base of light mineral oil, and white petrolatum.

Usual Dosage: Apply 2 to 4 times a day; see insert.

Keep out of reach of children.

Store at 20-25°C (68-77°F) [see USP Controlled Room Temperature]. For lot number and expiration date, see crimp of tube or carton.

Directions for puncturing tube seal: Remove cap. Turn cap upside down and place puncture tip onto tube seal; push down until seal is punctured. Screw cap back on to close.

Manufactured By
Perrigo
Bronx, NY 10457

Distributed By Perrigo
Allegan, MI 49010
www.perrigo.com

Rev. 11/12



N 3 45802-014-02 3

: 2C3A3 RC T3



HYDROCORTISONE

hydrocortisone cream

Product Information

Product Type	HUMAN PRESCRIPTION DRUG LABEL	Item Code (Source)	NDC:45802-004
Route of Administration	TOPICAL	DEA Schedule	

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HYDROCORTISONE (HYDROCORTISONE)	HYDROCORTISONE	25 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
WATER	
STEARYL ALCOHOL	
PROPYLENE GLYCOL	
CETYL ALCOHOL	
SODIUM LAURYL SULFATE	
METHYL PARABEN	
PROPYL PARABEN	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:45802-004-02	1 in 1 CARTON		
1		20 g in 1 TUBE		
2	NDC:45802-004-03	1 in 1 CARTON		
2		28 g in 1 TUBE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA085025	03/31/2006	

HYDROCORTISONE

hydrocortisone ointment

Product Information

Product Type	HUMAN PRESCRIPTION DRUG LABEL	Item Code (Source)	NDC:45802-014
Route of Administration	TOPICAL	DEA Schedule	

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HYDROCORTISONE (HYDROCORTISONE)	HYDROCORTISONE	25 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
PETROLATUM	
LIGHT MINERAL OIL	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:45802-014-02	1 in 1 CARTON		
1		20 g in 1 TUBE		
2	NDC:45802-014-05	454 g in 1 JAR		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA085027	06/13/2006	

