DRUG FACTS

Purpose Active ingredient(s)

Use(s)

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

if you have kidney disease, except under the advice and supervision of a doctor

- Ask a doctor before use if you have

 nausea, vomiting or abdominal pain
 - a sudden change in bowel habits that lasts over 2 weeks
 - irritable bowel syndrome

Ask a doctor or pharmacist before use if you are

taking a prescription drug

When using this product

you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

If pregnant or breast-feeding

Keep out of reach of children In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- do not take more than directed unless advised by your doctor
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
- fill to top of white section in cap which is marked to indicate the correct dose (17 g)
- stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
- use once a day
- use no more than 7 days children 16 years of age or under: ask a doctor

$Other\ information$

- store at 20#-25#C (68#-77#F)
- tamper-evident: do not use if printed foil seal under cap is missing, open or broken

Inactive ingredients