KETOROLAC TROMETHAMINE - ketorolac tromethamine injection KETOROLAC TROMETHAMINE - ketorolac tromethamine injection, solution

Fresenius Kabi USA, LLC

BOXED WARNING

Ketorolac tromethamine, a nonsteroidal anti-inflammatory drug (NSAID), is indicated for the shortterm (up to 5 days in adults) management of moderately severe acute pain that requires analgesia at the opioid level. It is NOT indicated for minor or chronic painful conditions. Ketorolac tromethamine is a potent NSAID analgesic, and its administration carries many risks. The resulting NSAID-related adverse events can be serious in certain patients for whom ketorolac tromethamine is indicated, especially when the drug is used inappropriately. Increasing the dose of ketorolac tromethamine beyond the label recommendations will not provide better efficacy but will result in increasing the risk of developing serious adverse events. Gastrointestinal Effects

- Ketorolac tromethamine can cause peptic ulcers, gastrointestinal bleeding and/or perforation. Therefore, ketorolac tromethamine is CONTRAINDICATED in patients with active peptic ulcer disease, in patients with recent gastrointestinal bleeding or perforation, and in patients with a history of peptic ulcer disease or gastrointestinal bleeding. Renal Effects
- Ketorolac tromethamine is CONTRAINDICATED in patients with advanced renal impairment and in patients at risk for renal failure due to volume depletion (see WARNINGS).
 Risk of Bleeding
- Ketorolac tromethamine inhibits platelet function and is, therefore, CONTRAINDICATED in patients with suspected or confirmed cerebrovascular bleeding, patients with hemorrhagic diathesis, incomplete hemostasis and those at high risk of bleeding (see WARNINGS and PRECAUTIONS).
- Ketorolac tromethamine is CONTRAINDICATED as prophylactic analgesic before any major surgery and is CONTRAINDICATED intraoperatively when hemostasis is critical because of the increased risk of bleeding.
 Hypersensitivity
- Hypersensitivity reactions, ranging from bronchospasm to anaphylactic shock, have occurred and appropriate
 counteractive measures must be available when administering the first dose of ketorolac tromethamine injection (see
 CONTRAINDICATIONS and WARNINGS). Ketorolac tromethamine is CONTRAINDICATED in patients with previously
 demonstrated hypersensitivity to ketorolac tromethamine or allergic manifestations to aspirin or other nonsteroidal antiinflammatory drugs (NSAIDs).

Intrathecal or Epidural Administration

- · Ketorolac tromethamine is CONTRAINDICATED for intrathecal or epidural administration due to its alcohol content. Labor, Delivery and Nursing
- The use of ketorolac tromethamine in labor and delivery is CONTRAINDICATED because it may adversely affect fetal circulation and inhibit uterine contractions.
- The use of ketorolac tromethamine is CONTRAINDICATED in nursing mothers because of the potential adverse effects of prostaglandin-inhibiting drugs on neonates.

Concomitant Use with NSAIDs

· Ketorolac tromethamine is CONTRAINDICATED in patients currently receiving ASA or NSAIDs because of the cumulative risk of inducing serious NSAID-related side effects.

DOSAGE AND ADMINISTRATION

Ketorolac Tromethamine Tablets

- Ketorolac tromethamine tablets are indicated only as continuation therapy to ketorolac tromethamine injection, and the combined duration of use of ketorolac tromethamine injection and ketorolac tromethamine tablets is not to exceed 5 days because of the increased risk of serious adverse events.
- The recommended total daily dose of ketorolac tromethamine tablets (maximum 40 mg) is significantly lower than for ketorolac tromethamine injection (maximum 120 mg) (see DOSAGE AND ADMINISTRATION).

 Special Populations
- Dosage should be adjusted for patients 65 years or older, for patients under 50 kg (110 lbs) of body weight (see DOSAGE AND ADMINISTRATION) and for patients with moderately elevated serum creatinine (see WARNINGS). Doses of ketorolac tromethamine injection are not to exceed 60 mg (total dose per day) in these patients. Ketorolac tromethamine injection is indicated as a single dose therapy in pediatric patients (see DOSAGE AND ADMINISTRATION); not to exceed 30 mg for IM administration and 15 mg for IV administration.

DESCRIPTION

Ketorolac tromethamine is a member of the pyrrolopyrrole group of nonsteroidal anti-inflammatory drugs (NSAIDs). The chemical name for ketorolac tromethamine is (\pm) -5-benzoyl-2,3-dihydro-1 \underline{H} pyrrolizine-1-carboxylic acid, compound with 2-amino-2-(hydroxymethyl)-1,3-propanediol (1:1), and the chemical structure is:

$C_{19}H_{24}N_2O_6$ M.W. 376.41

Ketorolac tromethamine is a racemic mixture of [-]S and [+]R ketorolac tromethamine. Ketorolac tromethamine may exist in three crystal forms. All forms are equally soluble in water. Ketorolac tromethamine has a pKa of 3.5 and an n-octanol/water partition coefficient of 0.26.

Ketorolac Tromethamine Injection, USP is available for intravenous (IV) or intramuscular (IM) administration as: 15 mg in 1 mL (1.5%) and 30 mg in 1 mL (3%) in sterile solution; 60 mg in 2 mL (3%) of ketorolac tromethamine in sterile solution is available for IM administration only. The solutions contain 0.1% citric acid, 10% (w/v) alcohol, USP, and 6.68 mg, 4.35 mg and 8.70 mg, respectively, of sodium chloride in sterile water. The pH is adjusted with sodium hydroxide or hydrochloric acid. The sterile solutions are clear and slightly yellow in color.

CLINICAL PHARMACOLOGY

Pharmacodynamics

Ketorolac tromethamine is a nonsteroidal anti-inflammatory drug (NSAID) that exhibits analgesic activity in animal models. Ketorolac tromethamine inhibits synthesis of prostaglandins and may be considered a peripherally acting analgesic. The biological activity of ketorolac tromethamine is associated with the S-form. Ketorolac tromethamine possesses no sedative or anxiolytic properties.

The peak analgesic effect of ketorolac tromethamine occurs within 2 to 3 hours and is not statistically significantly different over the recommended dosage range of ketorolac tromethamine. The greatest difference between large and small doses of ketorolac tromethamine by either route is in the duration of analgesia.

Pharmacokinetics

Ketorolac tromethamine is a racemic mixture of [-]S- and [+]R-enantiomeric forms, with the S-form having analgesic activity.

Comparison of IV, IM, and Oral Pharmacokinetics

The pharmacokinetics of ketorolac tromethamine, following IV, IM and oral doses of ketorolac tromethamine, are compared in Table 1. In adults, the extent of bioavailability following administration of the oral and IM forms of ketorolac tromethamine was equal to that following an IV bolus.

Linear Kinetics

In adults, following administration of single ORAL, IM or IV doses of ketorolac tromethamine in the recommended dosage ranges, the clearance of the racemate does not change. This implies that the pharmacokinetics of ketorolac tromethamine in adults, following single or multiple IM, IV or recommended oral doses of ketorolac tromethamine, are linear. At the higher recommended doses, there is a proportional increase in the concentrations of free and bound racemate.

Distribution

The mean apparent volume $(V\beta)$ of ketorolac tromethamine following complete distribution was approximately 13 liters. This parameter was determined from single-dose data. The ketorolac tromethamine racemate has been shown to be highly protein bound (99%). Nevertheless, even plasma concentrations as high as 10 mcg/mL will only occupy approximately 5% of the albumin binding sites. Thus, the unbound fraction for each enantiomer will be constant over the therapeutic range. A decrease in serum albumin, however, will result in increased free drug concentrations.

Ketorolac tromethamine is excreted in human milk (see PRECAUTIONS: Lactation and Nursing).

Metabolism

Ketorolac tromethamine is largely metabolized in the liver. The metabolic products are hydroxylated and conjugated forms of the parent drug. The products of metabolism, and some unchanged drug, are excreted in the urine.

Excretion

The principal route of elimination of ketorolac and its metabolites is renal. About 92% of a given dose is found in the urine, approximately 40% as metabolites and 60% as unchanged ketorolac. Approximately 6% of a dose is excreted in the feces. A single-dose study with 10 mg ketorolac tromethamine (n=9) demonstrated that the S-enantiomer is cleared approximately two times faster than the R-enantiomer and that the clearance was independent of the route of administration. This means that the ratio of S/R plasma concentrations decreases with time after each dose. There is little or no inversion of the R- to S-form in humans. The clearance of the racemate in normal subjects, elderly individuals and in hepatically and renally impaired patients is outlined in Table 2 (see

CLINICAL PHARMACOLOGY: Kinetics in Special Populations).

The half-life of the ketorolac tromethamine S-enantiomer was approximately 2.5 hours (SD \pm 0.4) compared with 5 hours (SD \pm 1.7) for the R-enantiomer. In other studies, the half-life for the racemate has been reported to lie within the range of 5 to 6 hours.

Accumulation

Ketorolac tromethamine administered as an IV bolus every 6 hours for 5 days to healthy subjects (n=13), showed no significant difference in C_{max} on Day 1 and Day 5. Trough levels averaged 0.29 mcg/mL (SD \pm 0.13) on Day 1 and 0.55 mcg/mL (SD \pm 0.23) on Day 6. Steady state was approached after the fourth dose.

Accumulation of ketorolac tromethamine has not been studied in special populations (elderly, pediatric, renal failure, or hepatic disease patients).

Kinetics in Special Populations

Geriatric Patients

Based on single-dose data only, the half-life of the ketorolac tromethamine racemate increased from 5 to 7 hours in the elderly (65 to 78 years) compared with young healthy volunteers (24 to 35 years) (see Table 2). There was little difference in the C_{max} for the two groups (elderly, 2.52 mcg/mL \pm 0.77; young, 2.99 mcg/mL \pm 1.03) (see **PRECAUTIONS**: *Geriatric Use*).

Pediatric Patients

Following a single intravenous bolus dose of 0.5 mg/kg in 10 children 4 to 8 years old, the half-life was 6 hours (range: 3.5 to 10 h), the average clearance was 0.042 L/hr/kg and the Vd was 0.26 L/kg (range: 0.19 to 0.44 L/kg). In a second study, following a single intravenous dose of 0.6 mg/kg in 24 children 3 to 18 years old, C_{max} was 4.3 ± 1.7 mcg/mL, T_{max} was 10.25 ± 1.15 minutes, half-life was 3.8 ± 2.6 hours, CI was 0.0678 L/hr/kg and Vd was 0.25 L/kg. The volume of distribution and clearance of ketorolac in pediatric patients was twice that observed in adult subjects (see Tables 1 and 2). There are no pharmacokinetic data available for ketorolac tromethamine administration by the IM route in pediatric patients.

Renal Insufficiency

Based on single-dose data only, the mean half-life of ketorolac tromethamine in renally impaired patients is between 6 and 19 hours and is dependent on the extent of the impairment. There is poor correlation between creatinine clearance and total ketorolac tromethamine clearance in the elderly and populations with renal impairment (r=0.5).

In patients with renal disease, the AUC∞ of each enantiomer increased by approximately 100% compared with healthy volunteers.

The volume of distribution doubles for the S-enantiomer and increases by 1/5th for the Renantiomer. The increase in volume of distribution of ketorolac tromethamine implies an increase in unbound fraction.

Table 1 Table of Approximate Average Pharmacokinetic Parameters (Mean \pm SD) Following Oral, Intramuscular and Intravenous Doses of Ketorolac Tromethamine

	Oral [*] Intramuscular [†]			Intravenous Bolus [‡]		
Pharmacokinetic Parameters (units)	10 mg	15 mg	30 mg	60 mg	15 mg	30 mg
Bioavailability (extent)		L	100	0%	<u> </u>	<u>I</u>
T _{max} ¹ (min)	44 ±34	33±21 [#]	44±29	33±21 [#]	1.1±0.7 [#]	2.9±1.8
C _{max} ² (mcg/mL) [single-dose]	0.87±0.22	1.14±0.32 [#]	2.42±0.68	4.55±1.27 [#]	2.47±0.51 [#]	4.65±0.96
C _{max} (mcg/mL) [steady state qid]	1.05±0.26 [#]	1.56±0.44 [#]	3.11±0.87 [#]	N/A ^{II}	3.09±1.17 [#]	6.85±2.61
C _{min} ³ (mcg/mL) [steady state qid]	0.29±0.07 [#]	0.47±0.13 [#]	0.93±0.26 [#]	N/A	0.61±0.21 [#]	1.04±0.35
C _{avg} ⁴ (mcg/mL) [steady state qid]	0.59±0.20 [#]	0.94±0.29 [#]	1.88±0.59 [#]	N/A	1.09±0.30 [#]	2.17±0.59
Vβ ⁵ (L/kg)			0.175±0.039		0.210-	±0.044

%Dose metabolized =<50 %Dose excreted in feces= 6 %Plasma protein binding= 99

Table 2

The Influence of Age, Liver and Kidney Function, on the Clearance and Terminal Half-life of Ketorolac Tromethamine $(IM^1 and\ ORAL^2)$ In Adult Populations

	Total Clearan	ce [in L/h/kg] ³	Terminal Half-life [in hours]	
Type of Subjects	IM	ORAL	IM	ORAL
	Mean (range)	Mean (range)	Mean (range)	Mean (range)
Normal Subjects IM (n=54) mean age= 32, range= 18-60 Oral (n=77) mean age= 32, range = 20-60	0.023	0.025	5.3	5.3
	(0.010-0.046)	(0.013-0.050)	(3.5-9.2)	(2.4-9)
Healthy Elderly Subjects IM (n=13),Oral (n=12) mean age= 72, range= 65-78	0.019 (0.013-0.034)	0.024 (0.018-0.034)	7 (4.7-8.6)	6.1 (4.3-7.6)
Patients with Hepatic Dysfunction IM and Oral (n=7) mean age= 51, range= 43-64	0.029	0.033	5.4	4.5
	(0.013-0.066)	(0.019-0.051)	(2.2-6.9)	(1.6-7.6)
Patients with Renal Impairment IM (n=25), Oral (n=9) serum creatinine= 1.9-5.0 mg/dL, mean age (IM)=54, range= 35-71 mean age (Oral)=57, range 39-70	0.015	0.016	10.3	10.8
	(0.005-0.043)	(0.007-0.052)	(5.9-19.2)	(3.4-18.9)
Renal Dialysis Patients IM and Oral (n=9) mean age = 40, range = 27-63	0.016 (0.003-0.036)		13.6 (8.0-39.1)	

^{*} Derived from PO pharmacokinetic studies in 77 normal fasted volunteers

[†] Derived from IM pharmacokinetic studies in 54 normal volunteers

[‡] Derived from IV pharmacokinetic studies in 24 normal volunteers

 $^{^{\#}}$ Mean value was simulated from observed plasma concentration data and standard deviation was simulated from percent coefficient of variation for observed C_{max} and T_{max} data

¹¹ Not applicable because 60 mg is only recommended as a single dose

¹ Time - to - peak plasma concentration

² Peak plasma concentration

³ Trough plasma concentration

⁴ Average plasma concentration

⁵ Volume of distribution

The AUC_{∞} - ratio of the ketorolac tromethamine enantiomers in healthy subjects and patients remained similar, indicating there was no selective excretion of either enantiomer in patients compared to healthy subjects (see WARNINGS: *Renal Effects*; see Table 2).

Hepatic Insufficiency

There was no significant difference in estimates of halflife, AUC_{∞} , and C_{max} in 7 patients with liver disease compared to healthy volunteers (see PRECAUTIONS: *General*, Hepatic Effects; see Table 2).

Race

Pharmacokinetic differences due to race have not been identified.

IV Administration

In normal adult subjects (n=37), the total clearance of 30 mg IV-administered ketorolac tromethamine was 0.030 (0.017-0.051) L/h/kg. The terminal half-life was 5.6 (4.0-7.9) hours (see **Kinetics in Special Populations** for use of ketorolac tromethamine injection in pediatric patients).

Clinical Studies

Adult Patients

The analgesic efficacy of intramuscularly, intravenously, and orally administered ketorolac tromethamine was investigated in two postoperative pain models: general surgery (orthopedic, gynecologic, and abdominal) and oral surgery (removal of impacted third molars). The studies were double-blind, single- and multiple-dose, parallel trial designs in patients with moderate to severe pain at baseline. Ketorolac tromethamine injection was compared as follows: IM to meperidine or morphine administered intramuscularly and IV to morphine administered either directly IV or through a PCA (Patient-Controlled Analgesia) pump.

Short-Term Use (up to 5 days) Studies

In adults, the comparisons of intramuscular administration during the first hour, the onset of analgesic action was similar for ketorolac tromethamine and the narcotics, but the duration of analgesia was longer with ketorolac tromethamine than with the opioid comparators meperidine or morphine.

In a multidose, postoperative (general surgery) double-blind trial of ketorolac tromethamine 30 mg IM versus morphine 6 and 12 mg IM, each drug given on an "as needed" basis for up to 5 days, the overall analysesic effect of ketorolac tromethamine 30 mg IM was between that of morphine 6 and 12 mg. The majority of patients treated with either ketorolac tromethamine or morphine were dosed for up to 3 days; a small percentage of patients received 5 days of dosing.

In clinical settings where perioperative morphine was allowed, ketorolac tromethamine 30 mg IV, given once or twice as needed, provided analgesia comparable to morphine 4 mg IV once or twice as needed.

There was relatively limited experience with 5 consecutive days of ketorolac tromethamine IV use in controlled clinical trials, as most patients were given the drug for 3 days or less. The adverse events seen with IV-administered ketorolac tromethamine were similar to those observed with IM-administered ketorolac tromethamine, as would be expected based on the similar pharmacokinetics and bioequivalence (AUC, clearance, plasma half-life) of IV and IM routes of ketorolac tromethamine administration.

Pediatric Patients

The analgesic efficacy of single doses of ketorolac tromethamine injection has been demonstrated by showing a decrease in the need for supplemental narcotic in pediatric patients receiving ketorolac as compared to placebo. See discussion of these results under **Clinical Studies with Concomitant Use of Opioids** below.

Clinical Studies with Concomitant Use of Opioids

Adult Patients

Clinical studies in postoperative pain management have demonstrated that ketorolac tromethamine injection, when used in combination with opioids, significantly reduced opioid consumption. This combination may be useful in the subpopulation of patients especially prone to opioid-related complications. Ketorolac tromethamine and narcotics should not be administered in the same syringe.

In a postoperative study, where all patients received morphine by a PCA device, patients treated with ketorolac tromethamine IV as fixed intermittent boluses (eg, 30 mg initial dose followed by 15 mg q3h), required significantly less morphine (26%) than the placebo group. Analgesia was significantly superior, at various postdosing pain assessment times, in the patients receiving ketorolac tromethamine IV plus PCA morphine as compared to patients receiving PCA-administered morphine alone.

Pediatric Patients

Ketorolac tromethamine intramuscular injection reduced the need for supplemental opioid (fentanyl) when a 1 mg/kg dose was administered immediately following tonsillectomy compared to saline controls (see WARNINGS: *Hemorrhage*). In another study, when a single bolus dose of 0.9 mg/kg of ketorolac tromethamine intravenous injection was given to pediatric patients ages 5 to 12 years, compared to saline, a reduction in supplemental opioid was needed following various surgical procedures. In a third study less supplemental morphine was needed in pediatric patients ages 8 to 16 years, who received a 0.8 mg/kg IV injection of ketorolac in conjunction with morphine following orthopedic surgical procedures, compared to morphine alone. In a study in pediatric patients

¹Estimated from 30 mg single IM doses of ketorolac tromethamine

²Estimated from 10 mg single oral doses of ketorolac tromethamine

³Liters/hour/kilogram

ages 3 to 12 years, ketorolac tromethamine intravenous injection demonstrated a slower onset of analgesia, but a longer duration of action compared to morphine. There is limited data available to support the use of multiple doses of ketorolac tromethamine injection in pediatric patients.

Postmarketing Surveillance Study

A large postmarketing observational, nonrandomized study, involving approximately 10,000 patients receiving ketorolac tromethamine, demonstrated that the risk of clinically serious gastrointestinal (GI) bleeding was dose-dependent (see Tables 3A and 3B). This was particularly true in elderly patients who received an average daily dose greater than 60 mg/day of ketorolac tromethamine (Table 3A).

Table 3

Incidence of Clinically Serious GI Bleeding as Related to Age, Total Daily Dose, and History of GI Perforation, Ulcer, Bleeding (PUB) after up to 5 Days of Treatment with Ketorolac Tromethamine Injection

A. Adult Patients without History of PUB

	Total Daily Dose of Ketorolac Tromethamine Injection			
Age of Patients	≤ 60 mg	> 60 to 90 mg	> 90 to 120 mg	> 120 mg
< 65 years of age	0.4%	0.4%	0.9%	4.6%
≥ 65 years of age	1.2%	2.8%	2.2%	7.7%

B. Adult Patients with History of PUB

	Total Daily Dose of Ketorolac Tromethamine Injection			
Age of Patients	≤ 60 mg	> 60 to 90 mg	> 90 to 120 mg	> 120 mg
< 65 years of age	2.1%	4.6%	7.8%	15.4%
≥ 65 years of age	4.7%	3.7%	2.8%	25%

INDICATIONS AND USAGE

Adult Patients

Ketorolac tromethamine injection is indicated for the short-term (≤ 5 days) management of moderately severe acute pain that requires analgesia at the opioid level, usually in a postoperative setting. Therapy should always be initiated with ketorolac tromethamine injection, and ketorolac tromethamine tablets are to be used only as continuation treatment, if necessary. Combined use of ketorolac tromethamine injection and tablets is not to exceed 5 days of use because of the potential of increasing the frequency and the severity of adverse reactions associated with the recommended doses (see WARNINGS, PRECAUTIONS, DOSAGE AND ADMINISTRATION, and ADVERSE REACTIONS). Patients should be switched to alternative analgesics as soon as possible, but ketorolac tromethamine therapy is not to exceed 5 days.

Pediatric Patients

The safety and effectiveness of single doses of ketorolac tromethamine injection have been established in pediatric patients between the ages of 2 and 16 years. Ketorolac, as a single injectable dose, has been shown to be effective in the management of moderately severe acute pain that requires analgesia at the opioid level, usually in the postoperative setting. There is limited data available to support the use of multiple doses of ketorolac in pediatric patients. Safety and effectiveness have not been established in pediatric patients below the age of 2 years. Use of ketorolac in pediatric patients is supported by evidence from adequate and well-controlled studies of ketorolac in adults with additional pharmacokinetic, efficacy and safety data on its use in pediatric patients available in the published literature (see CLINICAL PHARMACOLOGY: *Clinical Studies*, WARNINGS and **PRECAUTIONS**).

Ketorolac tromethamine injection has been used concomitantly with morphine and meperidine and has shown an opioid-sparing effect. For breakthrough pain, it is recommended to supplement the lower end of the ketorolac tromethamine injection dosage range with low doses of narcotics prn, unless otherwise contraindicated. Ketorolac tromethamine injection and narcotics should

not be administered in the same syringe (see **DOSAGE AND ADMINISTRATION**: *Pharmaceutical Information for Ketorolac Tromethamine Injection*).

CONTRAINDICATIONS

(see also Boxed WARNING)

- · Ketorolac tromethamine is CONTRAINDICATED in patients with active peptic ulcer disease, in patients with recent gastrointestinal bleeding or perforation and in patients with a history of peptic ulcer disease or gastrointestinal bleeding.
- · Ketorolac tromethamine is CONTRAINDICATED in patients with advanced renal impairment or in patients at risk for renal failure due to volume depletion (see **WARNINGS** for correction of volume depletion).
- · Ketorolac tromethamine is CONTRAINDICATED in labor and delivery because, through its prostaglandin synthesis inhibitory effect, it may adversely affect fetal circulation and inhibit uterine contractions, thus increasing the risk of uterine hemorrhage.
- The use of ketorolac tromethamine is CONTRAINDICATED in nursing mothers because of the potential adverse effects of prostaglandin-inhibiting drugs on neonates.
- · Ketorolac tromethamine is CONTRAINDICATED in patients with previously demonstrated hypersensitivity to ketorolac tromethamine, allergic manifestations to aspirin or other nonsteroidal anti-inflammatory drugs (NSAIDs).
- · Ketorolac tromethamine is CONTRAINDICATED as prophylactic analgesic before any major surgery and is CONTRAINDICATED intraoperatively when hemostasis is critical because of the increased risk of bleeding.
- · Ketorolac tromethamine inhibits platelet function and is, therefore, CONTRAINDICATED in patients with suspected or confirmed cerebrovascular bleeding, hemorrhagic diathesis, incomplete hemostasis and those at high risk of bleeding (see **WARNINGS** and **PRECAUTIONS**).
- · Ketorolac tromethamine is CONTRAINDICATED in patients currently receiving ASA or NSAIDs because of the cumulative risks of inducing serious NSAID-related adverse events.
- · Ketorolac tromethamine injection is CONTRAINDICATED for neuraxial (epidural or intrathecal) administration due to its alcohol content.
- The concomitant use of ketorolac tromethamine and probenecid is CONTRAINDICATED.

WARNINGS

(see also Boxed WARNING)

The combined use of ketorolac tromethamine injection and ketorolac tablets is not to exceed 5 days in adults. Only single doses of ketorolac tromethamine injection are recommended for use in pediatric patients. The most serious risks associated with ketorolac tromethamine are:

Gastrointestinal (GI) Effects - Risk of GI Ulcerations, Bleeding, and Perforation

Ketorolac tromethamine is CONTRAINDICATED in patientswith previously documented peptic ulcers and/or GI bleeding. Serious gastrointestinal toxicity, such as bleeding, ulceration and perforation, can occur at any time, with or without warning symptoms, in patients treated with ketorolac tromethamine. Studies to date with NSAIDs have not identified any subset of patients not at risk of developing peptic ulceration and bleeding. Elderly or debilitated patients seem to tolerate ulceration or bleeding less well than other individuals, and most spontaneous reports of fatal GI events are in this population. Postmarketing experience with parenterally administered ketorolac tromethamine suggests that there may be a greater risk of gastrointestinal ulcerations, bleeding and perforation in the elderly.

The incidence and severity of gastrointestinal complications increases with increasing dose of, and duration of treatment with, ketorolac tromethamine. In a non-randomized, in-hospital postmarketing surveillance study comparing parenteral ketorolac tromethamine to parenteral opioids, higher rates of clinically serious GI bleeding were seen in adult patients < 65 years of age who received an average total daily dose of more than 90 mg of ketorolac tromethamine injection per day (see **CLINICAL PHARMACOLOGY: Postmarketing Surveillance Study**).

The same study showed that elderly (\geq 65 years of age) and debilitated patients are more susceptible to gastrointestinal complications. A history of peptic ulcer disease was revealed as another risk factor that increases the possibility of developing serious gastrointestinal complications during ketorolac tromethamine therapy (see Tables 3A and 3B).

· Hemorrhage

Because prostaglandins play an important role in hemostasis and NSAIDs affect platelet aggregation as well, use of ketorolac tromethamine in patients who have coagulation disorders should be undertaken very cautiously, and those patients should be carefully monitored. Patients on therapeutic doses of anticoagulants (eg, heparin or dicumarol derivatives) have an increased risk of bleeding complications if given ketorolac tromethamine concurrently; therefore, physicians should administer such concomitant therapy only extremely cautiously. The concurrent use of ketorolac tromethamine and prophylactic low-dose heparin (2500 to 5000 units q12h), warfarin and dextrans have not been studied extensively, but may also be associated with an increased risk of bleeding. Until data from such studies are available, physicians should carefully weigh the benefits against the risks and use such concomitant therapy in these patients only extremely cautiously. In patients who receive anticoagulants for any reason, there is an increased risk of intramuscular hematoma formation from administered ketorolac tromethamine IM (see PRECAUTIONS: *Drug Interactions*). Patients receiving therapy that affects hemostasis should be monitored closely.

In postmarketing experience, postoperative hematomas and other signs of wound bleeding have been reported in association with the perioperative use of ketorolac tromethamine injection. Therefore, perioperative use of ketorolac tromethamine should be avoided and postoperative use be undertaken with caution when hemostasis is critical (see **WARNINGS** and **PRECAUTIONS**).

Pediatrics and Tonsillectomy

Physicians should consider the increased risk of bleeding before deciding to administer ketorolac in patients following tonsillectomy. Ketorolac tromethamine injection is not recommended for use in pediatric patients below the age of 2 years. In retrospective analysis of patients having undergone tonsillectomy with or without adenoidectomy, the risk of bleeding was 10.1% in patients administered ketorolac tromethamine injection compared to 2.2% in those receiving opioids. The postoperative hemorrhage rate in patients 12 years and younger was 6.5% and 3.3% with and without ketorolac, respectively. In a prospective study of ketorolac in pediatric patients (ages 3 to 9 years) undergoing tonsillectomy with or without adenoidectomy, the overall incidence of bleeding was similar between the patients receiving ketorolac and morphine (16.3% versus 17%, respectively). However, during the first 24 hours after surgery, a higher incidence of bleeding was observed in the ketorolac tromethamine injection group (14.3%) versus the morphine group (4.2%).

· Anaphylactoid Reactions

Anaphylactoid reactions may occur in patients without a known previous exposure or hypersensitivity to aspirin, ketorolac tromethamine or other NSAIDs, or in individuals with a history of angioedema, bronchospastic reactivity (eg, asthma) and nasal polyps. Anaphylactoid reactions, like anaphylaxis, may have a fatal outcome.

· Impaired Renal Function

Ketorolac tromethamine should be used with caution in patients with impaired renal function or a history of kidney disease because it is a potent inhibitor of prostaglandin synthesis. Renal toxicity with ketorolac tromethamine has been seen in patients with conditions leading to a reduction in blood volume and/or renal blood flow where renal prostaglandins have a supportive role in the maintenance of renal perfusion. In these patients administration of ketorolac tromethamine may cause a dose-dependent reduction in renal prostaglandin formation and may precipitate acute renal failure. Patients at greatest risk of this reaction are those with impaired renal function, dehydration, heart failure, liver dysfunction, those taking diuretics and the elderly. Discontinuation of ketorolac tromethamine therapy is usually followed by recovery to the pretreatment state.

· Renal Effects

Ketorolac tromethamine and its metabolites are eliminated primarily by the kidneys, which, in patients with reduced creatinine clearance, will result in diminished clearance of the drug (see **CLINICAL PHARMACOLOGY**). Therefore, ketorolac tromethamine should be used with caution in patients with impaired renal function (see **DOSAGE AND ADMINISTRATION**) and such patients should be followed closely. With the use of ketorolac tromethamine, there have been reports of acute renal failure, interstitial nephritis and nephrotic syndrome.

Because patients with underlying renal insufficiency are at increased risk of developing acute renal failure, the risks and benefits should be assessed prior to giving ketorolac tromethamine to these patients. Hence, in patients with moderately elevated serum creatinine, it is recommended that the daily dose of ketorolac tromethamine injection be reduced by half, not to exceed 60 mg/day. KETOROLAC TROMETHAMINE IS CONTRAINDICATED IN PATIENTS WITH SERUM CREATININE CONCENTRATIONS INDICATING ADVANCED RENAL IMPAIRMENT (see CONTRAINDICATIONS).

Hypovolemia should be corrected *before* treatment with ketorolac tromethamine is initiated.

• Fluid Retention and Edema

Fluid retention, edema, retention of NaCl, oliguria, elevations of serum urea nitrogen and creatinine have been reported in clinical trials with ketorolac tromethamine. Therefore, ketorolac tromethamine should be used only very cautiously in patients with cardiac decompensation, hypertension or similar conditions.

• Pregnancy

In late pregnancy, as with other NSAIDS, ketorolac tromethamine should be avoided because it may cause premature closure of the ductus arteriosus.

PRECAUTIONS

General

Hepatic Effects

Ketorolac tromethamine should be used with caution in patients with impaired hepatic function or a history of liver disease. Treatment with ketorolac tromethamine may cause elevations of liver enzymes, and, in patients with pre-existing liver dysfunction, it may lead to the development of a more severe hepatic reaction. The administration of ketorolac tromethamine should be discontinued in patients in whom an abnormal liver test has occurred as a result of ketorolac tromethamine therapy.

Hematologic Effects

Ketorolac tromethamine inhibits platelet aggregation and may prolong bleeding time; therefore, it is contraindicated as a preoperative medication, and caution should be used when hemostasis is critical. Unlike aspirin, the inhibition of platelet function by ketorolac tromethamine disappears within 24 to 48 hours after the drug is discontinued. Ketorolac tromethamine does not appear to affect platelet count, prothrombin time (PT) or partial thromboplastin time (PTT). In controlled clinical studies, where ketorolac

tromethamine was administered intramuscularly or intravenously postoperatively, the incidence of clinically significant postoperative bleeding was 0.4% for ketorolac tromethamine compared to 0.2% in the control groups receiving narcotic analgesics.

Information for Patients

Ketorolac tromethamine is a potent NSAID and may cause serious side effects such as gastrointestinal bleeding or kidney failure, which may result in hospitalization and even fatal outcome.

Physicians, when prescribing ketorolac tromethamine, should inform their patients or their guardians of the potential risks of ketorolac tromethamine treatment (see **Boxed WARNING**, **WARNINGS**, **PRECAUTIONS** and **ADVERSE REACTIONS** sections). *Advise patients not to give ketorolac tromethamine tablets to other family members and to discard any unused drug.*

Remember that the total duration of ketorolac therapy is not to exceed 5 days in adults or a single dose in pediatric patients ages 2 to 16 years.

Drug Interactions

Ketorolac is highly bound to human plasma protein (mean 99.2%).

Warfarin, Digoxin, Salicylate and Heparin: The *in vitro* binding of *warfarin* to plasma proteins is only slightly reduced by ketorolac tromethamine (99.5% control vs 99.3%) when ketorolac plasma concentrations reach 5 to 10 mcg/mL. Ketorolac does not alter *digoxin* protein binding. *In vitro* studies indicate that, at therapeutic concentrations of *salicylate* (300 mcg/mL), the binding of ketorolac was reduced from approximately 99.2% to 97.5%, representing a potential two-fold increase in unbound ketorolac plasma levels. Therapeutic concentrations of *digoxin*, *warfarin*, *ibuprofen*, *naproxen*, *piroxicam*, *acetaminophen*, *phenytoin*, and *tolbutamide* did not alter ketorolac tromethamine protein binding.

In a study involving 12 adult volunteers, ketorolac tromethamine tablets were coadministered with a single dose of 25 mg *warfarin*, causing no significant changes in pharmacokinetics or pharmacodynamics of warfarin. In another study, ketorolac tromethamine injection was given with two doses of 5000 U of *heparin* to 11 healthy volunteers, resulting in a mean template bleeding time of 6.4 minutes (3.2 to 11.4 min) compared to a mean of 6.0 minutes (3.4 to 7.5 min) for heparin alone and 5.1 minutes (3.5 to 8.5 min) for placebo. Although these results do not indicate a significant interaction between ketorolac tromethamine and warfarin or heparin, the administration of ketorolac tromethamine to patients taking anticoagulants should be done extremely cautiously, and patients should be closely monitored (see **WARNINGS** and **PRECAUTIONS**).

Furosemide: Ketorolac tromethamine injection reduced the diuretic response to *furosemide* in normovolemic healthy subjects by approximately 20% (mean sodium and urinary output decreased 17%).

Probenecid: Concomitant administration of ketorolac tromethamine tablets and *probenecid* resulted in decreased clearance of ketorolac and significant increases in ketorolac plasma levels (total AUC increased approximately threefold from 5.4 to 17.8 mcg/h/mL) and terminal half-life increased approximately twofold from 6.6 to 15.1 hours. Therefore, concomitant use of ketorolac tromethamine and probenecid is contraindicated.

Lithium: Inhibition of renal *lithium* clearance, leading to an increase in plasma lithium concentration, has been reported with some prostaglandin synthesisinhibiting drugs. The effect of ketorolac tromethamine on plasma lithium has not been studied, but cases of increased lithium plasma levels during ketorolac tromethamine therapy have been reported.

Methotrexate: Concomitant administration of *methotrexate* and some NSAIDs has been reported to reduce the clearance of methotrexate, enhancing the toxicity of methotrexate. The effect of ketorolac tromethamine on methotrexate clearance has not been studied.

Nondepolarizing muscle relaxants: In postmarketing experience there have been reports of possible interaction between ketorolac tromethamine injection and *nondepolarizing muscle relaxants* that resulted in apnea. The concurrent use of ketorolac tromethamine with muscle relaxants has not been formally studied.

ACE inhibitors: Concomitant use of *ACE inhibitors* may increase the risk of renal impairment, particularly in volume-depleted patients.

Antiepileptic drugs: Sporadic cases of seizures have been reported during concomitant use of ketorolac tromethamine and *antiepileptic drugs* (phenytoin, carbamazepine).

Psychoactive drugs: Hallucinations have been reported when ketorolac tromethamine was used in patients taking *psychoactive drugs* (fluoxetine, thiothixene, alprazolam).

Morphine: Ketorolac tromethamine injection has been administered concurrently with *morphine* in several clinical trials of postoperative pain without evidence of adverse interactions. Do not mix ketorolac tromethamine and morphine in the same syringe. There is no evidence in animal or human studies that ketorolac tromethamine induces or inhibits hepatic enzymes capable of metabolizing itself or other drugs.

Carcinogenesis, Mutagenesis and Impairment of Fertility

An 18-month study in mice with oral doses of ketorolac tromethamine at 2 mg/kg/day (0.9 times the human systemic exposure at the recommended IM or IV dose of 30 mg qid, based on area-under-the-plasmaconcentration curve [AUC]), and a 24-month study in rats at 5 mg/kg/day (0.5 times the human AUC) showed no evidence of tumorigenicity.

Ketorolac tromethamine was not mutagenic in the Ames test, unscheduled DNA synthesis and repair, and in forward mutation assays. Ketorolac tromethamine did not cause chromosome breakage in the *in vivo* mouse micronucleus assay. At 1590 mcg/mL and at higher concentrations, ketorolac tromethamine increased the incidence of chromosomal aberrations in Chinese hamster ovarian cells.

Impairment of fertility did not occur in male or female rats at oral doses of 9 mg/kg (0.9 times the human AUC) and 16 mg/kg (1.6 times the human AUC) of ketorolac tromethamine, respectively.

Pregnancy

Pregnancy Category C. Reproduction studies have been performed during organogenesis using daily oral doses of ketorolac tromethamine at 3.6 mg/kg (0.37 times the human AUC) in rabbits and at 10 mg/kg (1.0 times the human AUC) in rats. Results of these studies did not reveal evidence of teratogenicity to the fetus. Oral doses of ketorolac tromethamine at 1.5 mg/kg (0.14 times the human AUC), administered after gestation Day 17, caused dystocia and higher pup mortality in rats. There are no adequate and well-controlled studies of ketorolac tromethamine in pregnant women. Ketorolac tromethamine should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Labor and Delivery

The use of ketorolac tromethamine is contraindicated in labor and delivery because, through its prostaglandin synthesis inhibitory effect, it may adversely affect fetal circulation and inhibit uterine contractions, thus increasing the risk of uterine hemorrhage (see **CONTRAINDICATIONS**).

Lactation and Nursing

After a single administration of a 10 mg ketorolac tromethamine tablet to humans, the maximum milk concentration observed was 7.3 ng/mL, and the maximum milk-to-plasma ratio was 0.037. After 1 day of dosing (qid), the maximum milk concentration was 7.9 ng/mL, and the maximum milk-to-plasma ratio was 0.025. Because of the possible adverse effects of prostaglandin-inhibiting drugs on neonates, use in nursing mothers is contraindicated.

Pediatric Use

The safety and effectiveness of single doses of ketorolac tromethamine injection have been established in pediatric patients between the ages of 2 and 16 years. Ketorolac tromethamine injection has been shown to be effective in the management of moderately severe acute pain that requires analgesia at the opioid level, usually in a postoperative setting. Safety and efficacy in pediatric patients below the age of 2 have not been established. Therefore, ketorolac tromethamine injection is not recommended in pediatric patients below the age of 2. The risk of bleeding was greater in those patients administered ketorolac tromethamine injection following tonsillectomy. Physicians should consider the increased risk of bleeding before deciding to administer ketorolac tromethamine injection in patients following tonsillectomy (see WARNINGS: *Hemorrhage*, *Pediatrics and Tonsillectomy*).

The risks identified in the adult population with ketorolac tromethamine injection use also apply to pediatric patients. Therefore,

consult the **CONTRAINDICATIONS**, **WARNINGS**, **PRECAUTIONS**, and **ADVERSE REACTIONS** sections when prescribing ketorolac tromethamine injection to pediatric patients.

Geriatric Use

(≥65 Years of Age)

Because ketorolac tromethamine may be cleared more slowly by the elderly (see **CLINICAL PHARMACOLOGY**) who are also more sensitive to the adverse effects of **NSAIDs** (see **WARNINGS**: **Renal Effects**), extra caution and reduced dosages (see **DOSAGE AND ADMINISTRATION**) must be used when treating the elderly with ketorolac tromethamine injection. The lower end of the ketorolac tromethamine injection dosage range is recommended for patients over 65 years of age, and total daily dose is not to exceed 60 mg. The incidence and severity of gastrointestinal complications increases with increasing dose of, and duration of treatment with, ketorolac tromethamine.

ADVERSE REACTIONS

Adverse reaction rates increase with higher doses of ketorolac tromethamine. Practitioners should be alert for the severe complications of treatment with ketorolac tromethamine, such as GI ulceration, bleeding and perforation, postoperative bleeding, acute renal failure, anaphylactic and anaphylactoid reactions and liver failure (see **Boxed WARNING, WARNINGS, PRECAUTIONS**, and **DOSAGE AND ADMINISTRATION**). These NSAID-related complications can be serious in certain patients for whom ketorolac tromethamine is indicated, especially when the drug is used inappropriately.

The Adverse Reactions Listed Below Were Reported In Clinical Trials As Probably Related To Ketorolac Tromethamine:

Incidence Greater than 1%

Percentage of incidence in parentheses for those events reported in 3% or more patients.

Body as a Whole: edema (4%) Cardiovascular: hypertension Dermatologic: pruritus, rash

Gastrointestinal: nausea (12%), dyspepsia (12%), gastrointestinal pain (13%), diarrhea (7%), constipation, flatulence, gastrointestinal

fullness, vomiting, stomatitis **Hemic and Lymphatic:** purpura

Nervous System: headache (17%), drowsiness(6%), dizziness (7%), sweating **Injection-site pain** was reported by 2% of patients in multidose studies.

Incidence 1% or Less

Body as a Whole: weight gain, fever, infections, asthenia

Cardiovascular: palpitation, pallor, syncope

Dermatologic: urticaria

Gastrointestinal: gastritis, rectal bleeding, eructation, anorexia, increased appetite

Hemic and Lymphatic: epistaxis, anemia, eosinophilia

Nervous System: tremors, abnormal dreams, hallucinations, euphoria, extrapyramidal symptoms, vertigo, paresthesia, depression,

insomnia, nervousness, excessive thirst, dry mouth, abnormal thinking, inability to concentrate, hyperkinesis, stupor

Respiratory: dyspnea, pulmonary edema, rhinitis, cough

Special Senses: abnormal taste, abnormal vision, blurred vision, tinnitus, hearing loss

Urogenital: hematuria, proteinuria, oliguria, urinary retention, polyuria, increased urinary frequency

The Following Adverse Events Were Reported From Postmarketing Experience:

Body as a Whole: hypersensitivity reactions such as anaphylaxis, anaphylactoid reaction, laryngeal edema, tongue edema (see Boxed

WARNING, WARNINGS), angioedema, myalgia

Cardiovascular: hypotension, flushing

Dermatologic: Lyell's syndrome, Stevens-Johnson syndrome, exfoliative dermatitis, maculopapular rash, urticaria

Gastrointestinal: peptic ulceration, GI hemorrhage, GI perforation (see Boxed WARNING, WARNINGS), melena, acute

pancreatitis, hematemesis, esophaghitis

Hemic and Lymphatic: postoperative wound hemorrhage (rarely requiring blood transfusion-see Boxed WARNING, WARNINGS,

and **PRECAUTIONS**), thrombocytopenia, leukopenia **Hepatic:** hepatitis, liver failure, cholestatic jaundice

Nervous System: convulsions, psychosis, aseptic meningitis

Respiratory: asthma, bronchospasm

Urogenital: acute renal failure (see Boxed WARNING, WARNINGS), flank pain with or without hematuria and/or azotemia,

interstitial nephritis, hyponatremia, hyperkalemia, hemolytic uremic syndrome

OVERDOSAGE

Symptoms following acute NSAIDs overdoses are usually limited to lethargy, drowsiness, nausea, vomiting, and epigastric pain, which are generally reversible with supportive care. Gastrointestinal bleeding can occur. Hypertension, acute renal failure, respiratory depression and coma may occur, but are rare. Anaphylactoid reactions have been reported with therapeutic ingestion of NSAIDs, and may occur following an overdose.

Patients should be managed by symptomatic and supportive care following a NSAIDs overdose. There are no specific antidotes. Emesis and/or activated charcoal (60 g to 100 g in adults, 1g/kg to 2 g/kg in children) and/or osmotic cathartic may be indicated in patients seen within 4 hours of ingestion with symptoms or following a large oral overdose (5 to 10 times the usual dose). In controlled overdosage, daily doses of 360 mg of ketorolac tromethamine injection given for 5 days (three times the highest recommended dose), caused abdominal pain and peptic ulcers which healed after discontinuation of dosing. Metabolic acidosis has been reported following intentional overdosage.

Single overdoses of ketorolac tromethamine have been variously associated with abdominal pain, nausea, vomiting, hyperventilation, peptic ulcers, and/or erosive gastritis and renal dysfunction which have resolved after discontinuation of dosing. Dialysis does not significantly clear ketorolac tromethamine from the blood stream.

DOSAGE AND ADMINISTRATION

IN ADULTS, THE COMBINED DURATION OF USE OF KETOROLAC TROMETHAMINE INJECTION AND KETOROLAC TROMETHAMINE TABLETS IS NOT TO EXCEED 5 DAYS. IN ADULTS, THE USE OF KETOROLAC TROMETHAMINE TABLETS IS ONLY INDICATED AS CONTINUATION THERAPY TO KETOROLAC TROMETHAMINE INJECTION.

Ketorolac Tromethamine Injection

Adult Patients

Ketorolac tromethamine injection may be used as a single or multiple dose on a regular or "prn" schedule for the management of moderately severe acute pain that requires analgesia at the opioid level, usually in a postoperative setting. Hypovolemia should be corrected prior to the administration of ketorolac tromethamine (see WARNINGS: *Renal Effects*). Patients should be switched to alternative analgesics as soon as possible, but ketorolac tromethamine therapy is not to exceed 5 days.

When administering ketorolac tromethamine injection, the IV bolus must be given over no less than 15 seconds. The IM administration should be given slowly and deeply into the muscle. The analgesic effect begins in~30 minutes with maximum effect in 1 to 2 hours after dosing IV or IM. Duration of analgesic effect is usually 4 to 6 hours.

Single-Dose Treatment: The Following Regimen Should Be Limited To Single Administration Use Only

Adult Patients
IM Dosing:

- Patients <65 years of age: One dose of 60 mg.
- Patients ≥65 years of age, renally impaired and/or less than 50 kg (110 lbs) of body weight: One dose of 30 mg.

IV Dosing:

- · Patients <65 years of age: One dose of 30 mg.
- · Patients ≥65 years of age, renally impaired and/or less than 50 kg (110 lbs) of body weight: One dose of 15 mg. <u>Pediatric Patients (2 to 16 years of age)</u>: The pediatric population should receive only a single dose of ketorolac tromethamine injection as follows:

IM Dosing:

· One dose of 1 mg/kg up to a maximum of 30 mg.

IV Dosing:

• One dose of 0.5 mg/kg up to a maximum of 15 mg.

Multiple-Dose Treatment (IV or IM) In Adults:

- Patients <65 years of age: The recommended dose is 30 mg ketorolac tromethamine injection every 6 hours. The maximum daily dose should not exceed 120 mg.
- · For Patients ≥65 years of age, renally impaired patients (see WARNINGS) and patients less than 50 kg (110 lbs): The recommended dose is 15 mg ketorolac tromethamine injection every 6 hours. The maximum daily dose for these populations should not exceed 60 mg.

For breakthrough pain do not increase the dose or the frequency of ketorolac tromethamine. Consideration should be given to supplementing these regimens with low doses of opioids prn unless otherwise contraindicated.

Pharmaceutical Information for Ketorolac Tromethamine Injection

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration whenever solution and container permit.

Ketorolac tromethamine injection should not be mixed in a small volume (eg, in a syringe) with morphine sulfate, meperidine hydrochloride, promethazine hydrochloride or hydroxyzine hydrochloride; this will result in precipitation of ketorolac from solution.

HOW SUPPLIED

FOR IM or IV USE

Product	NDC No.	
No.	No.	
16010	63323-161-01	Ketorolac Tromethamine Injection, USP, 15 mg: 15 mg/mL, 1 ml fill in a 2 mL single dose vial.
160201	63323-162-01	Ketorolac Tromethamine Injection, USP, 30 mg: 30 mg/mL, 1 ml fill in a 2 mL single dose vial.

FOR IM USE ONLY, NOT FOR IV USE

Product No.	NDC No.	
106202		Ketorolac Tromethamine Injection, USP, 60 mg: 30 mg/mL in a 2 mL single dose vial.

The above products are packaged in twenty-fives.

Vial stoppers do not contain natural rubber latex.

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature], with protection from light.



45906E

Revised: April 2008

PACKAGE LABEL - PRINCIPAL DISPLAY - Ketorolac Tromethamine 1 mL Single Dose Vial Label

NDC 63323-161-01

160101

Ketorolac Tromethamine Injection, USP


15 mg/mL

For IM or IV USE

1 mL Single Dose Vial

Usual Dosage: See insert.

Rx only



PACKAGE LABEL - PRINCIPAL DISPLAY - Ketorolac Tromethamine 1 mL Single Dose Vial Tray Label

NDC 63323-161-01

160101

Ketorolac Tromethamine Injection, USP

15 mg/mL

For Intravenous or Intramuscular Use

Rx only

1 mL Single Dose Vial



160201

Ketorolac Tromethamine Injection, USP 30 mg/mL

For IM or IV USE

1 mL Single Dose Vial
Usual Dosage: See insert.

Rx only

NDC 63323-162-01 160201 **KETOROLAC TROMETHAMINE** *INJECTION, USP*

30 mg/mL

1 mL Single Dose Vial Usual Dosage: See insert. Rx only

PACKAGE LABEL - PRINCIPAL DISPLAY - Ketorolac Tromethamine 1 mL Single Dose Vial Tray Label NDC 63323-162-01

160201

Ketorolac Tromethamine Injection, USP 30 mg/mL

For Intravenous or Intramuscular Use

Rx only

1 mL Single Dose Vial

Sterile
EachmLcontains: 30mg
ketorolactromethamine, 10% (w/v)alcotho (USP), 4.35 mg
sod um chloride, 1 mgcitricacid, anhydroxide or hydrochloric acid.
UsualD caage: See insert.
Store at 20° to 25°C (68° to 77°F)
[see USP Controlled Room
Temperature].
PROTECT FROM LIGHT.
Retain in carton until time of use.
Vial stoppers do not contain natural rubber latex.

APP Pharmaceuticals, LLC
Schaumburg, Leon 33

42641 C