DRUG FACTS

Active ingredient(s) ***Purpose***

Use(s)

Uses: temporarily releives mnor aches and pains due to:
headache, backache, muscular aches, arthritis pain, colds, flu, fever, toothache, premenstrual cramps.

Warnings

Warnings:
Liver warning: This product contains acetaminophen. Severe liver damage may occur if: adult or child 12 years and older takes more than 12 tablets in 24 hours, which is the maximum daily amount; takn with other durgs containing acetaminophen; adult has 3 or more alcoholic drinks every day while using this product.

Do not use
Do not use
- wITH ANY OTHER DURG CONTAINING ACETAMINOPHEN(PRESCRIPTION OR NONPRESCRIPTION). IF YOU ARE NOT SURE WHETHER A DRUG CONTAINS ACETAMINOPHEN, ASK A DOCTOR OR PHARMACIST.

Ask a doctor before use if you have
Ask a doctor before use ifyou have liver disease.

Ask a doctor or pharmacist before use if you are
Ask a doctor or pharmacist before use if you are
- taking the blood thinning dur warfarin
- taking sedatives or tranquilizers

When using this product

Stop use and ask a doctor if
Stop use and ask a doctor if
pain prefor more than 10 days; fever persists for more than 3 days (unless directed by a doctor); condition worsens or new symptoms appear; redness or swelling is present. These could be signes of a serious condition.

If pregnant or breast-feeding

Keep out of reach of children
Keep this and all drugs out of reach of children

Directions

Directions
adults and children 12 years and over: take 1 to 2 tablets every six hours while symptoms last. Maximum daily dose is 9 tablets in 24 hours unless directed by a doctor.
children under 12 years: Do not use this product in children under 12 years of age; this will provide more than the recommended dose (overdose) and could cause serious health problems.

Other information

Other information
- store at 25° C (77° F) excursions permitted between 15°-30° C (59°-86° F)
- use by expiration date on package
- avoid excessive heat
- sodium free

Inactive ingredients

colloidal silicon dioxide, crospovidone, fd&C yellow #6 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, mineral oil, povidone, pregelatinized starch, stearic acid