#### GENTAMICIN PIGLET - gentamicin sulfate injection

Agri Laboratories, Ltd.

#### DESCRIPTION

Piglet Injection Veterinary

Antibacterial

For Piglets Up to 3 days of age

Gentamicin Piglet Injection is a sterile preparation for intramuscular administration in piglets up to 3 days of age. Each milliliter contains gentamicin sulfate, USP equivalent to 5 mg gentamicin base, 2 mg sodium bisulfite, 0.1 mg edetate disodium, 7.46 mg sodium acetate, 3 mg glacial acetic acid; 1.3 mg methylparaben and 0.2 mg propylparaben as preservatives, water for injection q.s.

#### WARNING

Piglets injected with Gentamicin Piglet Injection must not be slaughtered for food at least 40 days following treatment. Restricted Drug (California)--use only as directed.

Read accompanying directions carefully.

Store between  $2^{0}$  and  $30^{0}$ C ( $36^{0}$  and  $86^{0}$ F)

Protect from freezing.

For intramuscular use only.

ANADA #: 200-394 Approved by F.D.A.

#### **INDICATIONS**

Gentamicin Piglet Injection is recommended for the treatment of porcine colibacillosis caused by strains of *E.coli* sensitive to gentamicin.

#### DOSAGE AND ADMINISTRATION

The recommended dose is 1 mL (5 mg) administered intramuscularly. Gentamicin Piglet Injection is not to be used more than once in each piglet.

If piglets do not respond to treatment with Gentamicin Piglet Injection, diagnosis should be reevaluated. Clean and sterilize needles and syringes

by boiling in water for 15 minutes prior to use.

Prior to injection, disinfect the injection site and top of the vial with a suitable disinfectant, such as 70% alcohol.

Use all precautions to prevent contamination of vial contents.

#### CONTRAINDICATIONS

There are no known contraindications to this drug when used as directed.

#### **PRECAUTIONS**

Store between 2° and 30°C (36° and 86°F)

#### **CHEMISTRY**

Gentamicin is a mixture of aminoglycoside antibiotics derived from the fermentation of *Micromonospora purpurea*. Gentamicin sulfate, USP

is a mixture of sulfate salts of the antibiotics produced in this fermentation. The salts are weakly acidic and freely soluble in water. Gentamicin sulfate, USP is stable in solution for the period of time indicated on the labels by the expiration date.

Gentamicin sulfate, USP contains not less than 500 micrograms of gentamicin base per milligram.

#### HOW SUPPLIED

Gentamicin Piglet Injection, 5 mg/mL is available in 100 mL, 250 mL and 500 mL multiple dose vials.

**Antibacterial** 

100 Doses

Rev. 09-10

Manufactured for:

Agri Laboratories, Ltd.

**St. Joseph, MO 64503** 

# For use in piglets only, up to 3 days of age.

**Each mL contains:** gentamicin sulfate, USP equivalent to 5 mg gentamicin base, 2 mg sodium bisulfite, 0.1 mg edetate disodium, 7.46 mg sodium acetate, 3 mg glacial acetic acid; 1.3 mg methylparaben and 0.2 mg propylparaben as preservatives, water for injection q.s.

Indications: Gentamicin Piglet Injection is recommended for the treatment of porcine colibacillosis caused by strains

of *E. coli* sensitive to gentamicin.







Registered trademark of Agri Laboratories, Ltd.

LOT

**EXP** 

NDC 57561-638-05

# GENTAMICIN PIGLET INJECTION

(GENTAMICIN SULFATE, USP)

Piglet Injection
Antibacterial
250 Doses
5mg/mL

ANADA#: 200-394 Approved by F.D.A.

NET CONTENTS: 8.5 oz (250 mL)





**Dose:** 1 mL per pig cularly. Gentamicin I used more than once

WARNING: Pig Gentamicin Pigl be slaughtered 40 days followin

Restricted Drug as directed.

Read accompanyi

Store between 86'F)

Protect from fre

ANADA#: 200-39

Manuf Agri Lab St. Josep

G-6639-05

**Dose:** 1 mL per piglet, administered intramuscularly. Gentamicin Piglet Injection is not to be used more than once in each piglet.

WARNING: Piglets injected with Gentamicin Piglet Injection must not be slaughtered for food for at least 40 days following treatment.

Restricted Drug (California)--use only as directed.

Read accompanying directions carefully.

Store between 2° and 30°C (36° and 86°F)

Protect from freezing.

For intramuscular use only.

ANADA#: 200-394 Approved by F.D.A.

Manufactured for: Agri Laboratories, Ltd. St. Joseph, MO 64503

G-6639-05 Rev. 09-10

Ge Pigle

(GENTAI)

Piglet In

Piglet I

For Piglet

## **DESCRIPTI**

Injection is intramuscular to 3 days of gentamicin sul gentamicin ba 0.1 mg edetate acetate, 3 mg methylparaben as preservative

**WARNING:** For use in piglets up to 3 days of age only. Piglets injected with Gentamicin Piglet Injection must not be slaughtered for food for at least forty (40) days following treatment. Use of more than the recommended dose will result in gentamicin tissue residues beyond forty (40) days.

5

**INDICATIONS:** Gentamicin Piglet Injection is recommended for the treatment of porcine colibacillosis caused by strains of *E. coli* sensitive to gentamicin.

### **DOSAGE AND ADMINISTRATION:**

The recommended dose is 1 mL (5 mg) administered intramuscularly. Gentamicin Piglet Injection is not to be used more than once in each piglet. If piglets do not respond to treatment with Gentamicin Piglet Injection, diagnosis should be reevaluated. Clean and sterilize needles

2

and syringes by boiling in water for 15 minutes prior to use. Prior to injection, disinfect the injection site and top of the vial with a suitable disinfectant, such as 70% alcohol.

Use all precautions to prevent contamination of vial contents.

**CONTRAINDICATIONS:** There are no known contraindications to this drug when used as directed.

**PRECAUTIONS:** Store between 2° and 30°C (36° and 86°F)

**CHEMISTRY:** Gentamicin is a mixture of aminoglycoside antibiotics derived from the fermentation of *Micromonospora purpurea*. Gentamicin sulfate, USP is a mixture of sulfate salts of the antibiotics produced in this fermentation. The salts are weakly acidic and freely soluble in

water. Gentam in solution for on the labels b Gentamicin su than 500 micr per milligram.

HOW SUPI Injection, 5r 100 mL, 250 dose vials.

3