MEDIC BOOSTER DIP - iodine liquid
Animal Medic Incorporated

DRUG FACTS

ANIMAL MED

Medic Booster® D

Sanitizing Teat Dip

An Aid in Reducing the Spread of Organisms Which May Cause Mastitis

ACTIVE INGREDIENT:

Iodine.....0.50%

Contains nonylphenoxypolyethoxyethanol-iodine complex, providing 5000 ppm min. titratable iodine.

(Contains Aloe and other special emollients.)

FOR COMMERCIAL USE ONLY
KEEP OUT OF REACH OF CHILDREN

CAUTION

Not for internal use. May cause eye irritation. If swallowed, may cause irritation, nausea, stomach distress.

PRECAUTIONARY STATEMENTS

No protective equipment is necessary under normal use conditions. If splashing may occur, eye protection is recommended.

FIRST AID

If in Eyes: Immediately flush ey
running water. Remove contact
flushing for at least 15 minutes
If Swallowed: Rinse mouth; th
glasses of water. DO NOT induc
give anything by mouth to an u

IF IRRITATION OR DISCOMFOR
CALL A PHYSICIAN.

FOR EMERGENCY MEDICAL IN
OR CANADA, INFOTRAC (800)

Rev. 3/2004

Lot #

Expiration Date

Keep Out of Reach of Children
FOR COMMERCIAL USE ONLY
CAUTION
Not for internal use. May cause eye irritation.
If swallowed, may cause irritation, nausea, stomach distress.
PRECAUTIONARY STATEMENTS
No protective equipment is necessary under normal use conditions.
If splashing may occur, eye protection is recommended.
FIRST AID
If in Eyes: Immediately flush eyes with plenty of cool, running water. Remove contact lenses; then continue flushing for at least 15 minutes, holding eyelids apart.
If Swallowed: Rinse mouth; then drink 1 or 2 glasses of water. DO NOT induce vomiting. Never give anything by mouth to an unconscious person. IF IRRITATION OR DISCOMFORT PERSISTS, CALL A PHYSICIAN.
FOR EMERGENCY MEDICAL INFORMATION IN USA OR CANADA, INFOTRAC (800) 535-5053.
Saniting Teat Dip
An aid in reducing the spread of organisms that might cause mastitis.
Active Ingredient
Iodine 0.50%
Inactive Ingredients
Surfactants, glycerine, citric acid, aloe, water..... 99.50%
DIRECTIONS FOR USE
IMPORTANT: Do not mix *Medic Booster Dip* with water, any other teat dips, or other products unless specified in product use instructions. If transferred from this container to any other, make sure that the other container is clean and bears the proper container label for *Medic Booster Dip.* If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product.
Do not reuse or return to original container.
Before Milking: Wash teats thoroughly just prior to next milking with appropriate udder wash solution or pre-milking teat dip to avoid contamination of milk. Teats should be dried with single-service towels. Use proper procedures for udder washing or pre-dipping.
Pre-Milk Dipping: Fill teat cup with *Medic Booster Dip*. Do not dilute. Before each cow is milked, dip each teat full-length into the teat cup containing *Medic Booster Dip*. Wipe teats dry after dipping using single-service towels to avoid contamination of milk.
Post-Milk Dipping: Immediately after each milking, use *Medic Booster Dip* at full strength. Dip each teat full-length into the teat cup containing *Medic Booster Dip*. Allow to air dry. Do not wipe. Always use fresh, full-strength *Medic Booster Dip*. Do not turn cows out in freezing weather until *Medic Booster Dip* is completely dry.
Expanded Usage: When freshening cows, begin dipping teats twice daily for about ten days before calving.