LORATADINE D - loratadine and pseudoephedrine sulfate tablet, extended release

Bryant Ranch Prepack

ACTIVE INGREDIENTS (IN EACH TABLET)

Loratadine 5 mg

Pseudoephedrine sulfate 120 mg

PURPOSE

Antihistamine

Nasal decongestant

USES

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- sneezing
- · itchy, watery eyes
- runny nose
- · itching of the nose or throat
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- reduces swelling of nasal passages
- · temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

WARNINGS

Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- · heart disease
- · thyroid disease
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- liver or kidney disease. Your doctor should determine if you need a different dose.

When using this product do not take more than directed.

Taking more than directed may cause drowsiness.

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- · nervousness, dizziness or sleeplessness occurs

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

DIRECTIONS

• do not divide, crush, chew or dissolve the tablet

adults and children 12 years and over	1 tablet every 12 hours; not more than 2 tablets in 24 hours
children under 12 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

OTHER INFORMATION

- each tablet contains: calcium 25 mg
- · do not use if blister unit is broken or torn
- store between 20° to 25°C (68° to 77°F)
- · keep in a dry place

INACTIVE INGREDIENTS

croscarmellose sodium, dibasic calcium phosphate, hypromellose, lactose monohydrate, magnesium stearate, pharmaceutical ink, povidone, titanium dioxide

QUESTIONS OR COMMENTS?

1-800-719-9260

Pseudoephedrine/Loratadine 120mg/5mg

